

2023-06

Terrestrial carbon isotope stratigraphy of the EoceneOligocene transition, Petrockstowe and Bovey basins, Devon, UK

Chaanda, MS

<https://pearl.plymouth.ac.uk/handle/10026.1/21172>

10.1016/j.pgeola.2023.05.003

Proceedings of the Geologists' Association

Elsevier BV

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

Terrestrial carbon isotope stratigraphy of the Eocene–Oligocene transition, Petrockstowe and Bovey basins, Devon, UK

Mohammed S. Chaanda^a, Stephen T. Grimes^a, Rhodri M. Jerrett^b, Mark Anderson^a, Melanie J. Leng^c
Meriel E. Fitzpatrick^a, Gregory D. Price^{a*}

^aSchool of Geography, Earth and Environmental Sciences, Plymouth University, Drake Circus,
Plymouth, PL4 8AA, UK

^bDepartment of Earth and Environmental Sciences, University of Manchester, Oxford Road,
Manchester, M13 9PL, UK

^cBritish Geological Survey, Keyworth, Nottingham, NG12 5GG, UK.

ABSTRACT

The terrestrial sediments of the Petrockstowe and Bovey basins in Devon, UK were examined. Their age is considered to be Eocene and Oligocene. The sediments (kaolinitic clays, silts, sands, gravels, and lignites) from both basins were analysed for carbon isotopes of organic material, in conjunction with total organic carbon and palynological analyses used to unravel the type of and provenance of organic matter present. Within the Petrockstowe Basin, the lowermost interval examined shows a palynological distribution dominated by phytoclasts, whilst the upper part of the core is dominated by higher concentrations of palynomorphs (up to 90%) and an increase in amorphous organic matter consistent (up to 37%) with a change from sand-filled fluvial channels followed by an ephemeral lake or lake margin setting. Our palynological data from South John Acres Lane Quarry section, Bovey Basin, show that within the lignites palynomorphs are high again (up to 95%) consistent with them representing more ephemeral lakes or lake margins periodically exposed with mires. Our palynological data set further allows us to determine that isotope trends are not overly determined by the source of carbon in the basins. Our study suggests that the observed patterns were primarily produced by variations of the isotope ratios of terrestrial atmospheric carbon reservoirs. Even with our less than well constrained biostratigraphic control, the data indicate that the carbon isotope excursions seen in the Eocene and Oligocene could be associated with several transient carbon isotopic shifts (associated with the Palaeocene-Eocene Thermal Maximum). Our

29 findings therefore appear to lend support to the surface ocean and atmosphere behaving as
30 coupled reservoirs at this time.

31

32 Keywords: Eocene Oligocene Petrockstowe Bovey terrestrial carbon isotope, Palynological analyses

33

34 *Corresponding author.

35 E-mail address: g.price@plymouth.ac.uk (Gregory D. Price)

36

37

1. Introduction

The terrestrial sediments of the Petrockstowe and Bovey basins in Devon, UK offer an opportunity to examine the carbon cycle during the Eocene–Oligocene transition (~33.9 million years ago), an interval, that saw a climate shift from a largely ice-free greenhouse conditions to an icehouse world (Miller et al., 2009; Coxall and Wilson, 2011; Hutchinson et al. 2021). At this time major changes in fauna and flora record a shift toward more cold-climate-adapted species (e.g., Sun et al., 2014). For this pivotal interval in Earth's climate, our understanding the role of the carbon cycle, is much more limited (Coxall and Wilson, 2011; Armstrong McKay et al. 2016). Across the Eocene–Oligocene boundary a positive carbon isotope($\delta^{13}\text{C}$) excursion of ~1.0‰ is typically recorded in the marine record followed by a decline to pre-excursion values (~0.7‰) in the Oligocene (e.g., Nilsen et al. 2003; Armstrong McKay et al. 2016). This well-documented perturbation may be used to correlate marine and terrestrial sections from around the globe as previous studies have shown that $\delta^{13}\text{C}$ obtained from terrestrial organic material such as wood and lignites typically records a global signal (e.g., Heimhofer et al., 2003; Collinson et al. 2003; Gröcke et al. 2005; Bechtel et al. 2009; Hodgson et al. 2011; Jerrett et al. 2015; Lenz et al. 2022).

The Petrockstowe and Bovey basins lie on the Sticklepath - Lustleigh Fault and owe their origin to subsidence within this zone (Blyth 1962, Dearman 1963). About 600 m of Eocene–Oligocene sediments are present in the Petrockstowe Basin (Freshney et al. 1979) and ~ 1200 m in the Bovey Basin (Edwards 1976). In this study, we present new organic $\delta^{13}\text{C}$ data from terrestrial Eocene–Oligocene aged sediments from these basins. Our data is used to improve age constraints on the succession via comparison of our terrestrial $\delta^{13}\text{C}$ record with that of the extensively described time-equivalent marine sections. The similarity and magnitude of the $\delta^{13}\text{C}$ excursions between terrestrial and marine records can also be used to assess whether these archives behaved as coupled reservoirs during this time.

2. Location and Geological setting

The Petrockstowe Basin, nr Newton Abbott, Devon, UK (Fig. 1), lies on the Sticklepath – Lustleigh Fault (Dearman 1963; Holloway and Chadwick, 1986). The fault became active during the Paleogene and most activity ceased before deposition of the upper part of the Bovey Formation (Blyth 1962). Bristow and Robson (1994) proposed a structural model for the development of the basin – a pull-push model – and suggested that the development was in two phases: an early,

69 transitional phase, during which much of the sedimentation occurred, and a subsequent
 70 transpressional phase in which boundary thrust faults developed. Geophysical measurements,
 71 confirmed by a British Geological Survey (BGS) borehole from the centre of the basin proved a basin
 72 fill of 660 m of sands and silts (Freshney et al. 1979). The sediments, kaolinitic clays, silts, sands,
 73 gravels, and lignite, were likely to have been derived from weathering granite under warm
 74 temperate or sub-tropical conditions of the early Paleogene (Bristow 1968; Edwards 1976). The
 75 succession consists of fining-upwards cycles comprising of one or more of a gravel lag, gravelly
 76 sands, silty sands, and are probably representative of point bar and swale-fill deposits of a river
 77 system. These interstratify with clays and silts of lacustrine origin (Freshney 1970; Freshney et al.
 78 1979). Based on palynological evidence, Turner (cited by Freshney et al. 1979) suggested these
 79 were deposited in a subtropical climate with palms, ferns and heathers and many plants with
 80 swamp affinities.

81 The Bovey Basin is located between Newton Abbot, Kingsteignton and Bovey Tracey, Devon
 82 UK (Fig. 2) and is 45 km southeast of the Petrockstowe Basin. The Bovey Basin lies southeast of the
 83 Dartmoor granite and is approximately 7 km from east to west and 5 km from north to south. In the
 84 northern and eastern boundaries of the basin there are sedimentary contacts between the
 85 Dartmoor granite and the Upper Greensand and Aller Gravel. The bulk of the basin is filled by a
 86 thick (~ 1200 m) succession of Paleogene kaolinitic clays, silty clays, silts, lignites and sands,
 87 referred to as the Bovey Formation (Edwards 1976). Edwards (1976) proposed a morphological sub-
 88 division of the basin into two parts, lying to the north and to the south of Newton Abbot. The part
 89 between Bovey Tracey and Newton Abbot is considered as the main basin; the second part lies
 90 south of Newton Abbot and is referred to as Decoy Basin (Fig. 2). Edwards and Freshney (1982)
 91 proposed an informal sub-division of the Bovey Formation into 'lower', 'middle' and 'upper'. The
 92 'lower' is not exposed and the 'middle' and 'upper' Bovey Formation includes 14 members, some of
 93 uncertain stratigraphic position or lateral equivalents and are described in detail by Edwards and
 94 Freshney (1982) and Selwood et al. (1984). Of the top 350 exposed at the surface (Edwards 1976),
 95 48 m of the Abbots Clay and Sand and Southacre Clay and Lignite members of the 'middle' Bovey
 96 Formation were examined in the exposed working section at the South John Acre Lane Quarry.
 97 Chandler (1957) and Edwards (1976), suggest that during the Oligocene the lignites accumulated in
 98 swamps with associated fluvial sands and plant debris swept in from a warm hinterland into a lake

basin lying on Palaeozoic strata (see also Selwood et al. 1984). The lake was surrounded by marshland tree covered slopes (Chandler 1957).

3. Age of the Petrockstowe and Bovey basins

The age of the Bovey lignites has for a long time been debated (Chandler 1964). Based upon the macroflora in the lignite beds of the Bovey Formation, these were originally regarded as Miocene, but later assigned to the Oligocene (Chandler 1957, 1964). Likewise, Wilkinson (1979) cited by Selwood et al. (1984) noted that pollen from a borehole, near Heathfield that penetrated 185 m of Blatchford Sand (upper Bovey Formation), 69 m of South Acre Clay and Lignite and 51 m of Abbroom Clay and Sand and from below 290 m depth, Eocene indicators like *Anacolosidites* and *Pompeckjoidaepollenites* were observed (Fig. 3). It is also important to note that the Blatchford Sand is an obsolete unit name and has been replaced by the Woolley Grit Member. The South Acre Clay and Lignite Member is therefore likely to be early to middle Oligocene in age (Selwood et al. 1984) and the Abbroom Clay and Sand Member would contain the Eocene–Oligocene boundary. Freshney et al. (1982) also suggested that the lowermost ~ 700–800 m of the Bovey Formation could probably be assigned to Eocene (see also Wilkinson et al. 1980; Wilkinson and Boulter, 1981). For the Petrockstowe Basin, Turner (cited by Freshney et al. 1979) reported that pollen data indicate a boundary between the Oligocene and Eocene at ~120 m depth in the BGS Borehole No.1 (Fig. 3).

4. Methods

From Petrockstowe 2 borehole cores (Petrockstowe 1A and 1B) held in the core repository at the BGS, Keyworth, Nottingham, UK were logged, and sub sampled. The sampled section was 640 m long, and samples were collected, on average, every 6 m. Within the region of the Eocene Oligocene boundary as proposed by Turner (cited by Freshney et al. 1979), as well as the early Eocene of Core 1B higher resolution sampling was undertaken. In the Bovey Basin, the Abbroom Clay and Sand and Southacre Clay and Lignite members of the Bovey Formation from the accessible exposed working section at South John Acre Lane Quarry (Grid Reference SX 858758) were sampled. The sampled section was 48 m and samples were collected, on average, every 0.6 m. It was necessary to excavate the sediment surface by up to 0.5 m before sampling with a trowel to ensure fresh samples. All sediment types were sampled.

For the determination of the carbon isotope composition of total organic carbon ($\delta^{13}\text{C}_{\text{TOC}}$), samples were ground to a fine powder using an agate pestle and mortar. Powdered samples were decarbonated by placing each sample in a 50 ml polypropylene centrifuge tube and treating with 10% HCl for 1 h until any carbonate had reacted. Samples were then rinsed with deionized water, centrifuged, and rinsed again until neutrality was reached (using universal indicator paper). For $\delta^{13}\text{C}_{\text{TOC}}$ analysis, samples were weighed, to achieve ~ 0.5 mg TOC, into a tin capsule and placed into a Carlo Erba 1500 EA for analysis using an online VG Triple Trap Mass Spectrometer. The $\delta^{13}\text{C}_{\text{TOC}}$ results were calibrated against Vienna PeeDee Belemnite (V-PDB) through laboratory (BROC1) and International Standards (NBS19, NBS22, CH6). Standards were evenly distributed throughout the individual isotope runs to correct for daily drift. The mean standard deviation on replicate $\delta^{13}\text{C}_{\text{TOC}}$ analyses of laboratory standard (BROC1) and soil (SOILB) was between $\pm 0.1\text{‰}$ and 0.5‰ (1 Standard Deviation, σ) for $\delta^{13}\text{C}_{\text{TOC}}$. Replicate analyses showed an average precision of $\pm 0.1\%$. TOC content for each sample was measured using a Carlo Erba 1500 elemental analyser with acetanilide used as the calibration standard.

Palynological analyses were used to unravel the type of organic matter associated with the sediments and as a means of determining the source of carbon reservoir in the basins. Samples were processed using standard palynological processing (Brown, 2008) (hydrochloric acid followed by hydrofluoric acid for demineralisation). Slides were studied using a Zeiss standard microscope, normally using standard transmitted light. This is the first time such a method was used in both Petrockstowe and Bovey basins. To achieve this, counts of >300 organic matter types from each sample was made. There are several schemes to classify different components of the particulate organic matter (e.g., Tyson, 1995, Aggarwal et al., 2019). Four main categories of palynological matter were identified in this study (Fig. 4): (1) Non-opaque phytoclasts includes woody remains, tracheid material, poorly lignified, tissue fragments derived from higher plants, yellowish-brown organic remains (2) Opaque phytoclast includes palynodebris with irregular shapes and charcoal (3) Palynomorphs in this study include pollen, spores and undifferentiated forms (4) amorphous organic matter (AOM) and other palynodebris which appears grey, pale yellow or brown in colour, partly translucent masses of variable thickness and with no cellular detail. The AOM group probably originates from bacteria, phytoplankton and degraded organic aggregates. Their size varies from <5 to about $45\text{ }\mu\text{m}$ in diameter.

5. Results

5.1 Total Organic Carbon (%TOC)

The sediments from the Petrockstowe core have highly varying wt. % TOC values ranging from 0.02 to 42.7 wt. % TOC (Fig. 5). Unsurprisingly, the highest %TOC values coincide with the lignitic clays and lignites. These lignitic clays and lignites are seen in the middle and upper parts of the core (Core 1A). The lower part (core 1B) consists mostly of gravels, sands, and clays with very low wt. % TOC contents. The wt. % TOC values from the South John Acres Lane Quarry section, Bovey Basin, range from 0.1 to 61.8 % (Fig. 6). As for the Petrockstowe Basin, the highest wt. % TOC values coincide with either lignitic clays or the lignites. These sediments are seen within the Southacre Clay and Lignite Member whereas the underlying Abbrook Clay and Sand Member is dominated by sands and silty clays with fewer lignitic clay beds.

5.2 Palynology

Within the cores of the Petrockstowe Basin, the lowermost interval shows a palynological distribution dominated by phytoclasts with at certain intervals nearly 100% and low palynomorphs and AOM. The upper part of the core is dominated by much higher concentrations of palynomorphs and an increase in AOM (up to 37%) and low concentrations of phytoclasts.

With respect to South John Acres Lane Quarry section, Bovey Basin, phytoclasts are highest in the Abbrook Clay and Sand Member. When phytoclasts are high (opaque phytoclasts reach 91%), the palynomorphs shows lowest concentrations and vice versa. High AOM concentrations (up to 66%) are seen at the base of Abbrook Clay and Sand Member and decline upwards. In the overlying Southacre Clay and Lignite Member, dominated by lignites, opaque phytoclast concentrations are low, palynomorphs consistently high (spore-pollen and non-opaque phytoclasts reach 95%, Fig. 6).

5.3 Carbon Isotopes ($\delta^{13}C_{TOC}$)

The $\delta^{13}C_{TOC}$ values of samples in the Petrockstowe cores range from -28.5‰ to -23.5‰ with a mean value of -26.5‰ . As can be seen in Figure 5, values at the base of core 1B begins with a $\delta^{13}C_{TOC}$ value of $\sim -26.1\text{‰}$ at 645 m. A carbon isotope excursion with a magnitude of $\sim 2.5\text{‰}$ can be seen at ~ 586 m depth with $\delta^{13}C_{TOC}$ values reaching a minimum of -28.6‰ . The entire excursion occurs over a thickness of ~ 19 m, from 586 – 605 m. The data then shows a return to more positive values of -26.2‰ at 585 m. Thereafter, the $\delta^{13}C_{TOC}$ values remain relatively consistent between 584 m and 540 m with $\delta^{13}C_{TOC}$ in the range of -27.0‰ and -26.3‰ . There is a lack of core (because of

poor recovery) between 513.59 m to 431.60 m. In the upper part of the Petrockstowe 1A core, the $\delta^{13}\text{C}_{\text{TOC}}$ values generally vary around -26.0‰ . In the uppermost (Oligocene) part of the core the most positive $\delta^{13}\text{C}_{\text{TOC}}$ values are seen.

At South John Acres Lane Quarry section, Bovey Basin the $\delta^{13}\text{C}_{\text{TOC}}$ values range between -27.8‰ to -22.5‰ with a mean value of -26.0‰ (Fig. 6). In this succession, $\delta^{13}\text{C}_{\text{TOC}}$ values show limited variability. In the uppermost (Oligocene) part of the section within the lignitic clays and lignites the most positive $\delta^{13}\text{C}_{\text{TOC}}$ values (-22.5‰) are found.

6. Discussion

6.1 Palynological Interpretation

In the Petrockstowe 1A and 1B cores, close to the base of the succession the high opaque phytoclast content, together with low TOC values (down to 0.1 wt. %), and low AOM and non-opaque phytoclast contents may be related to local oxidation of organic matter (Figs. 5, 7) or diagenesis. Opaque phytoclasts are typically derived from the oxidation of structured organic matters (translucent brown wood, tracheids, cuticle, etc.) and along with a low proportion of the other organic matter types has been documented in oxic swamps and river sediments (e.g., Martín-Closas et al., 2005; Pieńkowski and Waksmundzka, 2009). In the upper part of the Petrockstowe section (Fig. 5) palynomorphs dominate which could indicate suboxic/anoxic waters (Tyson, 1995). Consequently, a restriction of water circulation rather than productivity, may serve as the controlling factor for the organic rich sediment accumulation. Also, the fluctuating, but relatively high percentages of palynomorphs and AOM (up to 37%), could suggest diverse source of areas of the organic matter (e.g., Martín-Closas et al. 2005) with deposition within an ephemeral lake or lake margin. These observations agree with the Petrockstowe 1A and 1B cores, representing a succession of sand-filled fluvial channels followed by an ephemeral lake or lake margin setting (Fig. 5, see Freshney et al., 1979). This represents an overall deepening-up sequence.

In the Abbbrook Clay and Sand and Southacre Clay and Lignite members, the base sees a diversity in palynological types (Fig. 6) with high percentage of AOM and phytoclasts potentially reflecting enhanced preservation in low energy, stagnant, oxygen depleted environment (Tyson 1995, Fig. 7). Only a single high opaque phytoclast level is seen, possibly associated with deposition

within an oxidising environment. In the overlying Southacre Clay and Lignite Member, dominated by lignites, palynomorph concentrations are consistently high and as such suggests a depositional environment associated with a swamp or ephemeral lake or marginal lake. These observations are in agreement that the Abbroom Clay and Sand and Southacre Clay and Lignite members of the Bovey Formation, represents a long-lived lake followed by sediments representing more ephemeral lakes or lake margins periodically exposed with mires (see Chandler, 1964; Edwards, 1976).

Our palynological data set further allows us to determine whether isotope trends are carbon source related. For example, within the Abbroom Clay and Sand Member, when phytoclasts are high, the palynomorphs shows lowest concentrations and vice versa. High, but variable, AOM concentrations are also seen. No correspondence is seen here with fluctuations in carbon isotopes, suggesting organic matter associated with the sediments is not overly determining the source of the carbon reservoir in the basins. Nevertheless, changes in the dominance of gymnosperms, angiosperms or pteridophytes/bryophytes within the vegetation could be of importance.

6.2 Carbon isotope trends

Carbon isotopic ratios from terrestrial organic materials have been previously used to study global carbon-isotope excursions in the Cenozoic (Collinson et al. 2003, Bechtel et al. 2009; Holdgate et al. 2009; Hodgson et al. 2011; Fang et al., 2013; Jerrett et al. 2015; Garel et al. 2020; Lenz et al. 2022). These studies (which use discrete plant fragments, lignites or disseminated organic matter) identify reproducible patterns in atmospheric carbon isotopic compositions. There are just a few terrestrially sourced high-resolution carbon isotope stratigraphies to compare our Eocene and Oligocene data to (e.g., Holdgate et al. 2009; Garel et al. 2020). Nevertheless, our $\delta^{13}\text{C}_{\text{TOC}}$ data are, consistent with terrestrially sourced $\delta^{13}\text{C}$ values of the Eocene (e.g., Collinson et al. 2003, Bechtel et al. 2009; Hodgson et al. 2011). Considering the carbon isotope excursion of -2.5‰ from the lower part of Petrockstowe core 1B (Fig. 5), the magnitude of this excursion falls within the lower limit of that is associated with the Palaeocene-Eocene Thermal Maximum (PETM), which ranges from -2.4 to -6.3‰ (see summary of McInerney and Wing 2011). This suggests it could be related to this event. However, biostratigraphically there is limited data from the Petrockstowe core (see Turner cited by Freshney et al. 1979). The biostratigraphic constraints allow the carbon isotope excursion to also be associated with one of the other transient carbon isotopic shifts that occurred after the Palaeocene-Eocene Thermal Maximum i.e., the Eocene Thermal Maximum (ETM-2). For example, the magnitude of the ETM-2 carbon isotope excursion documented in the

249 continental succession of the McCullough peaks, Bighorn Basin, Wyoming, USA, using paleosol
 250 carbonate is of -3.8‰ (Abels et al. 2012).

251 The presence of the Eocene – Oligocene boundary in the Petrockstowe core 1A has been
 252 proposed, based on pollen data, by Turner (cited by Freshney et al. 1979). The Abbrook Clay and
 253 Sand Member is also likely to contain the Eocene–Oligocene boundary (Selwood et al. 1984) but
 254 because of the limited biostratigraphic data the exact positioning of the boundary is less certain.
 255 The Eocene – Oligocene boundary is one of the most prominent abrupt climatic events in the
 256 Cenozoic and is considered to represent the initiation of major permanent Palaeogene ice sheets
 257 on Antarctica (Miller et al., 2009; Coxall and Wilson, 2011; Hutchinson et al. 2021). The glaciation of
 258 Antarctica is thought to result from the tectonic opening of Southern Ocean gateways, which
 259 enabled the formation of the Antarctic Circumpolar Current and the subsequent thermal isolation
 260 of the Antarctic continent (e.g., Zachos et al. 2001) Modelling studies implicate low atmospheric
 261 CO₂ also as an important factor (DeConto and Pollard, 2003). The carbon isotope changes across
 262 this boundary are, however, less pronounced, and certainly less well documented in the terrestrial
 263 system. This is perhaps due to the lack of suitable terrestrial sections to study.

264 Nevertheless, the marine records (Zachos et al. 2001; Coxall and Wilson, 2011) show a $\delta^{13}\text{C}$
 265 excursion of ~1.0‰ in benthic foraminifera, peaking in the earliest Oligocene and followed by a
 266 decline to ~0.5‰, 1 million years after the boundary. The Petrockstowe and Bovey $\delta^{13}\text{C}_{\text{TOC}}$ data do
 267 show some correspondence with this marine record whereby for the Eocene stable but the most
 268 negative carbon values are observed, whereas the most positive carbon isotope values are present
 269 in the Oligocene. More positive $\delta^{13}\text{C}_{\text{TOC}}$ values have been linked to increased organic carbon burial
 270 (Coxall and Wilson, 2011). Our study therefore supports the notion that the surface ocean and
 271 atmosphere behaved as coupled reservoirs at this time, similar to other times in the Cenozoic
 272 (Jerrett et al. 2015; Cui et al., 2021; Lenz et al. 2022), as opposed to a decoupled system (cf.
 273 Holdgate et al. 2009; Fang et al., 2013), but more data is required to fully test this possibility.

274 **7. Conclusions**

275 In conclusion, and in agreement with Freshney et al., (1979), our palynological observations
 276 show that the Petrockstowe 1A and 1B cores, represent a succession of sand-filled fluvial channels
 277 followed by an ephemeral lake or lake margin setting. The Abbrook Clay and Sand and Southacre
 278 Clay and Lignite members of the Bovey Formation, represents a long-lived lake followed by

sediments representing more ephemeral lakes or lake margins periodically exposed with mires (see Chandler, 1964; Edwards, 1976). Our palynological data set further allows us to determine that isotope trends are not overly determined by the source of carbon in the basins.

Our study suggests that the observed $\delta^{13}\text{C}_{\text{TOC}}$ trends in the Eocene–Oligocene of the Petrockstowe and Bovey basins were primarily produced by variations of the carbon isotope ratios of terrestrial atmospheric carbon reservoirs. Even with our less than well constrained biostratigraphic control, the data indicate that the carbon isotope excursions seen in the Eocene and Oligocene could be associated with a number of transient global carbon isotopic shifts (e.g., the PETM). Our findings therefore appear to lend support to the surface ocean and atmosphere behaving as coupled reservoirs at this time.

Acknowledgements

We would like to thank the British Geological Survey for access to cores and Sibelco UK for access and information regarding South John Acres Lane Quarry. MC was funded by the Tertiary Education Trust Fund (TETFund) Abuja, via the intervention of academic staff training and development programme to the Federal University of Petroleum Resources, Effurun, Nigeria. This manuscript benefited from the extensive and constructive reviews of two anonymous reviewers, who are duly thanked.

References

- Abels, H.A., Clyde, W.C., Gingerich, P.D., Hilgen, F.J., Fricke, H.C., Bowen, G.J., Lourens, L.J. 2012. Terrestrial carbon isotope excursions and biotic change during Palaeogene hyperthermals. *Nature Geoscience*, 5, 326–329.
- Armstrong McKay D.I., Tyrrell, T., Wilson, P.A. 2016 Global carbon cycle perturbation across the Eocene-Oligocene climate transition. *Paleoceanography*, 31, 311–329.
- Aggarwal N., Agrawal, S., Thakur, B. 2019. Palynofloral, palynofacies and carbon isotope of Permian coal deposits from the Godavari Valley Coalfield, South India: Insights into the age, palaeovegetation and palaeoclimate. *International Journal of Coal Geology*, 214, 103285
- Aggarwal N., Mathews, R.P., Ansarie, A.H., Thakur, B. Agrawal, S., 2022. Palaeoenvironmental reconstruction for the Permian (lower Gondwana) succession of the Godavari Valley

- 307 Coalfield in southern India based on a combined palynofacies, carbon isotope, and
308 biomarker study. *Journal of Palaeogeography*, 2022, 11, 123–144.
- 309 Bechtel, A., Gratzner, R., Sachsenhofer, R.F., Gusterhuber, J., Lucke, A., Puttmann, W., 2008.
310 Biomarker and carbon isotope variation in coal and fossil wood of Central Europe through
311 the Cenozoic. *Palaeogeography, Palaeoclimatology, Palaeoecology* 262, 166–175.
- 312 Blyth, F.G.H. 1962. The structure of the north-eastern tract of the Dartmoor granite. *Quarterly*
313 *Journal of the Geological Society*, 118, 435.
- 314 Bristow, C. 1968. The derivation of the Tertiary sediments in the Petrockstowe Basin, North Devon,
315 *The Proceedings of the Ussher Society*, 2, 29–35.
- 316 Bristow, C., Robson, J. 1994. Palaeogene basin development in Devon. *Transactions of the*
317 *Institution of Mining and Metallurgy. Section B. Applied Earth Science*, 103, B163–B173.
- 318 Brown C.A., 2008. *Palynological Techniques. Second Edition.* Riding J.B., Warny, S. (editors).
319 American Association of Stratigraphic Palynologists, Dallas, USA, 137 p.
- 320 Chandler, M.E.J. 1957. The Oligocene Flora of the Bovey Tracey Lake Basin, Devonshire. *British*
321 *Museum (Natural History)*, 73–123.
- 322 Chandler, M.E.J. 1964. *The Lower Tertiary Floras.* British Museum (Natural History), London, 151 pp.
- 323 Collinson, M., Hooker, J., Gröcke, D. 2003. Cobham lignite bed and penecontemporaneous
324 macrofloras of southern England: A record of vegetation and fire across the Paleocene-
325 Eocene Thermal Maximum. *Special Papers-Geological Society of America*, 333–350.
- 326 Collinson, M.E., Steart, D.C., Harrington, G., Hooker, J.J., Scott, A.C., Allen, L.O., Glasspool, I.J.,
327 Gibbison, S.J. 2009. Palynological evidence of vegetation dynamics in response to
328 palaeoenvironmental change across the onset of the Paleocene-Eocene Thermal Maximum
329 at Cobham, Southern England. *Grana*, 48, 38–66.
- 330 Coxall, H. K., Wilson, P.A., 2011, Early Oligocene glaciation and productivity in the eastern
331 equatorial Pacific: Insights into global carbon cycling, *Paleoceanography*, 26, PA2221,
332 doi:10.1029/2010PA002021.

- 333 Cui, Y., Diefendorf, A.F., Kump, L.R. Jiang, S.J. Freeman, K.H. 2021. Synchronous Marine and
 334 Terrestrial Carbon Cycle Perturbation in the High Arctic During the PETM. *Paleoceanography*
 335 and *Paleoclimatology* 36, Article Number e2020PA003942
- 336 DeConto, R.M., Pollard, D. 2001. Rapid Cenozoic glaciation of Antarctica induced by declining
 337 atmospheric CO₂ . *Nature* 421, 245–249.
- 338 Dearman, W., 1963. Wrench-faulting in Cornwall and south Devon. *Proceedings of the Geologists’*
 339 *Association*, 74, 265–287.
- 340 Edwards, R.A. 1976. Tertiary sediments and structure of the Bovey Basin, south Devon. *Proceedings*
 341 *of the Geologists Association*, 87, 1–26.
- 342 Fang, L., Bjerrum, C.J., Hesselbo, S.P., Kotthoff, U., McCarthy, F.M.G., Huang, B., Ditchfield, P.W.,
 343 2013, Carbon-isotope stratigraphy from terrestrial organic matter through the Monterey
 344 event, Miocene, New Jersey margin (IODP Expedition 313). *Geosphere*, 9, 1303–1318.
- 345 Freshney, E. 1970. Cyclical sedimentation in the Petrockstowe Basin. *Proceedings of the Ussher*
 346 *Society*, 2, 179–189.
- 347 Freshney, E.C., Beer, K.E. and Wright, J.E. 1979. *Geology of the Country around Chulmleigh.*
 348 *Institute of Geological Sciences* 69p.
- 349 Freshney, E.C., Edwards, R.A., Isaac, K.P., Witte, G., Wilkinson, G.C., Boulter, M.C. and Bain, J.A.
 350 1982. A Tertiary basin at Dutson, near Launceston, Cornwall, England. *Proceedings of the*
 351 *Geologists Association*, 93, 395–402.
- 352 Garel, S., Dupuis, C., Quesnel, F., Jacob, J., Yans, J., Magioncalda, R., Flehoc, C., Schnyder, J. 2020
 353 Multiple early Eocene carbon isotope excursions associated with environmental changes in
 354 the Dieppe-Hampshire Basin (NW Europe). *BSGF - Earth Sciences Bulletin*, 191, 33.
- 355 Gröcke, D.R., Price, G.D., Robinson, S.A., Baraboshkin, E., Ruffell, A.H., and Mutterlose, J., 2005, The
 356 Valanginian (Early Cretaceous) positive carbon-isotope event recorded in terrestrial plants:
 357 *Earth and Planetary Science Letters*, 240, 495–509.
- 358 Hacquebard, P.A., Donaldson, J.R., 1969. Carboniferous coal deposition associated with floodplain
 359 and limnic environments in Nova Scotia. In: Dapples, E.C., Hopkins, M.E. (Eds.), *Environment*
 360 *of Coal Deposition. Geol. Soc. America Spec. Papers* 114, 143–191.

- 361 Heimhofer, U., Hochuli, P.A., Burla, S., Andersen, N., Weissert, H., 2003. Terrestrial carbon-isotope
362 records from coastal deposits (Algarve, Portugal): a tool for chemostratigraphic correlation
363 on an intrabasinal and global scale. *Terra Nova*, 15, 8–13.
- 364 Hodgson, E., Grimes, S.T., Fitzpatrick, M.E.J., Price, G.D., Hart, M.B., Leng, M.J. 2011. Paleogene
365 carbon isotope excursions in the Bunkers Hill borehole: Hampshire Basin, UK. *Proceedings of*
366 *the Geologists Association*, 122, 460–471.
- 367 Holdgate, G.R., McGowran, B., Fromhold, T., Wagstaff, B.E. Gallagher, S.J., Wallace, M.W., Sluiter,
368 I.R.K., Whitelaw, M., 2009 Eocene–Miocene carbon-isotope and floral record from brown
369 coal seams in the Gippsland Basin of southeast Australia. *Global and Planetary Change*, 65,
370 89–103.
- 371 Holloway, S., Chadwick, R.A. 1986. The Sticklepath-Lustleigh fault zone: Tertiary sinistral
372 reactivation of a Variscan dextral strike-slip fault. *Journal of the Geological Society*, 143,
373 447–452.
- 374 Hutchinson, D.K., Coxall, H.K., Lunt, D.J., Steinthorsdottir, M de Boer, A.M., Baatsen, M., von der
375 Heydt, A., Huber, M., Kennedy-Asser, A.T., Kunzmann, L., Ladant, J.B., Lear, C.H., Moraweck,
376 K., Pearson, P.N., Piga, E., Pound, M.J., Salzmann, U., Scher, HD., Sijp, W.P., Sliwinska, K.K.,
377 Wilson, P.A., Zhang, Z.S., 2021 The Eocene-Oligocene transition: a review of marine and
378 terrestrial proxy data, models and model data comparisons. *Climate of the Past*, 17, 269–
379 315.
- 380 Jerrett, R.M., Price, G.D., Grimes, S.T., and Dawson, A.T., 2015, A paleoclimatic and
381 paleoatmospheric record from peatlands accumulating during the Cretaceous–Paleogene
382 boundary event, Western Interior Basin, Canada: *Geological Society of America Bulletin*,
383 127, 1564–1582.
- 384 Lenz, O. K., Montag, M., Wilde, V., Methner, K., Riegel, W., and Mulch, A. 2022 Early Eocene carbon
385 isotope excursions in a lignite-bearing succession at the southern edge of the proto-North
386 Sea (Schöningen, Germany). *Clim. Past*, 18, 2231–2254.
- 387 Marchionni, D.L., 1980. Petrography and depositional environments of the Liddel seam, Upper
388 Hunter Valley, New South Wales. *Int. J. Coal Geol.* 1, 36–61.

- 389 Martín-Closas, C., Permanyer, A. and Vila, M.-J. 2005. Palynofacies distribution in a lacustrine basin.
390 Geobios, 38, 197–210.
- 391 McInerney, F.A., Wing, S.L. 2011. The Paleocene-Eocene thermal maximum: a perturbation of
392 carbon cycle, climate, and biosphere with implications for the future. Annual Review of
393 Earth and Planetary Sciences, 39, 489–516.
- 394 Miller, K.G., Wright, J.D., Katz, M.E., Wade, B.S., Browning, J.V. and Cramer B.S. 2009, Climate
395 threshold at the Eocene-Oligocene transition: Antarctic ice sheet influence on ocean
396 circulation, in SPE452: The Late Eocene Earth—Hothouse, Icehouse, and Impacts, Geol. Soc.
397 of Am. Spec. Pap., 452, 80301.
- 398 Nilsen, E. B., Anderson, L. D., and Delaney, M. L. 2003. Paleoproductivity, nutrient burial, climate
399 change and the carbon cycle in the western equatorial Atlantic across the Eocene/Oligocene
400 boundary. Paleoceanography, 18, 1057. <https://doi.org/10.1029/2002PA000804>
- 401 Pieńkowski, G., Waksmundzka, M., 2009. Palynofacies in Lower Jurassic epicontinental deposits of
402 Poland: tool to interpret sedimentary environments. Episodes, 32, 21–32.
- 403 Selwood, E.B., Edwards, R., Chester, J., Hamblin, R., Henson, M., Riddolls, B. and Waters, R. 1984.
404 Geology of the Country around Newton Abbot. Natural Environment Research Council.
405 London: HMSO. 212 p.
- 406 Sun, J., Ni, X., Bi, S., Wu, W., Ye, J., Meng, J., Windley, B.F., 2014 Synchronous turnover of flora,
407 fauna, and climate at the Eocene–Oligocene Boundary in Asia. Scientific Reports, 4, 7463
- 408 Tyson, R.V. 1995. Sedimentary Organic Matter: Organic Facies and Palynofacies. Springer.
- 409 Wilkinson, G.C., Boulter, M.C. 1981. Oligocene pollen and spores from the western part of the
410 British Isles. Palaeontographica Abteilung B, 175, 27–83.
- 411 Wilkinson, G.C., Bazley, R.A.B., Boulter, M.C. 1980. The geology and palynology of the Oligocene
412 Lough Neagh Clays, Northern Ireland. Journal of the Geological Society, 137, 65–75.
- 413 Zachos, J.C., Pagani, M., Sloan, L., Thomas, E., Billups, K., 2001. Trends, rhythms, and aberrations in
414 global climate 65 Ma to present. Science, 292, 686–693.

417

418 **Figure 1.** A. Map showing the southern part of the UK and location of inset. B. General geological
 419 map of the southwest UK showing Petrockstowe and Bovey basins (modified from Bristow
 420 and Robson 1994).

421 **Figure 2.** A. Map showing the location of Petrockstowe and Bovey basins (modified from Bristow
 422 and Robson 1994). B. Geological map of the Petrockstowe Basin showing the location of
 423 Borehole 1 (cores 1A and 1B) and the relative positions of the axial trough and the marginal
 424 shelves with their dividing fault (modified from Freshney 1970; Freshney et al., 1979). C.
 425 Geological map of the Bovey Basin showing the location of the South John Acres Lane
 426 Quarry (modified from Selwood et al. 1984).

427 **Figure 3.** Summary stratigraphic logs (and correlation) of the Bovey succession with data derived
 428 from (Edwards 1976) and Wilkinson (1979) cited by Selwood et al. (1984) and Petrockstowe
 429 with age data from Turner (cited by Freshney et al. 1979).

430 **Figure 4.** Phytoclasts from the Petrockstowe and Bovey basins. (A) Opaque lath-shaped phytoclast,
 431 sample MC95 Petrockstowe Basin; (B) Large opaque lath shaped phytoclast; sample MC3
 432 Petrockstowe; (C) Multicellular fungal ‘fruiting body’, sample MC19 Petrockstowe; (D) Mass
 433 of melanised fungal hyphae; sample MC58, Petrockstowe. (E) Multicellular fungal ‘fruiting
 434 body’ sample MC19, Petrockstowe. (F) Well preserved, pale brown in colour, partly
 435 translucent AOM sample SJAL029, Bovey Basin. (G) well preserved pale yellow AOM seen in
 436 transmitted white light; sample SJAL002, Bovey Basin (H) is a cross section of plant fragment
 437 sample SJAL013, Bovey Basin; (I) Phytoclast (biostructured) composed of gymnosperm
 438 tracheids; sample MC76 Petrockstowe Basin.

439 **Figure 5.** TOC and $\delta^{13}\text{C}_{\text{TOC}}$ data, compared to palynological data, Petrockstowe Basin, Devon. Age
 440 assignments based on Wilkinson (1979) cited by Selwood et al. (1984).

441 **Figure 6.** TOC and $\delta^{13}\text{C}_{\text{TOC}}$ data, compared to palynological data, Bovey Basin, Devon. Age
 442 assignments of Turner (cited by Freshney et al. 1979).

443 **Figure 7.** Distribution of the different categories of palynological matter in the ternary diagram
 444 (proposed by Hacquebard and Donaldson, 1969 and modified by Marchionni, 1980; and
 445 Aggarwal et al., 2022).

Fig 1





Fig 2





Fig 3






Fig 5






Fig 6





Our m/s has not been published previously (except in the form of an abstract) and it is not under consideration for publication elsewhere. The publication is approved by all authors.

