Faculty of Health: Medicine, Dentistry and Human Sciences

Peninsula Medical School

2023-05

# Time for Dementia: Quantitative evaluation of a dementia education programme for healthcare students

## Daley, S

https://pearl.plymouth.ac.uk/handle/10026.1/21418

10.1002/gps.5922 International Journal of Geriatric Psychiatry Wiley

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

DOI: 10.1002/gps.5922

#### **RESEARCH ARTICLE**



Geriatric Psychiatry WILEY

# Time for Dementia: Quantitative evaluation of a dementia education programme for healthcare students

Stephanie Daley<sup>1</sup> | Molly Hebditch<sup>1</sup> | Christopher Jones<sup>2</sup> | Stephen Bremner<sup>2</sup> | Yvonne Feeney<sup>1</sup> | Georgia Towson<sup>1</sup> | Juliet Wright<sup>3</sup> | Sube Banerjee<sup>4</sup>

<sup>1</sup>Centre for Dementia Studies, Brighton and Sussex Medical School, University of Sussex, Falmer, UK

<sup>2</sup>Department of Primary Care and Public Health, Brighton and Sussex Medical School, Falmer, UK

<sup>3</sup>Faculty of Health and Medical Sciences, University of Surrey, Guildford, UK

<sup>4</sup>Faculty of Health, University of Plymouth, Plymouth, UK

#### Correspondence

Stephanie Daley, Centre for Dementia Studies, Brighton and Sussex Medical School, University of Sussex, Falmer BN1 9RY, UK. Email: S.Daley@bsms.ac.uk

Funding information Health Education England

#### Abstract

**Objectives:** The future healthcare workforce needs the skills, attitudes, and empathy to better meet the needs of those with dementia. Time for Dementia (TFD) is an educational programme in which healthcare students from a range of professional groups visit a person with dementia and their family carer over a two-year period. The aim of this study was to evaluate its impact on student attitudes, knowledge and empathy towards dementia.

**Methods:** Measures of dementia knowledge, attitudes and empathy were administered to healthcare students at five universities in the south of England before and after (24 months) they completed the TFD programme. Data were also collected at equivalent time points for a control group of students who had not taken part in the programme. Outcomes were modelled using multilevel linear regression models.

**Results:** 2,700 intervention group students, and 562 control group students consented to participate. Students undertaking the TFD programme had higher levels of knowledge and positive attitudes at follow-up compared to equivalent students who did not undertake the programme. Our findings indicate a positive relationship between the number of visits undertaken and increasing dementia knowledge and attitudes. No substantial differences in the development of empathy was observed between groups.

**Conclusion:** Our findings suggest that TFD may be effective across professional training programmes and universities. Further research into the mechanisms of action is needed.

### KEYWORDS

Dementia, healthcare students, longitudinal education models

Abbreviation: TFD, Time for Dementia.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2023 The Authors. International Journal of Geriatric Psychiatry published by John Wiley & Sons Ltd.

#### Key points

- The data suggests the Time for Dementia programme is effective in improving the knowledge and attitudes of healthcare students across different professional groups and universities.
- Dementia education that meaningfully involves people with dementia and their carers can be delivered at scale for healthcare professionals.
- The number of visits (contact with people with dementia and their family carers) completed by students is positively related to student outcomes.

#### 1 | INTRODUCTION

The delivery of healthcare to people with dementia is widely recognised as being sub-optimal and a lack of understanding about dementia in the workforce is seen as a contributing factor.<sup>1</sup> The increasing numbers of people with dementia, driven by population ageing, demand a future healthcare workforce with the necessary knowledge, positive attitudes, and skills to deliver effective assessment and support to those living with the condition. To date, dementia education for undergraduate healthcare students has generally failed to meet this aspiration.<sup>2</sup> With its reliance on didactic teaching and emphasis on acute episodes of care, there is a need for undergraduate teaching to adapt to produce a future multiprofessional workforce able to rise to the complex challenges of ageing and dementia for both generalists and specialists.

Educational innovation to meet this challenge has taken place through the involvement of people living with dementia and their carers. Allowing healthcare students at the beginning of training to engage with people with dementia and their carers provides an opportunity to build more positive attitudes towards dementia before more negative and pessimistic attitudes can take hold and become fixed. Newer models of dementia education which directly involve people with dementia include longitudinal approaches and activitybased interventions such as intergenerational art projects.<sup>3</sup> The longitudinal model developed by Mordhardt and colleagues,<sup>4</sup> the Buddy Program, is an elective curricular component, where medical students are paired with a person with dementia and undertake community visits and activities together. The original programme has grown across North America, and the model has been used across a number of different contexts, such as the 'Friend for Rachel' model for pre-medical students (undergraduate and post-baccalaureate students preparing for medical school).5

The advantages of such longitudinal models are that healthcare students can hear first-hand what it is like to live with dementia and gain an understanding of living with long-term chronic illness, and through continuity of contact build relationships that may foster positive attitudes and person-centred care.<sup>6,7</sup>

One programme that builds on this concept, is the TFD programme which was developed in the UK, with support from Health Education England, at Brighton and Sussex Medical School in 2014.<sup>8</sup> Unlike the *Buddy Program*, which is designed to accommodate only a relatively small number of interested students in any institution, TFD is delivered as a mandatory component of the curriculum. This means that the whole year participates, not just those with an interest in dementia. It was also explicitly developed to be delivered to all healthcare student groups, not just medical students. The learning objectives of TFD are to increase positive attitudes, knowledge, and empathy in students by developing a broader understanding of the impact and experience of dementia.

An initial evaluation of the TFD programme showed that: (a) it was possible to deliver the programme at scale as a mandatory component of the curriculum; and (b) there were improvements in dementia knowledge and attitude change in medical students who undertook the programme compared to students who did not.<sup>9</sup> Qualitative interviews and focus groups with medical, nursing and paramedic students also identified improvements in the depth of dementia understanding, challenging of negative attitudes, and increased confidence in delivering person-centred care in practice.<sup>10,11</sup> Due to these positive preliminary findings, TFD was expanded across the south of England to include a wider range of healthcare students. The aim of the study reported here was to evaluate this wider and more representative expansion of TFD in terms of its impact on student attitudes, knowledge and empathy towards dementia.

#### 2 | METHODS

#### 2.1 | Design

A longitudinal survey was administered to healthcare students before (Baseline) and at the end of the programme (24 months), and at equivalent time points in training for a control group of students who had not taken part in the programme. Measures of knowledge, attitudes, and empathy were completed at both timepoints; demographics and previous experience of dementia were ascertained at baseline.

#### 2.2 | Time for Dementia intervention

Students undertaking the TFD programme visit a family (person with dementia and their carer) over a 2-year period. Students attend a 1 hour introductory session; which introduces the aims of the programmes and involves a person with dementia and a student who has

already undertaken the programme describing their experience. Students then undertake a 2-h preparatory session which makes clear the expectations of the programme, structure and guidelines for visits, and communication with people with dementia. After this, pairs or trios of students visit a person with dementia and their family for 2 h every 3-4 months for a total of 5-6 visits. During the visits, students follow a suggested visit structure linked to core learning objectives. For example, the first visit is focussed on understanding the family's background, events leading up to diagnosis of dementia and the experience of diagnosis, whereas the second visit is focussed on life since the diagnosis, and experiences of services. Further details of the intervention can be found in previous publications.<sup>8,9</sup> In this study, the programme was delivered to nursing, medical, paramedic and allied health professional (AHP; including occupational therapy, physiotherapy, speech and language therapy, radiology) students. Depending on curriculum fit, the programme starts in either the first or second year of training.

#### 2.3 | Study setting and participants

The intervention group consisted of healthcare students participating in TFD between 2014 and 2020 at five universities in the south of England: Brighton and Sussex Medical School, University of Surrey, University of Brighton, University of Greenwich, and Canterbury Christ Church University. Participation in TFD was mandatory for students where TFD was included in the curriculum, but participation in the evaluation was not. The control group included cohorts of equivalent healthcare students who did not take part in TFD at these five universities as well as Norwich Medical School. In total 24 intervention and 8 control student cohorts were invited to take part in the study. All students were invited to take part in the evaluation within scheduled lectures. They were given an information sheet in advance, and written consent was obtained. The study was approved by the NHS Health Research Authority London Queen Square Research Ethics Committee (15/LO/0046).

In March of 2020 the COVID-19 pandemic led to the cessation of face-to-face visits to TFD families. The student cohorts affected by this disruption had a reduced number of visits and their last contact with their allocated families was conducted over the telephone. For the analysis, this disruption was recorded as those who completed the programme and their 24-month follow-ups before the 1<sup>st</sup> of March 2020 (no disruption) or post this date (possible disruption).

#### 2.4 | Outcomes measures

All students completed the same battery of instruments at baseline and 24 months later. For all outcomes, higher scores indicate a higher, or more positive, level of the attribute.

Approaches to Dementia Questionnaire (ADQ),<sup>12</sup> 19 items assessing attitudes towards dementia, each scored 1–5 (total: 19–95) with two subscales 'hopefulness' and 'person-centeredness',

Geriatric Psychiatry \_WILEY.

- 10991166, 2023, 5, Downloaded from https: //onlinelibrary.wiley.com/doi/10.1002/gps.5922 by Plymouth University, Wiley Online Library on [13/10/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/doi/10.1002/gps.5922 by Plymouth University, Wiley Online Library on [13/10/2023]. /onlinelibrary.wiley.com/terms -and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License
- Alzheimer's Disease Knowledge Scale (ADKS),<sup>13</sup> 30 true/false items (total: 0–30) assessing students' knowledge of Alzheimer's disease;
- Dementia Knowledge Questionnaire (DK-20),<sup>14</sup> 20 true/false items (total: 0–20) assessing dementia knowledge;
- Dementia Attitude Scale (DAS),<sup>15</sup> 20 items assessing attitudes toward dementia scored on a 7-point Likert-type scale (total: 20–140) with two subscales 'dementia knowledge' and 'social comfort';
- Medical Condition Regard Scale (MCRS),<sup>16</sup> 11 items scored on a 6-point Likert-type scale (total: 11–66) assessing the extent students find patients with a given condition to be enjoyable, treatable and worthy of medical resources; and
- Jefferson Scale of Empathy (JSE): Medical Student Version and Health Professional Version.<sup>17</sup> 20 items scored on a 7-point Likert-type scale (total: 20–140) measuring student empathy.

#### 2.5 | Statistical analysis

Demographic and outcome data were summarised by group (intervention or control) using means and standard deviations for normally distributed variables, medians and means for skewed continuous variables and frequencies and percentages in categories for categorical variables.

For the primary analysis, outcomes were modelled using twolevel multilevel linear regression models. Students attended visits in pairs or trios and therefore group membership was included as a random effect. Independent variables included in the models were: the outcome at baseline (continuous), student age (continuous), student gender (female/male), student previous experience of dementia (yes/no) and ethnicity (White British or European/other ethnic groups), COVID-19 disruption (yes/no), and intervention group (intervention/control).

The secondary analysis included replicating these models, and adjusting for the same variables, except the intervention group (intervention/control) which was replaced with the number of TFD visits completed by students to explore the relationship between the number of visits and outcomes. This was possible due to variation in the number of visits completed, influenced in part by the disruption caused by COVID-19. For both analyses assumptions for linear regression were examined, and robust standard errors were calculated as residuals deviated from normality. Analyses were performed in Stata 17.0.<sup>18</sup>

#### 3 | RESULTS

#### 3.1 | Student recruitment and demographics

In the intervention group, 3619 students were eligible<sup>1</sup> to take part and 2700 (74%) consented to participate in the evaluation. In the control group, 863 students were eligible and 562 (65%) consented. At baseline, 2570 (95%) of the consenting intervention group and 498 (89%) of the consenting control group contributed to the data.

#### TABLE 1 Descriptive summaries of demographics and other characteristics of student participants.

| DALEY ET A | L. |
|------------|----|
|------------|----|

|   | Control (n     | = 562)              | Intervention ( $n = 2,700$ ) |                     | Total (n = 3,262) |                     |  |
|---|----------------|---------------------|------------------------------|---------------------|-------------------|---------------------|--|
|   | Median<br>23.0 | IQR<br>20.0 to 29.0 | Median<br>21.0               | IQR<br>19.0 to 27.0 | Median<br>21.0    | IQR<br>19.0 to 28.0 |  |
| Age (years)                                 | INO.           | 70                  | INO.                         | 70                  | INU.              | 70                  |  |
| Nedical                                     | 110            | 10.0                | 601                          | 22.4                | 742               | 22.0                |  |
| Nurse                                       | 224            | 19.7                | 1 274                        | 23.4                | 1 5 0 9           | 22.0                |  |
| Deremodie                                   | 9 <i>4</i>     | 14.9                | 217                          | 47.2                | 1,590             | 49.0                |  |
|   | 04<br>42       | 14.7                | 479                          | 17.7                | 401               | 12.3                |  |
|   | 42             | 7.5                 | 470                          | 17.7                | 520               | 15.9                |  |
| Brighton and Suscey Medical School          | 0              | 0.0                 | 601                          | 22.4                | 421               | 10.2                |  |
|   | 0              | 0.0                 | 1 01 4                       | 23.4                | 1 055             | 17.5                |  |
|   | 41             | 7.3                 | 1,014                        | 37.0                | 1,055             | 32.3                |  |
|   | 220            | 39.1                | 445                          | 16.5                | 665               | 20.4                |  |
|   | 112            | 19.9                | 0                            | 0.0                 | 112               | 3.4                 |  |
| University of Greenwich                     | 68             | 12.1                | 149                          | 5.5                 | 217               | 6.7                 |  |
| Canterbury Christ Church University         | 121            | 21.5                | 461                          | 17.1                | 582               | 17.8                |  |
| Student gender                              |                |                     |                              |                     |                   |                     |  |
| Male  | 109            | 19.8                | 572                          | 21.4                | 681               | 21.1                |  |
| Female                                      | 442            | 80.2                | 2,095                        | 78.5                | 2,537             | 78.8                |  |
| Other                                       | 0              | 0.0                 | 2                            | 0.1                 | 2                 | 0.1                 |  |
| Student ethnicity                           |                |                     |                              |                     |                   |                     |  |
| White British/European                      | 383            | 70.7                | 1,899                        | 72.6                | 2,282             | 72.3                |  |
| Mixed/Multiple ethnic groups                | 14             | 2.6                 | 87                           | 3.3                 | 101               | 3.2                 |  |
| Asian/Asian British                         | 58             | 10.7                | 277                          | 10.6                | 335               | 10.6                |  |
| Black/African/Caribbean/Black British       | 77             | 14.2                | 282                          | 10.8                | 359               | 11.4                |  |
| Other                                       | 10             | 1.8                 | 70                           | 2.7                 | 80                | 2.5                 |  |
| Experience of knowing someone with demen    | ntia           |                     |                              |                     |                   |                     |  |
| Yes   | 324            | 60.0                | 1,433                        | 54.9                | 1,757             | 55.7                |  |
| No  | 216            | 40.0                | 1,179                        | 45.1                | 1,395             | 44.3                |  |
| Details of students' experience with dement | ia             |                     |                              |                     |                   |                     |  |
| Family member/Friend                        | 104            | 32.2                | 521                          | 36.6                | 625               | 35.8                |  |
| Paid/Unpaid work                            | 150            | 46.4                | 622                          | 43.7                | 772               | 44.2                |  |
| Both  | 69             | 21.4                | 279                          | 19.6                | 348               | 19.9                |  |
| COVID-19 disruption                         |                |                     |                              |                     |                   |                     |  |
| No  | 483            | 85.9                | 1,401                        | 51.9                | 1,884             | 57.8                |  |
| Yes   | 79             | 14.1                | 1,299                        | 48.1                | 1,378             | 42.2                |  |
| No. of completed TFD visits                 |                |                     |                              |                     |                   |                     |  |
| 0   | 562            | 100.0               | 3                            | 0.3                 | 565               | 32.6                |  |
| 1   | 0              | 0.0                 | 78                           | 6.7                 | 78                | 4.5                 |  |
| 2   | 0              | 0.0                 | 104                          | 8.9                 | 104               | 6.0                 |  |
| 3   | 0              | 0.0                 | 176                          | 15.0                | 176               | 10.2                |  |
| 4   | 0              | 0.0                 | 222                          | 19.0                | 222               | 12.8                |  |

#### TABLE 1 (Continued)

|             | Control ( <i>n</i> = 562) |                          | Interventio           | n ( <i>n</i> = 2,700)    | Total (n = 3,262)     |                          |
|-------------|---------------------------|--------------------------|-----------------------|--------------------------|-----------------------|--------------------------|
| Age (years) | Median<br>23.0<br>No.     | IQR<br>20.0 to 29.0<br>% | Median<br>21.0<br>No. | IQR<br>19.0 to 27.0<br>% | Median<br>21.0<br>No. | IQR<br>19.0 to 28.0<br>% |
| 5           | 0                         | 0.0                      | 309                   | 26.4                     | 309                   | 17.8                     |
| 6           | 0                         | 0.0                      | 279                   | 23.8                     | 279                   | 16.1                     |

Abbreviations: AHP, Allied health professionals (Occupational therapy; Physiotherapy, Speech and language therapy and Radiography); IQR, interquartile range; TFD, Time for Dementia.

## **TABLE 2** Descriptive summaries of scores on outcome measures at baseline.

|                                   | Baseline |      |     |              |      |       |       |      |       |
|-----------------------------------|----------|------|-----|--------------|------|-------|-------|------|-------|
|                                   | Control  |      |     | Intervention |      |       | Total |      |       |
|                                   | Mean     | SD   | n   | Mean         | SD   | n     | Mean  | SD   | n     |
| ADQ total (19-95)                 | 78.1     | 7.6  | 489 | 79.0         | 6.6  | 2,549 | 78.8  | 6.8  | 3,038 |
| ADQ-Hopeful (8-40)                | 28.2     | 4.3  | 489 | 28.9         | 4.0  | 2,549 | 28.8  | 4.1  | 3,038 |
| ADQ—Person-centredness<br>(11-55) | 49.9     | 4.9  | 489 | 50.1         | 4.3  | 2,549 | 50.0  | 4.4  | 3,038 |
| ADKS total (0-30)                 | 22.6     | 3.1  | 496 | 22.9         | 3.1  | 2,582 | 22.8  | 3.1  | 3,078 |
| DK-20 total (0-20)                | 14.5     | 3.0  | 498 | 15.1         | 2.9  | 2,544 | 15.0  | 2.9  | 3,042 |
| MCRS total (11-66)                | 53.9     | 7.3  | 489 | 54.2         | 6.8  | 2,570 | 54.2  | 6.8  | 3,059 |
| DAS total (20-140)                | 112.0    | 14.6 | 483 | 112.3        | 13.4 | 2,549 | 112.2 | 13.6 | 3,032 |
| DAS-Comfort (10-70)               | 50.5     | 9.8  | 483 | 50.4         | 9.5  | 2,549 | 50.4  | 9.6  | 3,032 |
| DAS—Knowledge (10-70)             | 61.5     | 7.5  | 483 | 61.8         | 6.4  | 2,549 | 61.8  | 6.6  | 3,032 |
| JSE total (20-140)                | 113.9    | 12.2 | 466 | 114.9        | 11.7 | 2,503 | 114.8 | 11.8 | 2,969 |

Abbreviations: ADKS, Alzheimer's Disease Knowledge Scale; ADQ, Approaches to Dementia Questionnaire; DAS, Dementia Attitude Scale; DK-20, Dementia Knowledge Questionnaire; JSE, Jefferson Scale of Empathy; MCRS, Medical Condition Regard Scale; SD, standard deviation.

## **TABLE 3** Descriptive summaries of scores on outcome measures at 2 year follow up.

|                                   | 24 Month follow up |      |              |       |      |       |       |      |       |
|-----------------------------------|--------------------|------|--------------|-------|------|-------|-------|------|-------|
|                                   | Control            |      | Intervention |       |      | Total |       |      |       |
|                                   | Mean               | SD   | n            | Mean  | SD   | n     | Mean  | SD   | n     |
| ADQ total (19-95)                 | 79.7               | 6.6  | 232          | 80.6  | 6.8  | 1,199 | 80.4  | 6.8  | 1,431 |
| ADQ-Hopeful (8-40)                | 29.4               | 3.9  | 232          | 29.9  | 4.0  | 1,199 | 29.8  | 4.0  | 1,431 |
| ADQ—Person-centredness<br>(11-55) | 50.3               | 4.1  | 232          | 50.7  | 4.2  | 1,199 | 50.7  | 4.2  | 1,431 |
| ADKS total (0-30)                 | 24.3               | 2.8  | 236          | 24.7  | 2.8  | 1,215 | 24.6  | 2.8  | 1,451 |
| DK-20 total (0-20)                | 15.5               | 2.8  | 232          | 16.3  | 2.4  | 1,199 | 16.2  | 2.5  | 1,431 |
| MCRS total (11-66)                | 53.6               | 6.9  | 233          | 54.0  | 6.7  | 1,199 | 53.9  | 6.8  | 1,432 |
| DAS total (20-140)                | 113.8              | 13.1 | 231          | 116.6 | 12.4 | 1,196 | 116.2 | 12.5 | 1,427 |
| DAS-Comfort (10-70)               | 51.6               | 9.1  | 231          | 53.8  | 8.5  | 1,196 | 53.5  | 8.6  | 1,427 |
| DAS—Knowledge (10-70)             | 62.2               | 6.5  | 231          | 62.8  | 6.0  | 1,196 | 62.7  | 6.1  | 1,427 |
| JSE total (20-140)                | 115.0              | 12.9 | 228          | 116.8 | 12.9 | 1,182 | 116.5 | 12.9 | 1,410 |

Abbreviations: ADKS, Alzheimer's Disease Knowledge Scale; ADQ, Approaches to Dementia Questionnaire; DAS, Dementia Attitude Scale; DK-20, Dementia Knowledge Questionnaire; JSE, Jefferson Scale of Empathy; MCRS, Medical Condition Regard Scale; SD, standard deviation.

5 of 9

WILEY Geriatric Psychiatry

| Outcome                 | Coefficient | 95% CI        | Scaled coefficient and<br>95% CI as % of scale | P-value |
|-------------------------|-------------|---------------|--|---------|
| Intervention vs Control |             |               |  |         |
| ADQ Total               | 0.22        | -0.73 to 1.17 | , <b>i</b> iii                                 | 0.652   |
| ADQ - Hopeful           | 0.18        | -0.36 to 0.72 | ⊢ i i i i i i i i i i i i i i i i i i i        | 0.515   |
| ADQ - Personcentrednes  | s 0.10      | -0.51 to 0.71 | ⊢ <b>i</b> −i                                  | 0.747   |
| ADKS                    | 0.29        | -1.25 to 0.66 | i i i i i i i i i i i i i i i i i i i          | 0.126   |
| DK-20                   | 0.59        | 0.20 to 0.97  | ••••   | 0.003   |
| DAS Total               | 2.27        | 0.61 to 3.94  | :⊢++   | 0.007   |
| DAS - Comfort           | 2.09        | 0.89 to 3.29  |  | ⊣ 0.001 |
| DAS - Knowledge         | 0.21        | -0.61 to 1.04 | ⊢÷+1   | 0.613   |
| MCRS                    | -0.26       | -1.25 to 0.73 | ⊢ + <mark>i</mark> − I                         | 0.605   |
| JSE                     | -0.18       | -1.87 to 1.51 | ⊢ <b>∔</b> −i                                  | 0.834   |
|                         |             |               | Favours TFD                                    | →       |
|                         |             |               | [+++++++++++++++++++++++++++++++++++++         | п       |
| Number of visite        |             |               | -6 -5 -4 -3 -2 -1 0 1 2 3 4 5                  | 6       |
| Number of visits        |             |               |  |         |
| ADQ Total               | 0.16        | 0.01 to 0.31  | <u>+</u>                                       | 0.038   |
| ADQ - Hopeful           | 0.09        | 0.00 to 0.18  |  | 0.055   |
| ADQ - Personcentrednes  | ss 0.08     | -0.01 to 0.18 | i <b>∔</b> -∎-1                                | 0.088   |
| ADKS                    | 0.08        | 0.01 to 0.14  | <b>;</b> ⊢                                     | 0.020   |
| DK-20                   | 0.13        | 0.07 to 0.19  | · · · · ·                                      | < 0.001 |
| DAS Total               | 0.53        | 0.25 to 0.82  | • • • • •                                      | < 0.001 |
| DAS - Comfort           | 0.41        | 0.20 to 0.61  | · · · · · · · · · · · · · · · · · · ·          | < 0.001 |
| DAS - Knowledge         | 0.13        | -0.01 to 0.27 | Ļ.   | 0.062   |
| MCRS                    | 0.14        | -0.02 to 0.30 | ÷-+I   | 0.082   |
| JSE                     | 0.27        | -0.03 to 0.57 | i-∎-i  | 0.078   |
|                         |             |               | Favours TFD                                    | ►       |
|                         |             |               |  | -       |
|                         |             |               | -1.0 -0.5 0.0 0.5 1.0                          | )       |

DALEY ET AL.

10991166, 2023, 5, Downloaded from https: //onlinelibrary.wiley.com/doi/10.1002/gps.5922 by Plymouth University, Wiley Online Library on [13/10/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/term on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

FIGURE 1 Adjusted multilevel models comparing 2-year outcomes in Time for Dementia (TFD) recipients with controls and association of 2-year outcomes with the number of TFD visits, with adjusted regression coefficients as a percentage of each scale. ADKS, Alzheimer's Disease Knowledge Scale; ADQ, Approaches to Dementia Questionnaire; DAS, Dementia Attitude Scale; DK-20, Dementia Knowledge Questionnaire; JSE, Jefferson Scale of Empathy; MCRS, Medical Condition Regard Scale; CI, confidence interval. Each multilevel model is adjusted for the student outcome at baseline, age, gender, previous experience of dementia, ethnicity, COVID-19 disruption, and a random effect is included for student group identity.

At 24 months 1215 (45%) in the intervention group and 236 (42%) control group completed a follow up assessment. Details of demographics, COVID-19 disruption, and the number of completed visits are presented in Table 1.

#### 3.2 | Intervention group

The scores on the study outcomes for baseline and follow up are presented in Tables 2 and 3. Figure 1 presents the results of the adjusted multilevel models comparing 2-year outcomes in TFD recipients to controls for the primary analysis. To illustrate the magnitude of the differences Figure 1 also displays the adjusted regression coefficients for the TFD group versus the control group as a percentage of each scale. Overall there was strong evidence supporting improvements for those receiving TFD in attitudes to dementia in the DAS total (coefficient: 2.27, 95% confidence interval [95% CI]: 0.61–3.94, p = 0.007) and its comfort scale (2.09, 95% CI:

0.89–3.29, p < 0.001). There was also strong evidence for more positive scores of dementia knowledge on the DK-20 (0.59, 95% CI: 0.20–0.97, p = 0.003) but not for the ADKS (0.29, 95% CI: -1.25 to 0.66, p = 0.126) or DAS Knowledge subscale (0.21, 95% CI: -0.61 to 1.04, p = 0.613). There was no evidence for differences on the other attitude measures; for the ADQ (0.22, 95% CI: -0.73 to 1.17, p = 0.652) or its subscales, MCRS (-0.26, 95% CI: -1.25 to 0.73, p = 0.605) and JSE (-0.18, 95% CI: -1.87 to 1.51, p = 0.834).

#### 3.3 | Number of visits

Figure 1 presents the results of the adjusted multilevel models exploring 2-year outcomes with the number of TFD visits completed and illustrates these differences with the adjusted regression coefficients for the number of visits as a percentage of each scale. There was evidence that the number of visits completed by students was associated with increased positive attitudes as measured by the ADQ (0.16, 95% CI: 0.01–0.31, p = 0.038) and weaker evidence for its hopefulness (0.09, 95% CI: 0.00–0.18, p = 0.055) and its personcentredness subscale (0.08, –0.01 to 0.18, p = 0.088). There was strong evidence of differences in the DAS (0.53, 95% CI: 0.25–0.82, p < 0.001) including its comfort subscale (0.41, 95% CI: 0.20–0.61, p < 0.001) and less strong evidence for its knowledge subscale (0.13–0.01 to 0.27, p = 0.062). Increases in knowledge were observed as measured by the DK-20 (0.13, 95% CI: 0.07–0.19, p < 0.001) and the ADKS (0.08, 95% CI: 0.01–0.14, p = 0.020). There was weak evidence for a positive association with visit numbers for the MCRS (0.14, 95% CI: –0.02 to 0.30, p = 0.082) and JSE (0.27, 95% CI: –0.03 to 0.57, p = 0.078).

#### 4 | DISCUSSION

Our findings indicate that, across professional backgrounds, students undertaking the TFD programme developed higher levels of knowledge and more positive attitudes to people with dementia compared with equivalent students not enrolled in the programme. Our results also suggest that these positive impacts increase with the number of visits undertaken. We believe these to be the first quantitative data indicating the potential value of a dementia training initiative across the range of healthcare professionals in training and applied at a whole cohort level. Taken with the positive multi-professional qualitative analyses of TFD we have completed<sup>10</sup> these data provide encouraging evidence for the positive impact of TFD on healthcare professionals in training.

These findings build on our earlier quantitative evidence of the effectiveness of TFD for medical students alone.<sup>9</sup> They indicate that TFD has a complementary positive impact on student learning about dementia in addition to traditional didactic class-room based teaching and placements which do not provide the profound level of deep learning that more experiential learning can offer.<sup>19</sup> In TFD, the experiential component is longitudinal first-hand exposure to the challenges that people living with dementia and their carers experience on a daily basis. The students are in effect taught by the people with dementia and their families, and it enables them to see the world and healthcare through their eyes.<sup>21</sup> This identification with the person with dementia and their carer and relational learning which takes place appears to be powerful in terms of knowledge and attitude change.

These findings are also consistent with positive outcomes identified by other experiential longitudinal models, such as the Buddy program.<sup>20</sup> In such programmes, participation in the programme is elective, and students are matched with a person with dementia based on similar interests. Therefore, there are notable differences between TFD and these programmes, specifically the mandatory nature of the programme, the pairing of 2-3 students with a family rather than 1 student and pairing based on geography rather common interests. This might explain the modest nature of TFD positive outcomes, with a focus on broadly upskilling the entire medical student cohort as opposed to more in-depth outcomes for a smaller number of interested students. It is interesting to note that there appears to be a 'dose response' relationship, in that student outcomes are higher for students who complete more visits. The importance of longitudinal contact is supported by our qualitative findings<sup>10</sup> which identified the importance of relational learning within TFD. Learning takes place within an authentic meaningful relationship between the person with dementia, their carer and students, which deepens over time. Further research is needed to understand the optimal number of visits, and how the longitudinal nature of the visits underpins the effects observed.

As in our earlier study of medical students,<sup>9</sup> TFD did not appear to change quantitative measures of empathy, unlike our two qualitative studies<sup>10,21</sup> which identified positive change in terms of empathy towards people with dementia. In terms of potential explanations, first JSE is a generic empathy measure, and its potential limitation in measuring empathy towards people with dementia has been identified in a wider systematic review of empathy interventions for people with dementia.<sup>22</sup> It may be that the measure itself is not sufficiently responsive to change in dementia. Second, there may be issues related to social desirability in students, due to a perception that it may be professionally unacceptable to admit to lower levels of empathy.<sup>23</sup> Third, it is interesting to note that there was a positive change in both intervention and control groups, which challenges the existing body of work that suggests a decline in empathy during healthcare training<sup>24</sup>; and it may be that other factors such as placement experiences including role models and ward culture might have positive impacts on empathy. Finally, it may be that alternative methods of assessing empathy, for example, patient or carer feedback, might have yielded a more accurate reflection of empathy in practice,<sup>25</sup> which could have been a more meaningful outcome.

There are two main limitations of this study. First, the study was affected by the COVID-19 pandemic through unplanned alterations to the TFD evaluation and programme delivery. Data collection sessions could not take place in person as planned. Despite active attempts to obtain follow up measures from students, overall this means that the loss to follow up was greater than expected. In terms of the TFD programme and altered delivery, we do not know how the final telephone interactions that took place after March 2020, compared to face-to-face visits. Life for both families and students would have been heavily impacted by the pandemic, and this is likely to have been the focus of the telephone contact, and not wider dementia learning. The impact of the pandemic therefore adds complexity to the interpretation of the research. Second, due to the study design, a number of potential confounders and biases cannot be accounted for in the analyses such as differences in the content of university curriculums, in control groups, and potential responder bias across intervention groups and at follow up. However, this study does have strengths. First, it is a real-life pragmatic study, which has managed to recruit a high number of participants. Second, the use of comparison groups and controlling for potential confounders (including COVID-19 disruption) within the statistical modelling strengthens the conclusions that can be drawn. Third, the inclusion of a broad range

WILEY\_Geriatric Psychiatry

of professional groups in training across different universities widens the generalisability of the findings.

#### 5 | CONCLUSION

There is a pressing global requirement that the future workforce is better equipped to support people living with dementia. Evidence based dementia education programmes, such as TFD illustrate that this aspiration can be met and how it can be met. The size and scope of this evaluation demonstrates that TFD can be delivered across a range of training programmes and universities, and suggests that in doing so, TFD can improve the dementia knowledge and attitudes of future healthcare professionals. The future of care for complex conditions such as dementia is teamwork, and strategies like TFD that enable symmetrical understanding of patient needs and experiences are likely to be of particular value in facilitating effective teamwork. The demonstration that people with dementia and their family carers can be deployed at scale as a teaching resource for healthcare students is also an encouragement to the meaningful involvement of patient groups in healthcare education. Further research is required to more fully understand the mechanisms of action of TFD, as well as differential impacts between student types, the optimal number of sessions, and whether change in knowledge and attitude translates into practice.

#### ACKNOWLEDGEMENTS

The authors would like to acknowledge all the contributing research assistants: Leila Morley, Gillian Sleater, Nazire Akarsu, Joanna Pooley, Serena Thomas, and Frankie Morrice. We would like to thank our collaborators: Dr Wendy Grosvenor, Professor Ramin Nilforooshan, Professor Chris Fox, Patricia Fordham, John Krohne, Mary Kirk and Dr Julie Cooper. Special thanks are due to all the families and students that have taken part, and Alzheimer's society for their work in recruiting and supporting families in the programme. This work was supported by Health Education England. Health Education England had no role in the design or completion of this study.

#### CONFLICT OF INTEREST STATEMENT

The authors declare no conflicts of interest.

#### DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

#### ORCID

Stephanie Daley D https://orcid.org/0000-0002-5842-0492 Sube Banerjee D https://orcid.org/0000-0002-8083-7649

#### ENDNOTE

<sup>1</sup> The number approached for research could not be calculated as registers for attendance at recruitment sessions or the numbers approached via email were not available. Therefore, the number presented for eligible students is based on intake numbers of student cohorts.

#### REFERENCES

- Lynch C. World Alzheimer Report 2019: attitudes to dementia, a global survey: public health: engaging people in ADRD research. *Alzheimer's and Dementia*. 2020;16(S10):e038255. https://doi.org/10. 1002/alz.038255
- Scott TL, Kugelman M, Tulloch K. How medical professional students view older people with dementia: implications for education and practice. *PLoS One*. 2019;14(11):e0225329. https://doi.org/10.1371/ journal.pone.0225329
- Lokon E, Li Y, Parajuli J. Using art in an intergenerational program to improve students' attitudes toward people with dementia. *Gerontology and Geriatr Educ.* 2017;38(4):407-424. https://doi.org/10. 1080/02701960.2017.1281804
- Morhardt D. Educating medical students on Alzheimer's disease and related disorders: an overview of the Northwestern University Buddy Program. *Dementia*. 2006;5(3):448-456. https://doi.org/10. 1177/147130120600500311
- Goldman JS, Trommer AE. A qualitative study of the impact of a dementia experiential learning project on pre-medical students: a friend for Rachel. *BMC Med Educ*. 2019;19(1):127. https://doi.org/10. 1186/s12909-019-1565-3
- Bartlett M, Muir F. A new model of undergraduate clinical education? Br J Gen Pract. 2018;68(670):216-217. https://doi.org/10.3399/ bjgp18x695885
- Walters L, Greenhill J, Richards J, et al. Outcomes of longitudinal integrated clinical placements for students, clinicians and society. *Med Educ.* 2012;46(11):1028-1041. https://doi.org/10.1111/j.1365-2923.2012.04331.x
- Banerjee S, Farina N, Daley S, et al. How do we enhance undergraduate healthcare education in dementia? A review of the role of innovative approaches and development of the Time for Dementia Programme. *Int J Geriatr Psychiatr.* 2017;32(1):68-75. https://doi.org/ 10.1002/gps.4602
- Banerjee S, Jones C, Wright J, et al. A comparative study of the effect of the Time for Dementia programme on medical students. *Int J Geriatr Psychiatr.* 2021;36(7):1011-1019. https://doi.org/10.1002/ gps.5532
- Daley S, Feeney Y, Grosvenor W, et al. A qualitative evaluation of the effect of a longitudinal dementia education programme on healthcare student knowledge and attitudes. *Age Ageing*. 2020;46(6): 1080-1086. https://doi.org/10.1093/ageing/afaa182
- 11. Grosvenor W. Student Nurse Perspectives on the Impact of Longitudinal Home Visits to People with Dementia and Their Carers Whole Sight: New Ways of Seeing. University of Surrey; 2019.
- 12. Lintern T, Woods B, Phair L. Before and after training: a case study of intervention. *J Dement Care*. 2000;8(1):15-17.
- Carpenter BD, Balsis S, Otilingam PG, Hanson PK, Gatz M. The Alzheimer's disease knowledge scale: development and psychometric properties. *Gerontologist*. 2009;49(2):236-247. https://doi. org/10.1093/geront/gnp023
- Shanahan N, Orrell M, Schepers AK, Spector A. The development and evaluation of the DK-20: a knowledge of dementia measure. *Int Psychogeriat*. 2013;25(11):1899-1907. https://doi.org/10.1017/s104 1610213001142
- O'Connor ML, McFadden SH. Development and psychometric validation of the dementia attitudes scale. *Int J Alzheimer's Dis.* 2010; 2010:1-10. https://doi.org/10.4061/2010/454218
- Christison GW, Haviland MG, Riggs ML. The medical condition regard scale: measuring reactions to diagnoses. *Acad Med.* 2002;77(3): 257-262. https://doi.org/10.1097/00001888-200203000-00017
- Hojat M, Mangione S, Nasca TJ, et al. The Jefferson scale of physician empathy: development and preliminary psychometric data. *Educ Psychol Meas.* 2001;61(2):349-365. https://doi.org/10.1177/00131 640121971158
- Stata 17.0. Stata Statistical Software: Release. 17 ed. StataCorp LLC; 2021.

- Dacre JE, Fox RA. How should we be teaching our undergraduates? Ann Rheum Dis. 2000;59(9):662-667. https://doi.org/10.1136/ard.59. 9.662
- Morhardt DJ. The Effects of an Experiential Learning and Mentorship Program Pairing Medical Students and Persons with Cognitive Impairment: A Qualitative Content Analysis. Loyola University Chicago; 2013.
- Grosvenor W, Gallagher A, Banerjee S. Reframing dementia: nursing Students' relational learning with rather than about people with dementia. A constructivist grounded theory study. *Int J Geriatr Psychiatr.* 2020;36(4):558-565. https://doi.org/10.1002/gps. 5452
- Brown EL, Agronin ME, Stein JR. Interventions to enhance empathy and person-centered care for individuals with dementia: a systematic review. *Res Gerontol Nurs.* 2020;13(3):158-168. https://doi.org/ 10.3928/19404921-20191028-01
- Chua JYX, Ang E, Lau STL, Shorey S. Effectiveness of simulation-based interventions at improving empathy among healthcare students: a

systematic review and meta-analysis. *Nurse Educ Today*. 2021;104: 105000. https://doi.org/10.1016/j.nedt.2021.105000

- Hojat M, Vergare MJ, Maxwell K, et al. The devil is in the third year: a longitudinal study of erosion of empathy in medical school. *Acad Med.* 2009;84(9):1182-1191. https://doi.org/10.1097/acm.0b013e3 181b17e55
- Batt-Rawden SA, Chisolm MS, Anton B, Flickinger TE. Teaching empathy to medical students: an updated, systematic review. *Acad Med.* 2013;88(8):1171-1177. https://doi.org/10.1097/acm.0b013e 318299f3e3

How to cite this article: Daley S, Hebditch M, Jones C, et al. Time for Dementia: quantitative evaluation of a dementia education programme for healthcare students. *Int J Geriatr Psychiatry*. 2023;e5922. https://doi.org/10.1002/gps.5922