

2023-10

# Quality standards for the management of alcohol-related liver disease: consensus recommendations from the British Association for the Study of the Liver and British Society of Gastroenterology ARLD special interest group

Parker, R

<https://pearl.plymouth.ac.uk/handle/10026.1/21576>

---

10.1136/bmjgast-2023-001221

BMJ Open Gastroenterology

BMJ

---

*All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.*

# Quality standards for the management of alcohol-related liver disease: consensus recommendations from the British Association for the Study of the Liver and British Society of Gastroenterology ARLD special interest group

Richard Parker <sup>1,2</sup>, Michael Allison,<sup>3</sup> Seonaid Anderson,<sup>4</sup> Richard Aspinall,<sup>5</sup> Sara Bardell,<sup>6</sup> Vikram Bains,<sup>7</sup> Ryan Buchanan,<sup>8</sup> Lynsey Corless,<sup>9</sup> Ian Davidson,<sup>10</sup> Pauline Dundas,<sup>11</sup> Jeff Fernandez,<sup>12</sup> Ewan Forrest,<sup>13</sup> Erica Forster,<sup>1</sup> Dennis Freshwater,<sup>6</sup> Ruth Gailer,<sup>14</sup> Robert Goldin,<sup>15</sup> Vanessa Hebditch,<sup>16</sup> Steve Hood,<sup>17</sup> Arron Jones,<sup>18</sup> Victoria Lavers,<sup>19</sup> Deborah Lindsay,<sup>20</sup> James Maurice,<sup>21</sup> Joanne McDonagh,<sup>6</sup> Sarah Morgan,<sup>22</sup> Tania Nurun,<sup>9</sup> Christopher Oldroyd,<sup>3</sup> Elizabeth Oxley,<sup>23</sup> Sally Pannifex,<sup>24</sup> Graham Parsons,<sup>25</sup> Thomas Phillips,<sup>26</sup> Nicole Rainford,<sup>7</sup> Neil Rajoriya,<sup>6</sup> Paul Richardson,<sup>27</sup> J Ryan,<sup>28</sup> Joanne Sayer,<sup>29</sup> Mandy Smith,<sup>30</sup> Ankur Srivastava,<sup>21</sup> Emma Stennett,<sup>31</sup> Jennifer Towey <sup>6</sup>, Roya Vaziri,<sup>32</sup> Ian Webzell,<sup>7</sup> Andrew Wellstead,<sup>33</sup> Ashwin Dhanda,<sup>34</sup> Steven Masson <sup>35</sup>

**To cite:** Parker R, Allison M, Anderson S, *et al*. Quality standards for the management of alcohol-related liver disease: consensus recommendations from the British Association for the Study of the Liver and British Society of Gastroenterology ARLD special interest group. *BMJ Open Gastroenterol* 2023;**10**:e001221. doi:10.1136/bmjgast-2023-001221

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjgast-2023-001221>).

Received 28 July 2023  
Accepted 29 August 2023



© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

## Correspondence to

Dr Richard Parker;  
[richardparker@nhs.net](mailto:richardparker@nhs.net)

## ABSTRACT

**Objective** Alcohol-related liver disease (ALD) is the most common cause of liver-related ill health and liver-related deaths in the UK, and deaths from ALD have doubled in the last decade. The management of ALD requires treatment of both liver disease and alcohol use; this necessitates effective and constructive multidisciplinary working. To support this, we have developed quality standard recommendations for the management of ALD, based on evidence and consensus expert opinion, with the aim of improving patient care.

**Design** A multidisciplinary group of experts from the British Association for the Study of the Liver and British Society of Gastroenterology ALD Special Interest Group developed the quality standards, with input from the British Liver Trust and patient representatives.

**Results** The standards cover three broad themes: the recognition and diagnosis of people with ALD in primary care and the liver outpatient clinic; the management of acutely decompensated ALD including acute alcohol-related hepatitis and the posthospital care of people with advanced liver disease due to ALD. Draft quality standards were initially developed by smaller working groups and then an anonymous modified Delphi voting process was conducted by the entire group to assess the level of agreement with each statement. Statements were included when agreement was 85% or greater. Twenty-four quality standards were produced from this

## WHAT IS ALREADY KNOWN ON THIS TOPIC

⇒ Alcohol related liver disease (ALD) is a common cause of ill health and premature death. Variations in the care of liver disease across the UK

## WHAT THIS STUDY ADDS

⇒ These consensus recommendations from the BASL/BSG ArLD special interest group are intended to improve quality and reduce variation in the management of ALD

## HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

⇒ Recommendations for all aspects of management of ArLD are provided. To support implementation an audit tool is provided, based on the quality standards, and a template for patient information leaflet

process which support best practice. From the final list of statements, a smaller number of auditable key performance indicators were selected to allow services to benchmark their practice and an audit tool provided.

**Conclusion** It is hoped that services will review their practice against these recommendations and key performance indicators and institute service development where needed to improve the care of patients with ALD.

## INTRODUCTION

Alcohol-related liver disease (ALD) is the major driver of liver-related morbidity and mortality in the UK. In 2019–2020, there were 74 220 admissions to hospital in England due to ALD<sup>1</sup> and 13 269 premature deaths (before the age of 75) between 2017 and 2019.<sup>2</sup> National Health Service (NHS) care for people with significant ALD is a major cost to the NHS—£1.8 billion for inpatient care related to alcohol.<sup>3</sup> ALD is intrinsically linked to alcohol misuse as the primary cause of liver disease, and in addition to the health costs of ALD, alcohol misuse is also causally associated to over 200 medical conditions and societal harms such as accidents, absenteeism and crime.<sup>4</sup> ALD is part of a wider spectrum of alcohol-related disease, which is linked to important comorbidities particularly mental health issues, cardiovascular disease and cancer,<sup>5</sup> and to other determinants of well-being particularly deprivation.<sup>6</sup> Recent data from the Office for Health Improvement and Disparities have highlighted the worsening situation: in 2020, 5608 alcoholic liver deaths were recorded in England, a rise of almost 21% compared with 2019. The spike in alcoholic liver deaths in 2020 occurred after a 43% increase in alcoholic liver deaths between 2001 and 2019.<sup>7,8</sup> The same publication highlighted that alcohol specific deaths have risen significantly since the onset of the COVID-19 pandemic. The treatment of ALD, therefore, requires the input of multiple practitioners and multiple different agencies.

ALD remains difficult to treat: a survey of outcomes from hospitals in England, Wales and Northern Ireland by the national confidential enquiry into patient outcomes and death found that death rates for patients remained high between two iterations of the survey in 2011 and 2019, and evidence of significant variations in care between centres.<sup>9,10</sup> This is consistent with NHS England data showing marked differences in the rates of death from liver disease across the country<sup>11</sup> and the observation that people with advanced liver disease admitted to hospital in emergency are 7–8 times more likely to die than those admitted for stroke or heart attack.<sup>12</sup>

Treatment guidelines for ALD have been produced by specialist hepatology societies to support the medical treatment of ALD, and the National Institute for Health and Care Excellence has also produced guidance in the UK<sup>13,14</sup> although this document has not been updated since 2017. The purpose of this initiative is to develop standards that would support the delivery of high-quality care for patients with ALD and reduce variation between areas.

## METHODS

A group of experts from the British Association for the Study of the Liver (BASL) and British Society of Gastroenterology (BSG) ALD Special Interest Group (SIG) developed the recommendations. RP chaired the group. All members of the ALD SIG were invited to participate via email and those expressing an interest were included in the working group. Ultimately, the group included a

multidisciplinary team of 46 individuals from hepatology, psychiatry and psychology, dietetics, hepatology specialist nursing, pathology, primary care, pharmacy and addiction specialists.

The group was subdivided into three working groups that led the writing of draft recommendations for one of three parts of the document: (1) prehospital assessment and management of ALD (lead AS); (2) management of decompensated ALD in secondary care (lead JS) and (3) management after hospitalisation (lead RP). Each group produced a list of draft quality statements for the management of ALD diagnosis and management pathway to address within the standards document. The draft statements from three working parties were combined and an anonymous modified Delphi voting process was conducted individually by each member of the working group using an online survey tool to assess the level of agreement with each statement on a five-point scale (strongly disagree, disagree, neutral, agree or strongly agree). Members could abstain from questions that related to areas outside their usual clinical practice. After a round of voting, statements were redrafted if necessary through discussions via teleconference meetings. ‘Agreement’ was predefined when statements received a score of ‘strongly agree’ or ‘agree’, consensus predefined as when agreement was  $\geq 85\%$  after exclusion of abstentions. The result of this process produced a series of 24 recommendations (table 1). From this final list of statements, nine auditable key performance indicators (KPIs) were selected to allow services to benchmark their practice. The KPIs were chosen based on their potential to influence patient outcomes as well as being easily measurable. An audit tool based on these KPIs was produced to allow for assessment, benchmarking and monitoring of services.

## Patient and public partnership involvement

The quality standards group included a representative from the British Liver Trust (VH) and a patient representative (VL).

Terminology used in the document for ALD, and cirrhosis due to (ALD cirrhosis) is based on the most recent guideline document from the European society for the study of the liver, which recognised that use of the word ‘alcoholic’ is stigmatising and sought to move away from its use.<sup>15</sup> We have also used the term alcohol-related hepatitis (AH) to replace the previous term alcoholic hepatitis.

## QUALITY STANDARDS

### Prehospital assessment and management of ALD

People who are being asked about their alcohol use should have a validated alcohol questionnaire completed to identify any need for intervention

All patients in primary care should have an accurate recording of alcohol intake, which includes both quantity and frequency, updated at registration and

**Table 1** Summary of quality standards with level of agreement from the final round of the Delphi process

	Agreement	Responses
People who are being asked about their alcohol use should have a validated alcohol questionnaire completed to identify any need for intervention.	91%	29% strongly agree 62% agree 6% disagree 2% strongly disagree
1. Assessment of liver fibrosis should be a. Offered to people who drink hazardously (35 units/week in women, 50 units/week in men). b. Considered in people drinking alcohol in excess of maximum recommended levels (14 units/week) who have cofactors for liver disease (eg, obesity).	94%	55% strongly agree 39% agree 3% neutral 3% disagree
Assessment of hepatic fibrosis should be done using validated non-invasive liver fibrosis markers.	100%	78% strongly agree 22% agree
Patients identified at high risk of advanced fibrosis or cirrhosis should be offered referral for assessment by a gastroenterologist or hepatologist.	97%	74% strongly agree 24% agree 2% neutral
Patients presenting to hospital with liver disease should be screened for alcohol use disorder (AUD) and an estimation of typical no of units of alcohol per week recorded.	76%	38% strongly agree 44% agree 15% neutral 3% disagree
Patients admitted to hospital with ALD should be reviewed by a clinician trained in hepatology and the management of alcohol withdrawal within 24 hours of admission.	88%	49% strongly agree 39% agree 9% neutral 3% disagree
Patients admitted to hospital with ALD and AUD should be assessed by a specialist addiction practitioner during their admission and offered appropriate intervention and referral.	97%	68% strongly agree 29% agree 3% neutral
Alcohol withdrawal syndrome in patients with ALD with advanced liver disease, especially jaundice and/or encephalopathy, should be treated in a symptom-triggered fashion using a recognised symptom scoring system to avoid overuse of benzodiazepines.	91%	66% strongly agree 25% agree 9% neutral
It should be documented that patients have been advised that complete abstinence from alcohol is associated with better prognosis in ALD and that stopping alcohol entirely should be their goal.	89%	65% strongly agree 24% agree 12% neutral
Patients presenting with decompensated ALD or AH should be screened for infection.	100%	79% strongly agree 21% agree
All patients with decompensated ALD should have a nutritional assessment.	100%	85% strongly agree 15% agree
A plan for escalation of care in patients with ALD who develop acute-on-chronic liver failure (grades 2 or 3) should be clearly documented.	100%	78% strongly agree 22% agree
AH should be diagnosed in keeping with recognised clinical criteria, and patients suspected as having AH but who have confounding factors or do not fulfil all criteria should be considered for liver biopsy.	93%	59% strongly agree 35% agree 7% neutral
Patients with AH should have their prognosis assessed using a recognised prognostic scoring system (GAHS; MELD).	100%	77% strongly agree 23% agree
Corticosteroid treatment should be considered in patients with indicators of likely beneficial response (GAHS $\geq$ 9; MELD 21–51; NLR 5–8) and without infection.	85%	58% strongly agree 27% agree 15% neutral
Response to treatment with corticosteroids should be assessed after 7 days and corticosteroid treatment discontinued if there is no response.	86%	45% strongly agree 41% agree 10% neutral 3% disagree
Patients should be provided with clear, written information about their liver disease in a manner that they can understand before they leave hospital.	94%	77% strongly agree 18% agree 6% neutral
The date and time of follow-up appointments should be arranged with patients before they leave hospital.	88%	56% strongly agree 32% agree 9% neutral 3% disagree
Patients hospitalised with decompensated ALD or AH should be followed up by clinicians with specialist interest in hepatology within 6 weeks of discharge.	97%	63% strongly agree 34% agree 3% neutral

Continued

**Table 1** Continued

	Agreement	Responses
Patients with ALD with AUD should be offered community-based alcohol support after discharge from hospital.	94%	77% strongly agree 18% agree 6% neutral
Access to addiction specialists should be available, when indicated, for all patients with decompensated ALD after leaving hospital.	91%	61% strongly agree 30% agree 9% neutral
Medicines to support abstinence are beneficial and should be continued in primary care after being started in hospital or in alcohol treatment.	85%	50% strongly agree 35% agree 15% neutral
Patients with ALD with ongoing hepatic failure and a UKELD score greater than 49 should be considered for liver transplant referral if they are abstinent from alcohol.	87%	47% strongly agree 40% agree 7% neutral 7% disagree
Patients with ALD with an expected survival of less than 12 months should have their condition discussed with palliative care services.	91%	73% strongly agree 18% agree 6% neutral 3% disagree
AH, alcohol-related hepatitis; ALD, alcohol-related liver disease; GAHS, Glasgow Alcoholic Hepatitis Score; MELD, model for end-stage liver disease; UKELD, UK Model for End-Stage Liver Disease.		

opportunistic with a special focus on people who have an alcohol-related condition or who are at increased risk of harm from alcohol.<sup>16</sup> This is in line with the Public Health England initiative 'Making every contact count'.<sup>17</sup> Patients who drink above the recommended limits should be screened for alcohol use disorder (AUD) using a validated formal tool, for example alcohol use disorders identification test-C (AUDIT-C) or fast alcohol screening test (FAST). Those identified as drinking at higher risk or hazardous levels (35 units/week in women, 50 units/week in men) should receive feedback; a brief intervention and written information on their alcohol intake and how to address it.<sup>18</sup> Screening should occur at least annually for those deemed by clinical judgement to be at high risk of alcohol-related morbidity including (but not limited to) significant mental health problems and metabolic risk factors.

**Assessment of liver fibrosis should be offered to people who drink hazariously (35 units/week in women, 50 units/week in men) and considered in people drinking alcohol in excess of maximum recommended levels (14 units/week) who have cofactors for liver disease (eg, obesity)**

The risk of liver disease increases with alcohol intake.<sup>19</sup> When males and females are stratified according to sex, the 35 units per week limit remains a reliable cut-off for increased advanced fibrosis risk in women, but for men, risk did not increase significantly until consuming 50 or more units per week.<sup>20</sup> When considering the value of fibrosis assessment in those who drink above recommended limits but below thresholds currently considered most harmful (35 units/week women; 50 units/week men), thought should be given to the presence of other significant drivers of liver disease, most notably obesity and diabetes mellitus.<sup>21–22</sup> The risk between alcohol intake and obesity is multiplicative<sup>23</sup>

in patients with a body mass index (BMI) >30 kg/m<sup>2</sup> drinking above UK recommended limits, the risk of chronic liver disease is five times greater than a patient with a normal BMI,<sup>24</sup> and those with severe obesity (BMI >35 kg/m<sup>2</sup>) have double the risk of liver disease for any level of alcohol intake,<sup>25</sup> therefore, any intake over recommended limits in this group might prompt assessment of fibrosis. This approach is supported by clinical guidelines<sup>26</sup> but direct evidence for fibrosis testing at this threshold is lacking.

**Assessment of hepatic fibrosis should be done using validated non-invasive liver fibrosis markers**

Identification of advanced fibrosis offers the opportunity for earlier referral to specialist support, earlier intervention for both the liver disease and significant cofactors/drivers, while avoiding unnecessary investigation in those without liver disease—only 20% of those with AUD for example, will develop significant liver disease.<sup>27</sup> Standard liver blood tests do not exclude the presence of significant liver fibrosis<sup>28</sup> and specific non-invasive tests (NITs) for liver fibrosis should therefore be used. NITs for detection of liver fibrosis are widely available in UK practice and are included in UK guidelines on investigation for fibrosis in AUD in primary care.<sup>25</sup> However, the evidence base to support the use of specific tests, or to provide cut-off values to exclude advanced fibrosis, is considerably smaller than in non-alcohol-related fatty liver disease.<sup>29–31</sup> NITs that have been used in ALD include panels based on standard blood tests (alanine transaminase (ALT), aspartate transaminase (AST), platelets) such as fibrosis-4 (FIB-4) and AST to platelet ratio index (APRI), stand-alone specialist blood tests such as Enhanced Liver Fibrosis (ELF) test, and imaging techniques such as transient elastography (TE).<sup>32</sup>



**Patients identified at high risk of advanced fibrosis or cirrhosis should be offered referral for assessment by a gastroenterologist or hepatologist**

All patients identified at high risk of advanced fibrosis or cirrhosis after assessment with NIT should be referred to secondary care for further evaluation. Patients should be informed that an initial positive result highlights the need for further evaluation rather than confirming the presence of significant liver disease, given the modest positive predictive value with some NIT results. Those at low risk of advanced fibrosis or cirrhosis do not need referral to secondary care and can be managed safely in primary care or in community settings, with a focus on tailored lifestyle advice and access to support services, as no liver-specific treatments are indicated. In patients with metabolic or cardiovascular risk factors, managing these risk factors is important to reduce the risk of disease progression. Patients should be reassessed for liver fibrosis after 3–5 years if they continue to drink above recommended limits, using the tests described above.<sup>14 33</sup>

**Assessment and management of patients with acutely decompensated ALD cirrhosis or AH**

**Patients presenting to hospital with liver disease should be screened for AUD and an estimation of typical number of units of alcohol per week recorded**

Evaluation of alcohol intake is important for accurate diagnosis of liver disease and assessing the need for management of alcohol withdrawal. Screening for AUD should be done with validated tools such as AUDIT or its abbreviated forms such as AUDIT-C or FAST. Integrated care of liver disease and AUD requires the engagement of multiple practitioners to be effective and relies on prompt identification of patients who need care.<sup>34</sup> All those experiencing alcohol dependence should be monitored for the early signs of alcohol withdrawal (for example sweats, tremor, craving or agitation). Those typically drinking daily >15 units of alcohol are likely to experience moderate dependence. Those typically drinking daily >30 units of alcohol are likely to experience severe alcohol dependence and at greater risk of severe alcohol withdrawal symptoms.<sup>35</sup> In addition to the direct management of ALD, there is value in systematic screening for alcohol misuse in all patients admitted to hospital to better manage alcohol withdrawal and to identify those at risk of liver disease.<sup>36</sup>

**Patients admitted to hospital with ALD should be reviewed by a clinician trained in hepatology and the management of alcohol withdrawal within 24 hours of admission**

Early consultant review improves outcomes in hospital admissions.<sup>37</sup> In patients with acutely decompensated ALD who may also require management of alcohol withdrawal, this complexity requires early specialist review. Timely review by senior specialist clinicians

has been shown to reduce early mortality in patients admitted with decompensated liver disease.<sup>38</sup> All patients with decompensated ALD cirrhosis should have their initial management guided by the BSG/BASL Decompensated Cirrhosis Care Bundle.<sup>39</sup>

**Patients admitted to hospital with ALD and AUD should be assessed by a specialist addiction practitioner during their admission and offered appropriate intervention and referral**

Alcohol treatment improves outcomes in patients with ALD and should be considered an integral part of holistic treatment.<sup>40</sup> Harmful alcohol use in the context of liver disease needs to be clearly addressed by practitioners with appropriate experience and training.<sup>35</sup> Specialist addiction practitioners may come from a variety of backgrounds including but not limited to nursing or medical training; their expertise in managing initial withdrawal and subsequent alcohol treatment is key. The international classification of alcohol use disorders (international classification of diseases (ICD) -10 or ICD-11) identify harmful drinking and alcohol dependence as conditions requiring access to specialist treatment interventions, which have shown to be effective in helping individuals to reduce or stop their alcohol use.<sup>16</sup> In centres with an alcohol care team (ACT), assessment and management of alcohol use is best done as part of the comprehensive care that ACTs provide.<sup>41</sup> Treatment and onward referral after an initial alcohol detoxification are supported by other guidelines.<sup>35</sup>

**Alcohol withdrawal syndrome in patients with ALD with advanced liver disease, especially jaundice and/or encephalopathy, should be treated in a symptom-triggered fashion using a recognised symptom scoring system to avoid overuse of benzodiazepines**

Symptom triggered management should be the gold standard for management of withdrawal in patients with liver disease to avoid the risk of oversedation with benzodiazepines.<sup>13</sup> However, it is recognised that this requires a highly vigilant, trained workforce who are able to maintain the schedule of review and react accordingly. Where this gold standard cannot be delivered, a modification to the benzodiazepine type/dose may be required for high risk patients using fixed dose regimens.

Management of alcohol withdrawal syndrome in those with advanced liver disease should be undertaken in a symptom triggered fashion using recognised symptom scoring systems. These include the CIWA-Ar scale<sup>42</sup> and the Glasgow Modified Alcohol Withdrawal Score (GMAWS).<sup>43</sup> Metabolism of benzodiazepines is reduced in advanced liver disease with the risk of accumulation and subsequent oversedation and precipitation of hepatic encephalopathy if fixed dose regimens are used. However, staff should be competent in monitoring symptoms effectively and should have sufficient resources to allow them to do so frequently and safely. In those with decompensated liver disease,

especially severe jaundice and/or encephalopathy, modification to the benzodiazepine regimen may be required. This might include a reduction in the dose of benzodiazepine triggered by symptoms or perhaps the use of shorter acting benzodiazepines such as oxazepam or lorazepam.<sup>16</sup>

**It should be documented that patients have been advised that complete abstinence from alcohol is associated with better prognosis in ALD and that stopping alcohol entirely should be their goal**

It is clear that abstinence from alcohol is associated with better outcomes compared with people who continue to drink.<sup>44 45</sup> Alcohol abstinence is also a prerequisite for consideration of liver transplantation should this become necessary. All patients with decompensated ALD cirrhosis including AH should therefore be advised that complete abstinence from alcohol should be the target of treatment. This may be an overwhelming goal and patients should be supported to work towards this target to avoid alienating them from treatment. This should be documented in patient notes, records and on electronic portals where possible with access for all clinicians involved in going care. A record of this advice should also be given to patients and general practitioners (usually via a discharge letter).

**Patients presenting with decompensated ALD or AH should be screened for infection**

Alcohol consumption increases the risk of infection, and infection is a common cause of decompensated liver disease.<sup>46</sup> Infection may be present at presentation or occur during admission particularly in AH, and is associated with poorer outcomes.<sup>47</sup> This advice regarding screening for infection is consistent with the BSG/BASL decompensated cirrhosis bundle. Monitoring for signs of infection should continue throughout an admission with particular vigilance in patients with AH who are treated with corticosteroids where incident infection is associated with poor outcomes.<sup>48</sup> It is important to note that standard laboratory markers of infection for example, leukocytosis or elevated C reactive protein may be part of the syndrome of AH and not necessarily reflect active infection; these should be interpreted with caution and additional evidence for active infection should be sought. This said, given the adverse outcome with infection a low threshold for prescribing antibiotics is appropriate. The possibility of fungal infection should be considered: this may be more frequent in AH and associated with poor outcomes.<sup>49</sup> Prophylactic antibiotic treatment in AH has been shown not to improve mortality or morbidity in AH and should not replace careful monitoring for signs of infection.<sup>50</sup>

**All patients with decompensated ALD should have a nutritional assessment**

Malnutrition (deficiencies in individual nutrients), sarcopenia (a reduction in muscle mass, strength and/or

function<sup>51</sup>) and frailty (a clinical state of decreased physiological reserve and increased vulnerability to health stressors, predisposing individuals to adverse clinical outcomes<sup>52</sup>) are common in end-stage liver disease and in patients with AH.<sup>53</sup> Malnutrition risk can be assessed using subjective global assessment; however, there is risk of underestimation so caution is advised. Gold standard assessment of sarcopenia is through use of CT with analysis of Skeletal Muscle Index, however, if inaccessible then bedside measures through obtaining mid-arm muscle circumference are advised<sup>54</sup>; if malnutrition risk or sarcopenia/frailty are identified, patients should be seen by a dietician with specialist training and experience in hepatology.<sup>55</sup> Patients' food security should be considered when developing dietary plans for the post-discharge period and referral to social care made when appropriate.

Enteral nutrition via nasogastric tube has often been considered part of the treatment of patients with AH. More recent trial data failed to show an advantage of NG feeding over standard care but it was clear that adequate caloric intake was associated with better outcomes<sup>56</sup>; close attention should be paid to dietary intake.

**A plan for escalation of care in patients with ALD who develop acute-on-chronic liver failure (grades 2 or 3) should be clearly documented**

Escalation decisions should be made on a patient-by-patient basis. Escalation decisions for patients with ALD must take account of factors such as pre-admission patient function, comorbidity, frailty, transplant eligibility and severity of liver disease; patient preferences must also be taken into account. Alcohol remains the the most common aetiology of acute on chronic liver failure (ACLF) in the UK<sup>57</sup>; ACLF grade and the need for renal replacement therapy are independently associated with 28-day mortality. Without transplantation, 28-day mortality falls between 18%–25% in grade 1 and 68%–89% grade 3.<sup>58</sup> With such high mortality early decision making and assessing appropriate candidacy is vital. A national cohort study of patients with ALD from the Scottish ITU registry confirmed a high mortality among this cohort, along with a significant long-term burden on subsequent healthcare resource utilisation.<sup>59</sup> Those admitted to ITU with ALD experience higher readmission rates, with more days in hospital and higher consequent costs, compared with ITU patients with other severe comorbidities. Therefore, careful liaison and shared decision-making about escalation, between the medical and ITU teams, alongside patients and their families is required. Escalation to intensive care units is often considered in the context of 'first presentation' of liver disease and the potential for patients to be a candidate for liver transplantation. Neither of these factors are known to be relevant to short term outcomes and may not be appropriate to inform care planning.

## Assessment and management of AH

AH should be diagnosed in keeping with recognised clinical criteria; patients suspected as having AH but who have confounding factors or do not fulfil all criteria should be considered for liver biopsy

The diagnosis of AH has been controversial in the past. Previously it had been argued that a diagnosis of AH could only be made with confirmatory histology. For most AH patients this would require a transjugular liver biopsy the availability of which is variable and for which there is a technical failure rate of 3.2% and a diagnostically suboptimal sample obtained in up to 12%.<sup>60</sup> In addition, histological features of steatohepatitis may be present in patients with chronic liver disease without an acute illness.<sup>61 62</sup> AH is primarily a clinical diagnosis with recent onset jaundice being a cardinal feature. In order to standardise the clinical features of AH the National Institute on Alcohol Abuse and Alcoholism has published criteria which have received general acceptance.<sup>63</sup> Onset of jaundice (serum bilirubin >3 mg/dL or 50 µmol/L) within 8 weeks and excessive alcohol consumption within 60 days of presentation are key. The liver biochemistry should be compatible with AH with a raised AST, an AST-to-ALT ratio of >1.5 and neither value >400 IU/L. Fulfilment of these criteria without any confounding factors equates to a diagnosis of 'probable' AH which suffices for most clinical situations. A 'definite' diagnosis of AH is when these clinical features are fulfilled and there is additional confirmatory histology. However, if confounding factors are present then the level of diagnostic certainty falls to 'possible' AH in which case additional confirmatory histology would be recommended to make the diagnosis. Examples of such factors would be incompatible liver biochemistry, possible ischaemic liver injury (secondary to hypotension or cocaine use within 7 days), possible metabolic liver injury (Wilson's disease), possible drug induced liver injury (suspect drug within 30 days of jaundice) and uncertainty regarding timing of onset of jaundice. Liver biopsy rarely changes the diagnosis in individuals who meet the clinical criteria or history of alcohol excess.<sup>64</sup>

It should be noted that while these criteria have been accepted and form part of guidelines,<sup>65</sup> the threshold of serum bilirubin recommended is lower than that of most historical clinical studies of AH. These have tended to use a threshold of 80 µmol/L (4.7 mg/dL) or more. Therefore, the applicability of previous clinical studies to those with lower levels of serum bilirubin is unclear.

Patients with AH should have their prognosis assessed using a recognised prognostic scoring system (Glasgow Alcoholic Hepatitis Score; model for end-stage liver disease)

Estimation of likely outcome from AH is important to identify those at greatest benefit from intervention as well as to allow informed discussion with the patient and their families regarding their expectations. 'Static' scores are those derived from variables available at a single point in time. The discriminant function (DF) has been used for

many years with a threshold  $\geq 32$  identifying those with severe disease.<sup>66 67</sup> However, concerns have been raised regarding the reliability of the DF as it uses the absolute value of prothrombin time rather than a ratio metric value such as the international normalised ratio (INR).<sup>68</sup> Alternative scores have been proposed including the Glasgow Alcoholic Hepatitis Score (GAHS, and also its modified version, mGAHS using the neutrophil-to-lymphocyte ratio (NLR)),<sup>69 70</sup> the age, bilirubin, INR and creatinine score<sup>71</sup> and the model for end-stage liver disease (usually the pre-2016 UNOS variation of MELD, but also the modified MELD-Na version)<sup>72</sup> score. Recent studies have shown that these more recent scores are superior to the original DF.<sup>72 73</sup> In international guidelines, the GAHS and MELD have been most commonly cited as indicators of prognosis in clinical practice.<sup>15 74</sup> In the STOPAH trial, the 90-day mortality of AH with consistently low scores was: GAHS <9 110.2%; MELD <25 140.4%. A 90-day mortality in those with high scores (excluding those presenting with sepsis and/or gastrointestinal bleeding) was: GAHS  $\geq 9$  380.4%; MELD  $\geq 25$  390.6%.<sup>73</sup>

Corticosteroid treatment should be considered in patients with indicators of likely beneficial response (GAHS  $\geq 9$ ; MELD 21–51; NLR 5–8) and without infection

Corticosteroids are the only currently recommended pharmacological treatment. In the STOPAH trial, corticosteroids were associated with a modest improvement in mortality at 28 days, though the improvement did not reach conventional statistical significance (OR 0.72, 95% CI 0.52 to 1.01,  $p=0.06$ ) and any benefit was not sustained at 90 days.<sup>75</sup> Subsequent meta-analysis supports the use of corticosteroids in selected patients to improve 28-day survival (HR 0.64, 95% CI 0.48 to 0.86,  $p=0.003$ ), but also confirmed longer-term benefit was lost at 6 months.<sup>76</sup> Importantly, the rate of infection is significantly higher in patients with AH treated with steroids and in those who develop infection with AH, steroids increase 90-day mortality (OR 2.26, 95% CI 1.41 to 4.30,  $p=0.002$ ).<sup>48 77</sup> Therefore, if steroids are to be used, they should be used cautiously after careful screening and aggressive treatment of infection at baseline. Furthermore, if a significant spontaneous decrease in bilirubin is observed after 7 days ( $<0.9 \times$  baseline bilirubin), 28-day survival may not be improved with steroids, and in this group the risks of steroid treatment could outweigh the benefits.<sup>78</sup> The standard dose of prednisolone is 40 mg daily, this is continued for 28 days in responders and then discontinued without weaning.

Response to treatment with corticosteroids should be assessed after 7 days and corticosteroid treatment discontinued if there is no response

Dynamic prognostic scores, such as the Lille score,<sup>79</sup> predict mortality based on a reduction in bilirubin level in patients who have received corticosteroid therapy. The Lille score combines several variables, including age, renal function, albumin, prothrombin time and bilirubin



at baseline and 7 days. A non-response to corticosteroids is defined as a Lille score  $\geq 0.45$ , and this is associated with a much poorer survival at 6 months compared with responders (25% vs 85%) and it is recommended that steroids are discontinued in this group. Calculating the Lille score at day four may give similar diagnostic accuracy in predicting 28-day and 90-day mortality compared with the day seven score, limiting unnecessary steroid exposure and reducing the risk of infection, but this requires further validation.<sup>80</sup>

### Posthospital management of ALD

**Patients should be provided with clear, written information about their liver disease in a manner that they can understand before they leave hospital**

Patients should be supported to understand their health in the transition from hospital to the community.<sup>81</sup> Written information should include details about their liver disease, the reasons for hospital admission, treatment received and plans for follow-up including advice about abstinence and sources of alcohol support. Written information should take into account literacy level as the median UK reading age is 9 years old. A template information leaflet for decompensated alcohol-related cirrhosis is provided in the supporting information. The British Liver Trust also offers a range of resources designed to be accessible for patients (<https://britishlivertrust.org.uk/>). Where appropriate and with patients' consent, information should be shared with people significant or important to patients.

**The date and time of follow-up appointments should be arranged with patients before they leave hospital**

Patients who attend outpatient hepatology outpatient clinics have been shown in retrospective reviews to have improved survival.<sup>82 83</sup> Early follow-up also facilitates discharge and may help prevent readmission, both of which can reduce pressure on inpatient services. Necessary investigations should where possible be completed during an index admission to reduce journeys and costs to patients. Measures to allow patients to lead follow-up arrangements by picking dates or times of appointments may also support engagement. Outpatient hepatology clinics have traditionally been delivered in hospitals, but could be placed in other centres which may improve attendance.

**Patients hospitalised with decompensated ALD or AH should be followed up by clinicians with specialist interest in hepatology within 6 weeks of discharge**

In clinical studies, follow-up has been difficult.<sup>84</sup> Special efforts to engage patients with follow-up may be needed in excess of other patient groups. These efforts should ideally ensure that further engagement is made as easy as possible and, importantly, continually informed by patient experience. Additional efforts to engage patients through for example peer support, dedicated specialist nursing staff and using text messages have shown to

improve attendance in people with hepatitis C<sup>85 86</sup>; similar initiatives may also be valuable in ALD.

**Patients with ALD with AUD should be offered community-based alcohol support after discharge from hospital**

When accepted, contact with alcohol services should be established before leaving hospital. Team working including joint clinics between hepatology and alcohol practitioners may allow alcohol support to be offered to those declining community services. The management plan must address alcohol reduction and cessation.<sup>15</sup> Patients who have initially declined referral to alcohol services should be able to access this support if they change their minds. Motivational enhancement work by hepatology staff can help and reinforce the need for links between the services. Other substance use should also be addressed in addition to alcohol use. For example, there is a growing body of evidence that cigarette smoking is associated with increased progression of fibrosis and development of hepatocellular carcinoma. It is recommended that all patients with ALD be offered pharmacological and psychological smoking cessation support, particularly those being considered for liver transplantation.<sup>87 88</sup>

**Access to addiction specialists should be available, when indicated, for all patients with decompensated ALD after leaving hospital**

Addiction specialist psychiatric specialists improve outcomes for people with advanced disease, for example, through reduced relapse rates after transplantation.<sup>89</sup> Most literature around the impact of this service is in the context of transplantation; the quality standard development group was clear that access to specialist addiction psychiatrists should not be reserved for such a highly selected group of patients. Good practice would include access to specialist psychiatrists for all patients.

**Medicines to support abstinence are beneficial and should be continued in primary care after being started in hospital or in alcohol treatment**

A recent systematic review has demonstrated the benefits of either integrated or colocated addiction therapy for patients with ALD.<sup>90</sup> Compared with standard of care patients who received addiction treatment were less likely to suffer liver decompensation, require readmission (within 30 days) and had a lower risk of mortality. A similar pattern is seen in the use of relapse prevention medications: patients with cirrhosis were less likely to suffer hepatic decompensation.<sup>40</sup> This positive effect remained even when treatment was started after a diagnosis of cirrhosis indicating the ongoing value of treatment of AUD. Indications and medication doses should be reviewed in light of changes in liver function and patients' needs, and their engagement with addiction services. Disulfiram is associated with potentially serious hepatotoxicity and should be avoided in patients with advanced liver disease; its use in earlier disease may be

safe with careful monitoring. Acamprosate is renally excreted and baclofen has evidence of safety and efficacy in cirrhosis,<sup>91 92</sup> these agents may be preferred in cirrhosis. The use of relapse prevention medications is covered in other national UK guidance currently in development.

**Patients with ALD with ongoing hepatic failure and a UK Model for end-stage liver disease score greater than 49 should be considered for liver transplant referral if they are abstinent from alcohol**

Liver transplant is an effective treatment option for selected patients with decompensated alcohol-related cirrhosis.<sup>93</sup> The development of decompensated disease (ascites, oedema, encephalopathy, bleeding) should prompt the consideration of transplant referral. The UK Liver Advisory Group<sup>94</sup> have recently provided updated guidelines for transplant assessment and referral which are specific to patients with alcohol-related disease. These supplement existing national criteria and guidelines for liver transplant.<sup>95 96</sup> Early discussion with a transplant centre should be considered in all cases, particularly where patients have demonstrated at least 3 months of alcohol abstinence or even earlier if they show good engagement with addiction services and an assessment may be prolonged because of medical complexity.<sup>95</sup> Referral should not be based on the 'rule of thumb' of 6 months abstinence before referral to a transplant unit.

The UK Model for End-Stage Liver Disease (UKELD) score is a compound measure of liver function intended to guide transplant candidacy in the UK. UKELD underpins minimal listing criteria for the elective waitlist. This is derived from the patient's serum sodium, creatinine, bilirubin and INR.<sup>97</sup> In chronic liver disease (CLD), a UKELD score  $\geq 49$  indicates survival advantage for LT over conservative management in patients with irreversible decompensation. A UKELD score of 49 is the equipoise at which the predicted 1-year mortality without liver transplantation (9%) matches that expected after liver transplantation, and is therefore the threshold of minimum listing criteria for elective liver transplantation in those with irreversible decompensation in the UK.<sup>95</sup> Calculators for the UKELD score are available on the NHS Blood and Transplant website.

**Patients with ALD with an expected survival of less than 12 months should have their condition discussed with palliative care services**

Core palliative care is best delivered by the hepatology team in parallel with active disease management. This includes ensuring that discussions about disease trajectory and advanced care planning occur alongside active management of disease complications such as transplantation: one should not preclude the other.<sup>98</sup> Patients should be given the opportunity to be introduced to the palliative and supportive care team in hospital and a discussion in a suitable environment offered. If this is declined by the patient an opportunity to meet with the

team postdischarge to discuss advanced care planning should be offered.

Planned admissions for paracentesis as a day case are well evidenced to improve patient and carer satisfaction. This can be done by a nurse-led service with the support of the medical team. The opportunity to use a multidisciplinary approach can be used including dietitian input, addictions specialist/ACT and introduction to supportive and palliative care. It allows continuity and advanced care planning. For patients enrolled in day case services, improvements in outcomes correlated with the proportion of large volume paracentesis procedures done in a day case (vs unplanned) setting. There are significant cost savings to planned admissions.<sup>99</sup>

Long-term indwelling ascitic drains (LTAD) drains could also be considered on a case-to-case basis in patients with refractory ascites who are not under consideration for/listed for liver transplantation or transjugular intrahepatic portosystemic shunt (TIPSS). The risks of indwelling drains need to be carefully weighed and decisions for LTAD insertion should be made by a multidisciplinary team.<sup>100</sup>

## RECOMMENDATIONS FOR FUTURE RESEARCH, AUDIT AND SERVICE DEVELOPMENT

### Earlier detection of liver disease

It has been clear for some time that people with ALD are more likely to present late, with more advanced disease, compared with those with other causes of liver disease.<sup>101</sup> Recent initiatives in the UK have sought to find cases of disease at an earlier stage, for example, by the use of TE in people at risk of disease.<sup>102</sup> Understanding the value of this approach and the best way to implement methods of early detection will be essential for future service development.

### NITs for risk assessment in ALD

There is a clear unmet need to understand which NITs have the greatest utility for the risk assessment of ALD, and the cut-off levels which are best suited to confirming or excluding advanced fibrosis and risk of liver-related ill health. The optimal timeframe for repeat testing in patients at low risk of disease at baseline also needs to be established.

### Treatments and services for decompensated disease and AH

There have been no new treatments for decompensated ALD cirrhosis or AH since the introduction of corticosteroids several decades ago. Given the increasing number of admissions and deaths from ALD this is an important gap in our ability to care for patients and improve outcomes. Research into new treatments is necessary. Research into new methods of delivering care is also required for example the value of 'hub and spoke' relationships between secondary and tertiary care for improved access to specialist services, and delivery of hepatology services outside of traditional hospital settings such as outpatient clinics based in community or alcohol treatment centres.

Innovative research methods may be necessary to effectively deliver research in this cohort since concomitant health and social issues may reduce the value of traditional trial methods.

## CONCLUSION

ALD continues to be a major cause of ill health and premature mortality. These quality standards should be used alongside clinical guidelines from other organisations and local protocols to direct the treatment of patients with ALD. Despite the prevalence of ALD, research has historically lagged behind other less common causes of liver disease. As such there remains some uncertainty about the best management of ALD. We have highlighted particular aspects of the care of patients with ALD that we feel are most in need of examination (online supplemental file 1). As information from clinical audit and research emerges, it is likely that these standards will need to be updated so that they remain relevant. It is hoped that these evidence-based quality standards will improve the care of ALD and reduce variation between units.

## Author affiliations

- <sup>1</sup>Leeds Liver Unit, Leeds Teaching Hospitals NHS Trust, Leeds, UK
- <sup>2</sup>Leeds Institute of Medical Research at St James's University Hospital, University of Leeds, Leeds, UK
- <sup>3</sup>Cambridge Liver Unit, Addenbrooke's Hospital, Cambridge, UK
- <sup>4</sup>Angus Integrated Drug and Alcohol Recovery Service (AIDARS), Ninewells Hospital and Medical School, Dundee, UK
- <sup>5</sup>Gastroenterology & Hepatology, Portsmouth Hospitals NHS Trust, Portsmouth, UK
- <sup>6</sup>Birmingham Liver Services Unit, University Hospitals Birmingham NHS Foundation Trust, Birmingham, UK
- <sup>7</sup>Liver Transplant Unit, King's College Hospital NHS Foundation Trust, London, UK
- <sup>8</sup>Faculty of Medicine, University of Southampton, Southampton, UK
- <sup>9</sup>Department of Gastroenterology, Hepatology and Endoscopy, Hull University Teaching Hospitals NHS Trust, Hull, UK
- <sup>10</sup>NHS Fife Addiction Services, NHS Fife, Kirkcaldy, UK
- <sup>11</sup>Peter Brunt Centre, Aberdeen Royal Infirmary, NHS Grampian, Aberdeen, UK
- <sup>12</sup>Alcohol and Drug Liaison, Royal Free London NHS Foundation Trust, London, UK
- <sup>13</sup>Dept of Gastroenterology, Glasgow Royal Infirmary, Glasgow, UK
- <sup>14</sup>Islington Primary Care Federation, London, UK
- <sup>15</sup>Department of Digestive Diseases, Department of Medicine, Imperial College London, London, UK
- <sup>16</sup>British Liver Trust, Ringwood, UK
- <sup>17</sup>Digestive Diseases Unit, Liverpool University Hospitals NHS Foundation Trust, Liverpool, UK
- <sup>18</sup>Pharmacy, Barts and The London NHS Trust, London, UK
- <sup>19</sup>Independent researcher, Plymouth, UK
- <sup>20</sup>Alcohol Care Team, East Lancashire Hospitals NHS Trust, Blackburn, UK
- <sup>21</sup>Gastroenterology and hepatology, North Bristol NHS Trust, Westbury on Trym, UK
- <sup>22</sup>Hampstead Group Practice, London, UK
- <sup>23</sup>St Martins Practice, Leeds, UK
- <sup>24</sup>Hepatology, St George's Healthcare NHS Trust, London, UK
- <sup>25</sup>Hepatology, Plymouth Hospitals NHS Trust, Plymouth, UK
- <sup>26</sup>Faculty of Health Sciences, University of Hull, Hull, UK
- <sup>27</sup>Gastroenterology and Hepatology, Royal Liverpool and Broadgreen University Hospitals NHS Trust, Liverpool, UK
- <sup>28</sup>Sheila Sherlock Liver Centre, Royal Free London NHS Foundation Trust, London, UK
- <sup>29</sup>Gastroenterology, Doncaster and Bassetlaw Hospitals NHS Foundation Trust, Doncaster, UK
- <sup>30</sup>Alcohol care team, Southport and Ormskirk Hospital NHS Trust, Southport, UK
- <sup>31</sup>Gastroenterology, Guy's and St Thomas' Hospitals NHS Trust, London, UK
- <sup>32</sup>Humankind, Durham, UK

<sup>33</sup>Gastroenterology, University Hospitals Dorset NHS Foundation Trust, Poole, UK

<sup>34</sup>Faculty of health, University of Plymouth, Plymouth, UK

<sup>35</sup>Liver unit, The Newcastle upon Tyne Hospitals NHS Foundation Trust, Newcastle, UK

**Twitter** Jennifer Towey @liver\_dietitian

**Contributors** RP led the project, chaired meetings of the whole group and one of the working groups, coordinated Delphi process and drafted the manuscript. MA, SA, RA, SB, VB, RB, LC, ID, AD, PD, JF, EForrest, EForster, DF, RGailor, RGoldin, SH, AJ, DL, JMaurice, JMcDonagh, SMorgan, TN, CO, EO, SP, GP, TP, NRainford, NRajoriya, PR, JR, ES, JT, RV, IW, AW and SMasson attended meetings, contributed to Delphi process, drafted quality standards and reviewed draft versions of the manuscript. VH reviewed the draft manuscript. VL contributed to the Delphi process and reviewed the draft manuscript. JS and AS chaired meetings of working groups, coordinated Delphi process and drafted the manuscript.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** RP: research support from NIHR and Leeds Hospital Charity. Consulting fees from Durect. Fees for speaking from Norgine Pharmaceuticals. Advisory board fees from Novo Nordisk. LC: consulting fees from Novo Nordisk relating to non-alcohol-related fatty liver disease. TP: research grant support from NIHR, Society for the Study of Addiction, Office of Police and Crime Commissioner. SMorgan: research support from NIHR. Consulting fees from Norgine Pharmaceuticals, Payment or honoraria from Sandoz UK and Dr Falk. LEadership roles with British Society of Gastroenterology, Alcohol Health Alliance, Medical Council on Alcohol. These authors declare no conflicts of interest: SA, MA, RA, SB, VB, RB, ID, AD, PD, RGailor, EForrest, EForster, RGoldin, VH, AJ, VL, DL, JMcDonagh, JMaurice, SMasson, TN, EO, GP, NRajoriya, NRainford, PR, JR, JS, MS, AS, ES, JT, RV, IW and AW.

**Patient consent for publication** Not applicable.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** No data are available.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

## ORCID iDs

Richard Parker <http://orcid.org/0000-0003-4888-8670>

Jennifer Towey <http://orcid.org/0000-0002-7320-5328>

Steven Masson <http://orcid.org/0000-0003-1041-9844>

## REFERENCES

- 1 NHS Digital. Statistics on alcohol. England; 2021.
- 2 Fingertips public health data (office for health improvement and disparities). In: *Local Alcohol Profiles for England*. Available: <https://fingertips.phe.org.uk/profile/local-alcohol-profiles/data#page/1/gid/1938132832/pat/159/par/K02000001/ati/15/are/E92000001/yr/1/cid/4/tbm/1> [accessed 20 Oct 2022].
- 3 Williams R, Alexander G, Armstrong I, et al. Disease burden and costs from excess alcohol consumption, obesity, and viral hepatitis: fourth report of the lancet standing Commission on liver disease in the UK. *The Lancet* 2018;391:1097–107.
- 4 Institute for Alcohol Studies. *The Costs of Alcohol to Society*. IAS Briefing, 2020.



- 5 Theodoreson MD, Aithal GP, Allison M, *et al.* Extra-hepatic morbidity and mortality in alcohol-related liver disease: systematic review and meta-analysis. *Liver Int* 2023;43:763–72.
- 6 Major JM, Sargent JD, Graubard BI, *et al.* Local geographic variation in chronic liver disease and hepatocellular carcinoma: contributions of socioeconomic deprivation, alcohol retail outlets, and lifestyle. *Ann Epidemiol* 2014;24:S1047–2797(13)00444–4:104–10..
- 7 Office for National Statistics. Home people, population and community births, deaths and marriages deaths alcohol-specific deaths in the UK alcohol-specific deaths in the UK: registered in 2021. 2021.
- 8 Public Health England. Research and analysis monitoring alcohol consumption and harm during the COVID-19 pandemic: summary. 2021.
- 9 National Confidential Enquiry into Patient Outcome and Death. Remeasuring the units an update on the Organisation of alcohol-related liver disease services. 2022.
- 10 Allison MED, Verne J, Bernal W, *et al.* Deaths from alcohol-related liver disease in the UK: an escalating tragedy. *Lancet* 2023;401:S0140-6736(22)02583-1:418–20..
- 11 Public Health England. The 2ND Atlas of variation in risk factors and Healthcare for liver disease in England. 2017.
- 12 Williams R, Alessi C, Alexander G, *et al.* New dimensions for hospital services and early detection of disease: a review from the lancet Commission into liver disease in the UK. *Lancet* 2021;397:S0140-6736(20)32396-5:1770–80..
- 13 Alcohol-use disorders: diagnosis and management of physical complications. London: National Institute for Health and Care Excellence (NICE),
- 14 National Guideline Centre (UK). *Cirrhosis in Over 16s: Assessment and Management*. London: National Institute for Health and Care Excellence (NICE),
- 15 Thursz M, Gual A, Lackner C, *et al.* EASL clinical practice guidelines: management of alcohol-related liver disease. *Journal of Hepatology* 2018;69:154–81.
- 16 National Collaborating Centre for Mental Health (Great Britain). Alcohol use disorders: the NICE guideline on the diagnosis. *Assessment and Management of Harmful Drinking and Alcohol Dependence RCPsych Publications* 2011.
- 17 Meade O, O'Brien M, Mc Sharry J, *et al.* Enhancing the implementation of the making every contact count brief behavioural intervention programme in Ireland: protocol for the making MECC work research programme. *HRB Open Res* 2022;5:6.
- 18 Kaner E, Bland M, Cassidy P, *et al.* Effectiveness of screening and brief alcohol intervention in primary care (SIPS trial): pragmatic cluster randomised controlled trial. *BMJ* 2013;346:e8501.
- 19 Llamas-Falcón L, Probst C, Buckley C, *et al.* Sex-specific association between alcohol consumption and liver cirrhosis: an updated systematic review and meta-analysis. *Front Gastroenterol (Lausanne)* 2022;1:1005729.
- 20 Rhodes FA, Cococcia S, Patel P, *et al.* Is there scope to improve the selection of patients with alcohol-related liver disease for referral to secondary care? A retrospective analysis of primary care referrals to a UK liver centre, incorporating simple blood tests. *BMJ Open* 2021;11:e047786.
- 21 UK Chief Medical Officers. Alcohol consumption: advice on low risk drinking. 2016.
- 22 Whitfield JB, Masson S, Liangpunsakul S, *et al.* Obesity, diabetes, coffee, tea, and Cannabis use alter risk for alcohol-related cirrhosis in 2 large cohorts of high-risk drinkers. *Am J Gastroenterol* 2021;116:106–15.
- 23 Inan-Eroglu E, Huang B-H, Ahmadi MN, *et al.* Joint associations of Adiposity and alcohol consumption with liver disease-related morbidity and mortality risk: findings from the UK Biobank. *Eur J Clin Nutr* 2022;76:74–83.
- 24 Glyn-Owen K, Böhning D, Parkes J, *et al.* The combined effect of alcohol and body mass index on risk of chronic liver disease: A systematic review and meta-analysis of cohort studies. *Liver Int* 2021;41:1216–26.
- 25 Newsome PN, Cramb R, Davison SM, *et al.* Guidelines on the management of abnormal liver blood tests. *Gut* 2018;67:6–19.
- 26 European Association for the Study of the Liver. Electronic address: Easloffic@Easloffic.EU, clinical practice guideline panel, chair; EASL governing board representative; panel members: EASL clinical practice guidelines on non-invasive tests for evaluation of liver disease severity and prognosis - 2021 update. *J Hepatol* 2021;75:659–89.
- 27 Parker R, Aithal GP, Becker U, *et al.* Natural history of histologically proven alcohol-related liver disease: A systematic review. *J Hepatol* 2019;71:S0168-8278(19)30306-X:586–93..
- 28 Rhodes F, Cococcia S, Panovska-Griffiths J, *et al.* Uncovering unsuspected advanced liver fibrosis in patients referred to alcohol nurse specialists using the ELF test. *BMC Gastroenterol* 2021;21:143.
- 29 Moreno C, Mueller S, Szabo G. Non-invasive diagnosis and biomarkers in alcohol-related liver disease. *J Hepatol* 2019;70:S0168-8278(18)32576-5:273–83..
- 30 Rhodes FA, Tremblay P, Panovska-Griffiths J, *et al.* Systematic review: investigating the Prognostic performance of four non-invasive tests in alcohol-related liver disease. *J Gastroenterol Hepatol* 2021;36:1435–49.
- 31 Hinkson A, Lally H, Gibson H, *et al.* Meta-analysis: enhanced liver fibrosis test to identify hepatic fibrosis in chronic liver diseases. *Aliment Pharmacol Ther* 2023;57:750–62.
- 32 Pavlov CS, Casazza G, Nikolova D, *et al.* Systematic review with meta-analysis: diagnostic accuracy of transient Elastography for staging of fibrosis in people with alcoholic liver disease. *Aliment Pharmacol Ther* 2016;43:575–85.
- 33 Rasmussen DN, Thiele M, Johansen S, *et al.* Prognostic performance of 7 biomarkers compared to liver biopsy in early alcohol-related liver disease. *J Hepatol* 2021;75:S0168-8278(21)00411-6:1017–25..
- 34 Winder GS, Fernandez AC, Mellinger JL. Integrated care of alcohol-related liver disease. *J Clin Exp Hepatol* 2022;12:1069–82.
- 35 Nice. Diagnosis, assessment and management of harmful drinking and alcohol dependence (CG 115). In: *National Institute for Health and Clinical Excellence (NICE)*. London, 2011.
- 36 Westwood G, Meredith P, Atkins S, *et al.* Universal screening for alcohol misuse in acute medical admissions is feasible and identifies patients at high risk of liver disease. *J Hepatol* 2017;67:S0168-8278(17)31997-9:559–67..
- 37 Bell D, Lambourne A, Percival F, *et al.* Consultant input in acute medical admissions and patient outcomes in hospitals in England: a multivariate analysis. *PLoS One* 2013;8:e61476.
- 38 Roberts SE, John A, Brown J, *et al.* Early and late mortality following unscheduled admissions for severe liver disease across England and Wales. *Aliment Pharmacol Ther* 2019;49:1334–45.
- 39 Dyson JK, Rajasekhar P, Wetten A, *et al.* "Implementation of a "care bundle" improves the management of patients admitted to hospital with decompensated cirrhosis". *Aliment Pharmacol Ther* 2016;44:1030–8.
- 40 Vannier AGL, Shay JES, Fomin V, *et al.* Incidence and progression of alcohol-associated liver disease after medical therapy for alcohol use disorder. *JAMA Netw Open* 2022;5:e2213014.
- 41 Moriarty KJ. Alcohol care teams: where are we now *Frontline Gastroenterol* 2020;11:293–302.
- 42 Sullivan JT, Sykora K, Schneiderman J, *et al.* Assessment of alcohol withdrawal: the revised clinical Institute withdrawal assessment for alcohol scale (CIWA-Ar). *Addiction* 1989;84:1353–7. 10.1111/j.1360-0443.1989.tb00737.x Available: <http://www.blackwell-synergy.com/toc/add/84/11>
- 43 McPherson A, Benson G, Forrest EH. Appraisal of the Glasgow assessment and management of alcohol guideline: a comprehensive alcohol management protocol for use in general hospitals. *QJM* 2012;105:649–56.
- 44 Lackner C, Spindelboeck W, Haybaeck J, *et al.* Histological parameters and alcohol abstinence determine long-term prognosis in patients with alcoholic liver disease. *J Hepatol* 2017;66:S0168-8278(16)30685-7:610–8..
- 45 Hofer BS, Simbrunner B, Hartl L, *et al.* Alcohol abstinence improves prognosis across all stages of portal hypertension in alcohol-related cirrhosis. *Clin Gastroenterol Hepatol* 2023;21:S1542-3565(22)01113-2:2308–2317..
- 46 Trebicka J, Fernandez J, Papp M, *et al.* The PREDICT study Uncovers three clinical courses of acutely decompensated cirrhosis that have distinct pathophysiology. *J Hepatol* 2020;73:S0168-8278(20)30384-6:842–54..
- 47 Parker R, Im G, Jones F, *et al.* Clinical and Microbiological features of infection in alcoholic hepatitis: an international cohort study. *J Gastroenterol* 2017;52:1192–200.
- 48 Louvet A, Wartel F, Castel H, *et al.* Infection in patients with severe alcoholic hepatitis treated with steroids: early response to therapy is the key factor. *Gastroenterology* 2009;137:541–8.
- 49 Gustot T, Maillart E, Bocci M, *et al.* Invasive Aspergillosis in patients with severe alcoholic hepatitis. *J Hepatol* 2014;60:S0168-8278(13)00667-3:267–74..
- 50 Louvet A, Labreuche J, Dao T, *et al.* Effect of prophylactic antibiotics on mortality in severe alcohol-related hepatitis: A randomized clinical trial. *JAMA* 2023;329:1558–66.



- 51 Carey EJ, Lai JC, Sonnenday C, *et al.* A North American expert opinion statement on Sarcopenia in liver transplantation. *Hepatology* 2019;70:1816–29.
- 52 Fried LP, Tangen CM, Walston J, *et al.* Frailty in older adults: evidence for a phenotype. *J Gerontol A Biol Sci Med Sci* 2001;56:M146–56.
- 53 McClain CJ, Barve SS, Barve A, *et al.* Alcoholic liver disease and malnutrition. *Alcohol Clin Exp Res* 2011;35:815–20.
- 54 Dhaliwal A, Armstrong MJ. Sarcopenia in cirrhosis: A practical overview. *Clin Med (Lond)* 2020;20:489–92.
- 55 European Association for the Study of the Liver. Electronic address: Easloffice@Easloffice. EU, European Association for the study of the liver. EASL clinical practice guidelines on nutrition in chronic liver disease. *J Hepatol* 2018;70:172–93.
- 56 Moreno C, Deltenre P, Senterre C, *et al.* Intensive Enteral nutrition is ineffective for patients with severe alcoholic hepatitis treated with corticosteroids. *Gastroenterology* 2016;150:S0016-5085(15)01866-1:903–10..
- 57 da Silva Boteon APC, Chauhan A, Boteon YL, *et al.* Predictive factors for 28-day mortality in acute-on-chronic liver failure patients admitted to the intensive care unit. *Dig Liver Dis* 2019;51:S1590-8658(19)30547-X:1416–22..
- 58 Moreau R, Jalan R, Gines P, *et al.* Acute-on-chronic liver failure is a distinct syndrome that develops in patients with acute Decompensation of cirrhosis. *Gastroenterology* 2013;144:S0016-5085(13)00291-6:1426–37, .
- 59 Lone NI, Lee R, Walsh TS. Long-term mortality and hospital resource use in ICU patients with alcohol-related liver disease. *Crit Care Med* 2019;47:23–32.
- 60 Kalambokis G, Manousou P, Vibhakorn S, *et al.* Transjugular liver biopsy--indications, adequacy, quality of specimens, and complications--a systematic review. *J Hepatol* 2007;47:284–94.
- 61 Ventura-Cots M, Argemi J, Jones PD, *et al.* Clinical, histological and molecular profiling of different stages of alcohol-related liver disease. *Gut* 2022;71:1856–66.
- 62 Katoonizadeh A, Laleman W, Verslype C, *et al.* Early features of acute-on-chronic alcoholic liver failure: a prospective cohort study. *Gut* 2010;59:1561–9.
- 63 Crabb DW, Bataller R, Chalasani NP, *et al.* Standard definitions and common data elements for clinical trials in patients with alcoholic hepatitis: recommendation from the NIAAA alcoholic hepatitis consortia. *Gastroenterology* 2016;150:S0016-5085(16)00233-X:785–90..
- 64 Forrest E, Goldin R. The diagnostic and Prognostic significance of liver histology in alcoholic hepatitis. *Aliment Pharmacol Ther* 2021;54:864.
- 65 Crabb DW, Im GY, Szabo G, *et al.* Diagnosis and treatment of alcohol-associated liver diseases: 2019 practice guidance from the American Association for the study of liver diseases. *Hepatology* 2020;71:306–33.
- 66 Maddrey WC, Boitnott JK, Bedine MS, *et al.* Corticosteroid therapy of alcoholic hepatitis. *Gastroenterology* 1978;75:193–9.
- 67 Carithers RL Jr, Herlong HF, Diehl AM, *et al.* Methylprednisolone therapy in patients with severe alcoholic hepatitis. A randomized multicenter trial. *Ann Intern Med* 1989;110:685–90.
- 68 Robert A, Chazouillères O. Prothrombin time in liver failure: time, ratio, activity percentage, or international normalized ratio. *Hepatology* 1996;24:1392–4.
- 69 Forrest EH, Evans CDJ, Stewart S, *et al.* Analysis of factors predictive of mortality in alcoholic hepatitis and derivation and validation of the Glasgow alcoholic hepatitis score. *Gut* 2005;54:1174–9.
- 70 Forrest EH, Storey N, Sinha R, *et al.* Baseline neutrophil-to-lymphocyte ratio predicts response to corticosteroids and is associated with infection and renal dysfunction in alcoholic hepatitis. *Aliment Pharmacol Ther* 2019;50:442–53.
- 71 Dominguez M, Rincón D, Abalde JG, *et al.* A new scoring system for Prognostic stratification of patients with alcoholic hepatitis. *Am J Gastroenterol* 2008;103:2747–56.
- 72 Morales-Arráez D, Ventura-Cots M, Altamirano J, *et al.* The MELD score is superior to the Maddrey discriminant function score to predict short-term mortality in alcohol-associated hepatitis: A global study. *Am J Gastroenterol* 2022;117:301–10.
- 73 Forrest EH, Atkinson SR, Richardson P, *et al.* Application of Prognostic scores in the STOPAH trial: discriminant function is no longer the optimal scoring system in alcoholic hepatitis. *Journal of Hepatology* 2018;68:511–8.
- 74 Singal AK, Bataller R, Ahn J, *et al.* ACG clinical guideline: alcoholic liver disease. *Am J Gastroenterol* 2018;113:175–94.
- 75 Thursz MR, Richardson P, Allison M, *et al.* Prednisolone or Pentoxifylline for alcoholic hepatitis. *N Engl J Med* 2015;372:1619–28.
- 76 Louvet A, Thursz MR, Kim DJ, *et al.* Corticosteroids reduce risk of death within 28 days for patients with severe alcoholic hepatitis, compared with Pentoxifylline or placebo-a meta-analysis of individual data from controlled trials. *Gastroenterology* 2018;155:458–468.
- 77 Vergis N, Atkinson SR, Knapp S, *et al.* In patients with severe alcoholic hepatitis, prednisolone increases susceptibility to infection and infection-related mortality, and is associated with high circulating levels of bacterial DNA. *Gastroenterology* 2017;152:1068–1077.
- 78 Parker R, Cabezas J, Altamirano J, *et al.* Trajectory of serum bilirubin predicts spontaneous recovery in a real-world cohort of patients with alcoholic hepatitis. *Clin Gastroenterol Hepatol* 2022;20:S1542-3565(21)00092-6:e289–97..
- 79 Louvet A, Naveau S, Abdelnour M, *et al.* The Lille model: a new tool for therapeutic strategy in patients with severe alcoholic hepatitis treated with steroids. *Hepatology* 2007;45:1348–54.
- 80 Garcia-Saenz-de-Sicilia M, Duvoor C, Altamirano J, *et al.* A Day-4 Lille model predicts response to corticosteroids and mortality in severe alcoholic hepatitis. *Am J Gastroenterol* 2017;112:666:306–15..
- 81 GOV UK. Hospital discharge and community support guidance, Available: <https://www.gov.uk/government/publications/hospital-discharge-and-community-support-guidance/hospital-discharge-and-community-support-guidance>
- 82 Majc D, Tepes B. The impact of outpatient clinical care on the survival and Hospitalisation rate in patients with alcoholic liver cirrhosis. *Radiol Oncol* 2018;52:75–82.
- 83 Kanwal F, Asch SM, Kramer JR, *et al.* Early outpatient Follow-Up and 30-Day outcomes in patients hospitalized with cirrhosis. *Hepatology* 2016;64:569–81.
- 84 Atkinson SR, Way MJ, McQuillin A, *et al.* Homozygosity for Rs738409:G in Pnpl3 is associated with increased mortality following an episode of severe alcoholic hepatitis. *Journal of Hepatology* 2017;67:120–7.
- 85 Stagg HR, Surey J, Francis M, *et al.* Improving engagement with Healthcare in hepatitis C: a randomised controlled trial of a peer support intervention. *BMC Med* 2019;17:71.
- 86 Harris M, Bonnington O, Harrison G, *et al.* Understanding hepatitis C intervention success-qualitative findings from the Hepcatt study. *J Viral Hepat* 2018;25:762–70. 10.1111/jvh.12869 Available: <http://doi.wiley.com/10.1111/jvh.12869> Available: <http://doi.wiley.com/10.1111/jvh.12869> Available: <http://doi.wiley.com/10.1111/jvh.12869>
- 87 Rutledge SM, Asgharpour A. Smoking and liver disease. *Gastroenterol Hepatol* 2020;16:617–25.
- 88 Hagström H. Alcohol, smoking and the liver disease patient. *Best Pract Res Clin Gastroenterol* 2017;31:S1521-6918(17)30098-7:537–43..
- 89 Addolorato G, Mirijello A, Leggio L, *et al.* Liver transplantation in alcoholic patients: impact of an alcohol addiction unit within a liver transplant center. *Alcohol Clin Exp Res* 2013;37:1601–8.
- 90 Elfeki MA, Abdallah MA, Leggio L, *et al.* Simultaneous management of alcohol use disorder and liver disease: A systematic review and meta-analysis. *J Addict Med* 2023;17:e119–28.
- 91 Addolorato G, Leggio L, Ferrulli A, *et al.* Effectiveness and safety of baclofen for maintenance of alcohol abstinence in alcohol-dependent patients with liver cirrhosis: randomised, double-blind controlled study. *Lancet* 2007;370:1915–22.
- 92 Morley KC, Baillie A, Fraser I, *et al.* Baclofen in the treatment of alcohol dependence with or without liver disease: Multisite, randomised, double-blind, placebo-controlled trial. *Br J Psychiatry* 2018;212:362–9.
- 93 Burra P, Senzolo M, Adam R, *et al.* Liver transplantation for alcoholic liver disease in Europe: a study from the ELTR (European liver transplant Registry). *Am J Transplant* 2010;10:138–48.
- 94 Masson S, Aldersley H, Leithead JA, *et al.* Liver transplantation for alcohol-related liver disease in the UK: revised UK liver advisory group recommendations for referral. *Lancet Gastroenterol Hepatol* 2021;6:S2468-1253(21)00195-3:947–55..
- 95 Millson C, Considine A, Cramp ME, *et al.* Adult liver transplantation: A UK clinical guideline - part 1: pre-operation. *Frontline Gastroenterol* 2020;11:375–84.
- 96 NHS blood and transplant (liver advisory group). Liver transplantation: selection criteria and recipient registration. *Policy POL* 2018. Available: <https://nhs.uk/blood-and-transplant/policies/liver-selection-policy.pdf>

- 97 Barber K, Madden S, Allen J, *et al.* Elective liver transplant list mortality: development of a United Kingdom end-stage liver disease score. *Transplantation* 2011;92:469–76.
- 98 Woodland H, Hudson B, Forbes K, *et al.* British Association for the study of the liver (BASL) end of life special interest group. palliative care in liver disease: what does good look like. *Frontline Gastroenterol* 2020;11:218–27.
- 99 Hudson B, Round J, Georgeson B, *et al.* Cirrhosis with Ascites in the last year of life: a nationwide analysis of factors shaping costs, health-care use, and place of death in England. *The Lancet Gastroenterology & Hepatology* 2018;3:95–103.
- 100 Macken L, Corrigan M, Prentice W, *et al.* Palliative long-term abdominal drains for the management of refractory Ascites due to cirrhosis: a consensus document. *Frontline Gastroenterol* 2022;13:e116–25.
- 101 Shah ND, Ventura-Cots M, Abraldes JG, *et al.* Alcohol-related liver disease is rarely detected at early stages compared with liver diseases of other Etiologies worldwide. *Clinical Gastroenterology and Hepatology* 2019;17:2320–2329.
- 102 Chalmers J, Wilkes E, Harris R, *et al.* Development and implementation of a commissioned pathway for the identification and stratification of liver disease in the community. *Frontline Gastroenterol* 2020;11:86–92.