

2023-10

Genome-wide Analysis of Motor Progression in Parkinson Disease

Martinez Carrasco, A

<https://pearl.plymouth.ac.uk/handle/10026.1/21618>

10.1212/nxg.0000000000200092

Neurology Genetics

Ovid Technologies (Wolters Kluwer Health)

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

Genome-wide Analysis of Motor Progression in Parkinson Disease

Alejandro Martínez Carrasco, MSc, Raquel Real, PhD, Michael Lawton, PhD, Regina Hertfelder Reynolds, PhD, Manuela Tan, PhD, Lesley Wu, MSc, Nigel Williams, PhD, Camille Carroll, MD, Jean-Christophe Corvol, MD, PhD, Michele Hu, PhD, Donald Grosset, MD, John Hardy, PhD, Mina Ryten, PhD, Yoav Ben-Shlomo, PhD, Maryam Shoai, PhD, and Huw R. Morris, PhD

Correspondence

Mr. Martínez Carrasco
alejandro.carrasco.20@ucl.ac.uk

Neurol Genet 2023;9:e200092. doi:10.1212/NXG.0000000000200092

Abstract

Background and Objectives

The genetic basis of Parkinson disease (PD) motor progression is largely unknown. Previous studies of the genetics of PD progression have included small cohorts and shown a limited overlap with genetic PD risk factors from case-control studies. Here, we have studied genomic variation associated with PD motor severity and early-stage progression in large longitudinal cohorts to help to define the biology of PD progression and potential new drug targets.

Methods

We performed a GWAS meta-analysis of early PD motor severity and progression up to 3 years from study entry. We used linear mixed-effect models with additive effects, corrected for age at diagnosis, sex, and the first 5 genetic principal components to assess variability in axial, limb, and total Movement Disorder Society–Unified Parkinson's Disease Rating Scale (MDS-UPDRS) III scores.

Results

We included 3,572 unrelated European ancestry patients with PD from 5 observational cohorts and 1 drug trial. The average AAO was 62.6 years (SD = 9.83), and 63% of participants were male. We found an average increase in the total MDS-UPDRS III score of 2.3 points/year. We identified an association between PD axial motor progression and variation at the *GJA5* locus at 1q12 ($\beta = -0.25$, SE = 0.04, $p = 3.4e^{-10}$). Exploration of the regulation of gene expression in the region (*cis*-expression quantitative trait loci [eQTL] analysis) showed that the lead variant was associated with expression of *ACP6*, a lysophosphatidic acid phosphatase that regulates mitochondrial lipid biosynthesis (*cis*-eQTL p -values in blood and brain RNA expression data sets: $<10^{-14}$ in eQTLGen and 10^{-7} in PsychEncode).

Discussion

Our study highlights the potential role of mitochondrial lipid homeostasis in the progression of PD, which may be important in establishing new drug targets that might modify disease progression.

From the Department of Clinical and Movement Neurosciences (A.M.C., R.R., L.W., H.R.M.), UCL Queen Square Institute of Neurology; UCL Movement Disorders Centre (A.M.C., R.R., L.W., H.R.M.), University College London, United Kingdom; Aligning Science Across Parkinson's (ASAP) Collaborative Research Network (A.M.C., R.R., R.H.R., L.W., M.R., M.S., J.H., H.R.M.), Chevy Chase, MD; Population Health Sciences (M.L., Y.B.-S.), Bristol Medical School, University of Bristol; Genetics and Genomic Medicine (R.H.R., M.R.), UCL Great Ormond Street Institute of Child Health, University College London, United Kingdom; Department of Neurology (M.T.), Oslo University Hospital, Norway; Institute of Psychological Medicine and Clinical Neurosciences (N.W.), MRC Centre for Neuropsychiatric Genetics and Genomics, Cardiff University; Faculty of Health (C.C.), University of Plymouth, United Kingdom; Sorbonne Université (J.-C.C.), Institut du Cerveau - Paris Brain Institute - ICM, INSERM, CNRS; Assistance Publique Hôpitaux de Paris (J.-C.C.), Department of Neurology, Hôpital Pitié-Salpêtrière, France; Division of Clinical Neurology (M.H.), Nuffield Department of Clinical Neurosciences; Oxford Parkinson's Disease Centre (M.H.), University of Oxford; School of Neuroscience and Psychology (D.G.), University of Glasgow; Department of Neurodegenerative Diseases (J.H., M.S.), UCL Queen Square Institute of Neurology; UK Dementia Research Institute (J.H., M.S.), University College London; Reta Lila Weston Institute (J.H., M.S.), UCL Queen Square Institute of Neurology; National Institute for Health Research (NIHR), University College London Hospitals Biomedical Research Centre (J.H.); Institute for Advanced Study (J.H.), The Hong Kong University of Science and Technology, Hong Kong SAR, China; and NIHR Great Ormond Street Hospital Biomedical Research Centre (M.R.), University College London, United Kingdom.

Go to [Neurology.org/NG](https://www.neurology.org/NG) for full disclosures. Funding information is provided at the end of the article.

The Article Processing Charge was funded by MJFF.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Glossary

eQTL = expression quantitative trait loci; **GD** = Gaucher disease; **GWA** = genome-wide association; **LD** = linkage disequilibrium; **LMMs** = linear mixed-effect models; **MDS-UPDRS** = Movement Disorder Society–Unified Parkinson’s Disease Rating Scale; **PD** = Parkinson disease; **QC** = quality control; **TSS** = transcription start site; **WGS** = whole genome sequencing.

Introduction

Parkinson disease (PD) is a progressive neurodegenerative disorder with motor and nonmotor symptoms, clinically manifesting with rigidity, postural instability, and slowness of movement (bradykinesia). The motor deficits are linked to the loss of dopaminergic (DA) neurons in the substantia nigra pars compacta, their projection to the striatum, and the accumulation of alpha-synuclein aggregates in Lewy bodies in the nigral and other neurons.¹

PD is heterogeneous in its progression and onset. The predominant motor phenotype is influenced by age at onset, with tremor being more prominent in older patients, and the risk of dystonia at presentation increasing in younger individuals.² With respect to progression, younger patients tend to have a slower rate of motor progression as measured using the Movement Disorder Society–Unified Parkinson’s Disease Rating Scale (MDS-UPDRS) and Hoehn and Yahr assessments.³ Functional imaging studies have also shown a slower rate of decline in the loss of nigrostriatal terminals in early onset compared with late-onset PD.⁴

To date, most PD genetic studies have focused on the risk of developing PD.⁵ Relatively little is known about the genetic factors that contribute to variation in the onset and progression of motor and nonmotor symptoms. Studies of PD age at onset have shown that the genetic determinants of age at onset are different to the genetic factors determining case-control status, with the *MAPT* and *GCH-1* loci associated with disease risk but not age at onset.^{6–8} With respect to common variability explaining differences in disease progression, Iwaki et al.⁹ performed a GWAS looking at 25 different outcome measures in a meta-analysis of 12 longitudinal cohorts, including mortality, dementia, disease severity, and patient disability. They reported an association between an intronic variant in *SLC44A1*, a mitochondrial choline transporter, and motor progression, reflected by reaching a Hoehn and Yahr score higher or equal than 3 (HY3). A more recent study¹⁰ took a different approach using a principal components (PC)–based measure that combined multiple assessments for composite motor and cognitive progression. They found a novel association between *ATP8B2* and PC-based motor progression in a gene-based analysis.

These studies suggest that pathways specifically related to the progression of PD can be understood from genotype/phenotype analysis, which may ultimately lead to the development of new disease-modifying therapies. We modeled the early stages of

motor Parkinson disease, using the total score from the MDS-UPDRS part III, a validated scale recommended for clinical trials to measure both response to levodopa treatment and the rate of change over time.¹¹ We have also derived and analyzed separate axial and limb motor stages from the scale, as they may relate in part to distinct pathology.¹² We have used a genome-wide association (GWA) approach to define genetic determinants associated with variation in motor progression and severity, identifying genetic variation that is significantly associated with change in the MDS-UPDRS, the primary outcome measure for many clinical PD trials. Finally, we have performed functional annotation and fine-mapping to understand how the nominated genetic variants are associated with the regulation of gene expression and the underlying biology of PD motor traits.

Methods

The workflow for all methods applied in the manuscript is available on a Zenodo repository (doi.org/10.5281/zenodo.7258985).

Study Design and Quality Control at the Sample and Genetic Level

We studied 6 observational and interventional longitudinal PD cohorts with either genotyping or whole genome sequencing (WGS) data available, totaling 4,971 patients (eTable 1, [links. lww.com/NXG/A624](https://www.nxg.org/A624)). We selected cohorts which included longitudinal MDS-UPDRS part III assessments from the MDS-UPDRS¹¹ and applied quality control (QC) at the clinical data level (eFigure 1). To study the motor progression of PD, we derived limb and axial phenotypes from the MDS-UPDRS part III scale based on previously accepted definitions.¹³ In addition, we used the MDS-UPDRS III total score as an overall measure of PD motor state (eTable 2). We included data up to 36 months from the baseline visit, within longitudinal observational and therapeutic cohorts, as we had high rates of data completion up to 36 months. Imputation of patients’ missing motor outcomes was performed when possible¹⁴ (eMethods; eTable 3).

We applied genetic QC at the sample and variant level followed by imputation in the Michigan Imputation Server (RRID:SCR_017579)¹⁵ and postimputation QC (eMethods; eFigure 1, [links. lww.com/NXG/A624](https://www.nxg.org/A624)). Table 1 summarizes the demographics of the data after QC.

Previous studies have reported that levodopa improves motor state examination and may possibly slow disease progression.¹⁶ Because the motor improvement is noticeable a few hours after treatment, and affects the MDS-UPDRS measure, we have

Table 1 Cohort Demographics and Motor Scores Rate of Change

Study	N patients ^a	N Observations ^b	Visit interval, mo	ON/OFF	No. (%) male	AAD, y Mean ± SD	AAB, y Mean ± SD	Duration ^c , y Mean ± SD	Total rate of change per year Mean ± SD	Total rate of change centered Mean ± SD	Limb rate of change centered Mean ± SD	Axial rate of change centered Mean ± SD
TPD	1,699	4,349	18	ON	1101 (64.8)	66.20 ± 9.24	67.50 ± 9.31	3.12 ± 3.04	2.7 ± 4.69	0.48 ± 0.92	0.54 ± 1.25	0.71 ± 1.37
OPDC	797	1,978	18	ON	513 (64.37)	66.04 ± 9.46	67.25 ± 9.57	2.86 ± 1.84	2.85 ± 4.27	0.50 ± 0.77	0.50 ± 0.93	1.02 ± 1.76
PPMI	287	1,653	3 & 6 & 12	OFF	184 (64.11)	61.01 ± 9.73	61.59 ± 9.70	2.02 ± 1.97	3.01 ± 3.65	0.84 ± 0.94	1.03 ± 1.23	0.67 ± 1.05
PD STAT	124	358	12	OFF	76 (61.29)	NA	66.08 ± 9.37	10.14 ± 4.06	1.70 ± 5.76	0.06 ± 0.41	0.07 ± 0.58	0.27 ± 0.61
DIGPD	305	1,005	12	ON	184 (60.33)	59.68 ± 9.85	62.59 ± 9.70	3.74 ± 1.55	2.37 ± 3.41	0.56 ± 0.87	0.67 ± 1.40	0.52 ± 0.95
PDBP	360	2,090	6	ON	222 (61.67)	59.9 ± 10.90	64.73 ± 9.15	NA	1.05 ± 4.11	0.21 ± 0.61	0.30 ± 0.90	0.40 ± 1.46

Abbreviations: AAB = Age at Baseline; AAD = Age at Diagnosis; N = Number.

Total rate of change per year: (Last visit score - Baseline score) ÷ number of years

Total/Limb/Axial rate of change centered: (Last visit score - Baseline score) ÷ Baseline Score.

^a N patients = number of patients with PD after QC was applied.

^b N observations = number of observations including MDS-UPDRS part III motor assessment after QC was applied.

^c Duration = disease duration from onset to study entry.

compared individuals from cohorts in the same state at each assessment. If cohorts had data in the “OFF” state available, we used longitudinal “OFF” vs “OFF” MDS-UPDRS part III scores, otherwise “ON” vs “ON” comparisons were made. In addition, we performed sensitivity analyses adjusting the motor scores by levodopa dosage (eMethods; eFigure 2).

Statistical Approaches

To study the effect of genetics on motor progression as well as on baseline variability, we used linear mixed-effect models (LMMs) which can be used to study longitudinal measures as the variability in the outcome can be studied at 2 levels, within groups (i.e., individual patient longitudinal visits) or between groups (across patients). We studied the changes in disease limb and axial motor severity and progression associated with genetic variants under an additive genetic effect (eMethods, links.lww.com/NXG/A624). In the disease severity model, we studied the additive genetic effect of SNPs on the average motor outcome. In the disease progression model, we studied the genetic additive effect of SNPs on the outcomes’ rate of change (eMethods). Both progression and severity models met the LMMs model assumptions (eFigure 3). We quantified the power of LMMs to investigate the effect of SNPs in changes in the motor signs of PD (eMethods). To run the disease progression model, we used lmerTest R package (v. 3.1-3; RRID:SCR_015656) and the Satterthwaite approach to approximate degrees of freedom and derive *p*-values and fitted them using restricted maximum likelihood (REML), as it produces acceptable type-1 error rates.¹⁷ In addition, we used SCEBE¹⁸ algorithm (v. 0.1.0) with REML and lme4 R package (v. 1.1-30; RRID:SCR_015654) to decrease the computational expense of adding unexplained variability at the slope level in the disease severity model (eMethods). We validated SCEBE in 2 separate cohorts (eFigure 4). All tests were two-tailed. We used METAL software (version released on the 25/03/2011; RRID:SCR_002013) for meta-analysis of genome-wide association summary statistics, using a fixed effects model weighted by β coefficients and the inverse of the standard errors.^{19,20} We also applied QC to the meta-analysis results (eMethods). Statistical significance was assessed at the genome-wide level ($p = 5 \times 10^{-8}$).

Fine-Mapping and Functional Annotation

We performed a conditional and a stepwise model selection procedure to determine whether there were independently associated SNPs in each locus of interest. To nominate causal variants, we applied fine-mapping. To further understand regulatory mechanisms in nominated loci, we mapped each locus against (1) cell type-specific and “bulk” genome enhancer marks, (2) enhancer-transcription start site (TSS) interaction marks from FANTOM5, and (3) brain cell type-specific transcriptional regulatory marks and distal enhancer-promoter interactions.^{21,22} To determine whether causal variants could be linked to the motor phenotypes by dysregulation of gene expression, we performed colocalization against cis-expression quantitative trait loci (eQTL) data sets.²³⁻²⁵ Software and packages to conduct the analyses and access the data are described in eMethods. In addition, we used FUMA, a web-based platform that integrates a wide range of

functional annotation data (RRID:SCR_017521; version 1.3.8).²⁶ We used LocusZoom (RRID:SCR_009257; version 0.12)²⁷ to display the linkage disequilibrium (LD) structure of a given locus against the locus lead SNP, as well as the protein coding genes and rRNAs nearby.

Standard Protocol Approvals, Registrations, and Patient Consents

Each participant provided written informed consent for participation. TPD has multicenter research ethics approval from the West of Scotland Research Ethics Committee: IRAS 70980, MREC 11/AL/0163 (ClinicalTrials.gov, NCT02881099). OPDC has multicenter research ethics approval from the South Central Oxford Research Ethics Committee 16/SC/0108. PD STAT has been approved by the North East-Newcastle and North Tyneside 2 Research Ethics Committee (ClinicalTrials.gov NCT02787590). The DIGPD study was sponsored by Assistance Publique Hôpitaux de Paris, approved by French regulatory authorities and an ethics committee and conducted according to good clinical practices (ClinicalTrials.gov NCT01564992).

Data Availability

We have made our summary statistics available (doi.org/10.5281/zenodo.7257484). TPD data are available on access request from trackingparkinsons.org.uk/about-1/data/. The PDBP and PPMI data were accessed from Accelerating Medicines Partnership: Parkinson's Disease (AMP-PD), and data are available on registration at amp-pd.org/. OPDC data are available on request from the Dementias Platform UK (portal.dementiasplatform.uk/Apply). DIGPD data are available on request to the principal investigator (JC Corvol, Assistance Publique Hôpitaux de Paris; clinicaltrials.gov/ct2/show/NCT01564992). PD STAT is available on request to the principal investigator (C Carroll, Plymouth University, pencu.psmd.plymouth.ac.uk/pdstat/#:~:text=PD%20STAT%20%2D%20Simvastatin%20as%20a,brain%20from%20injury%20or%20loss). HapMap phase 3 data (HapMap3) is available for download at <ftp://ftp.ncbi.nlm.nih.gov/hapmap/>. Cis-QTL data were obtained from eQTLGen (eqtlgen.org/cis-eqtls.html) and PsychENCODE (resource.psychencode.org). MetaBrain cis-eQTL data can be accessed on access request form (metabrain.nl/cis-eqtls.html). eQTL data from eQTL catalog can be ftp-accessed (ebi.ac.uk/eqtl/Data_access/). FANTOMS CAGE-seq and Nott brain cell type-specific enhancer-promoter interactome data were accessed through echolocator (github.com/RajLabMSSM/echolocator).

Code Availability

Code used in the analysis is available from github.com/AMCalejandro/EMPD (doi.org/10.5281/zenodo.7258985). Analysis was performed using open-source tools as described in the Methods section.

Results

We explored the overall rate of change in MDS-UPDRS part III total, limb, and axial scores (Table 1). There was variation


across studies. We specifically studied the amount of change for the motor measures in each study by comparing the final score with the baseline score, divided by the baseline score, for MDS-UPDRS-total, axial, and limb. We found that the axial score rate of change was the highest in TPD, OPDC, PD STAT, and PDBP. The limb rate of change was the highest in PPMI and DIGPD. PD STAT and PDBP had a lower rate of changes, which may be due either to longer disease duration or to selection effects related to the inclusion of “benign” PD in patients with longer disease duration. We assessed this by fitting an LMM using data from TPD and found a significant interaction between time and disease duration related to MDS-UPDRS total progression ($\beta = -0.11$, SE = 0.04, $p = 0.01$). Longer disease duration was associated with a lower total rate of change in MDS-UPDRS, which appears to be nonlinear with extended disease durations. Overall, we confirmed that the MDS-UPDRS-derived measures increased, reflecting worsening motor impairment, from study entry up to 3 years (Figure 1). The MDS-UPDRS part III total yearly rate of change ranged between 2.37 and 3.01 points/year, which is consistent with previous reports.²⁸

Our power calculation showed that the current LMM was well powered to detect high effect sizes ($\beta \geq 0.2$) for a wide range of different MAFs, with a limit for variants with an allele frequency $\leq 1\%$ (eFigure 5, links.lww.com/NXG/A624). We performed a GWAS on each cohort to study PD motor progression and meta-analyzed results separately using a genomic control to correct the test statistics of those cohorts that had genomic inflation ($\lambda > 1$ & $\lambda < 1.2$) (eTable 4).

We evaluated disease progression and disease severity models for total, limb, and axial progression. We did not find any significant genetic association with the PD limb motor progression or severity. For axial motor progression, we found 1 haplotype block that reached genome-wide significance (*GJA5* in chromosome 1) (Figure 2A; eFigure 6, links.lww.com/NXG/A624). This association was also found, at a lower significance level, with the MDS-UPDRS part III total. Given that there was no association with PD limb motor progression and severity, this relates to the inclusion of axial components in the overall MDS-UPDRS-III total score. Although the lead variant in the *GJA5* locus was not captured in the PPMI WGS data, we found proxy variants that were present in all cohorts. The lead proxy variant was rs12037169 ($\beta = -0.25$, SE = 0.04, $p = 3.93e^{-10}$) (Table 2). The association test statistic and directionality of each of these variants was consistently replicated across cohorts (eTable 5; Figure 2B).

To test whether levodopa was a major confounder of our study of motor progression, we corrected the patient motor scores using an equation described in eMethods (links.lww.com/NXG/A624) that best predicted the effect of levodopa dose on MDS-UPDRS part III total over time to correct the motor scores by levodopa usage. To weight the effect of levodopa usage on the limb and axial motor states, we used data from Tracking Parkinson's Levodopa challenge²⁹ with MDS-UPDRS part III scores recorded before and after treatment and used these

Figure 1 MDS-UPDRS III Motor Scores Trajectories



Trajectory of the MDS-UPDRS III-derived motor scores across cohorts. In the x-axis, the time point at which the MDS-UPDRS III assessment was measured. Each plot shows the motor scores trajectories on each cohort highlighted in the label. The y-axis represents the average scores for each of the motor states. The bars represent the SD of the average motor scores.

weights to correct the motor scores by levodopa usage. We did not find significant changes either in the SNPs significance level or the direction of effects. In addition, rs120371169 remained significantly associated with axial motor progression (eTable 5).


Hoehn and Yahr (HY), a measure of a patient's disability, measures motor progression including loss of balance and provides an alternative measure to the axial score from MDS-UPDRS part III. To further validate the GWAS association linked to PD axial motor progression, we ran the disease progression statistical model to study the contribution of SNPs to motor changes over time using HY as our longitudinal outcome. We found an LD block approaching genome-wide significance within *GJAS* locus, the same locus found to be significantly associated with axial motor progression (eFigure 7, links.lww.com/NXG/A624). The lead variant was rs36005900 ($\beta = -0.08$, $SE = 0.0078$, $p = 5.7e^{-7}$). We found the directionality of the effects to be the same as in the axial motor progression GWAS. In addition, rs36005900 is in LD with the lead variant reported in the same locus for MDS-UPDRS III axial motor progression ($D' = 0.8$, $R^2 = 0.6$).

We then investigated whether there were independently associated SNPs at the *GJAS* locus. We did not find any signal other than the lead SNP in the selection procedure under a conditional and stepwise selection approach using GCTA-

COJO. Under a single causal variant assumption, we then performed statistical fine-mapping. We did not resolve consensus SNPs (a SNP nominated to be causal by 2 different fine-mapping tools) at the *GJAS* locus. We found a total of 12 SNPs with support for causality of changes in motor axial progression, nominated from at least 1 fine-mapping tool (eTable 6, links.lww.com/NXG/A624). We did not find an overlap between the *GJAS* locus haplotype block and regulatory marks from functional annotation data sets described in eMethods.

We also explored eQTLs data sets through the FUMA platform. We found that many of the GWAS-significant SNPs within the *GJAS* locus were significant cis-eQTLs for *ACP6*, in PsychEncode, and eQTLGen. In particular, we found that the lead variant was a significant eQTL in PsychEncode, and eQTLGen, and also rs12037169, the proxy significant variant found in all cohorts, was a significant cis-eQTL in eQTLGen (eTable 7, links.lww.com/NXG/A624). We then performed a colocalization analysis to evaluate whether there was colocalization between the GWAS axial progression results and eQTL GWAS for gene expression at the *GJAS* locus (eMethods). We used cis-eQTL data from eQTLGen and Metabrain cortex tissue cis-eQTLs data sets and performed a colocalization test for any gene within ± 1 Mb from the *GJAS* lead SNP. We did not find direct colocalization evidence for any gene, including *ACP6*. We found PPH3 (indicating separate significant

Figure 2 GWAS Meta-analysis of Motor Axial Progression



(A) Manhattan plot of the rate of axial change GWAS meta-analysis. On the x-axis, each of the 22 chromosomes and each SNP p -value on the Y axis. The red dashed line indicates the genome-wide significance threshold p -value = $5e^{-8}$. The LD block that reached genome-wide significance on chromosome 1 is on the *GJA5* locus. Each dot corresponds to the p -value of the conditional likelihood interaction term between SNP and time (SNP*time). (B) Forest plots for proxy variant rs12037169 within *GJA5* locus under the GWAS meta-analysis using disease progression model for the axial outcome ($I^2 = 40.1$; Cochran Q test: $\chi^2 = 9.64$, $df = 5$, $p = 0.10$), annotated by study, effect size, and the corresponding 95% confidence interval.

associations for GWAS and eQTL analysis) to be the highest for the *ACP6* gene using default SNP priors (eQTLGen = 0.98, MetaBrain = 0.88). The PPH3 remained the highest for these 2 genes (PPH3 > 0.8), after we adjusted the priors according to the number of overlapping SNPs (eMethods).

In a separate analysis, we also studied the genome-wide effect that SNPs had on average changes of limb and axial motor states using the disease severity model (see eMethods, links.

lww.com/NXG/A624). We did not find any haplotype block that reached genome-wide significance. However, there were 2 distinct signals approaching genome-wide significance associated with changes in the average axial motor scores (*MAD1L1* in chromosome 7 and *LINC00511* in chromosome 17) (eFigure 8A). The lead SNP in *MAD1L1* was rs4721411 (Beta = 0.54, SE = 0.11, $p = 1.6e^{-7}$), and the lead variant in the long noncoding RNA *LINC00511* was rs36082764 (Beta = -0.62, SE = 0.11, $p = 6.3e^{-8}$) (Table 2). We found the association test

Table 2 Lead SNPs on the Disease Progression and Severity GWASs

rsID	Chr	Pos	A1	A2	MAF	Beta	SE	p Value	Nearest gene	Distance (Kb)	Type of variant	Model
rs6593808	1	147219250	A	G	0.23	-0.28	0.04	$1.35e^{-10}$	<i>GJA5</i>	0	Intergenic	Disease progression
rs12037169	1	147248057	A	G	0.25	-0.25	0.04	$3.93e^{-10}$	<i>GJA5</i>	0	Intergenic	Disease progression
rs4073509	2	192611013	C	T	0.02	0.52	0.10	$2.12e^{-7}$	<i>AC098872.3</i>	47,137	Intergenic	Disease progression
rs117239007	13	30550016	C	T	0.01	0.68	0.14	$4.71e^{-7}$	<i>LINC00544</i>	25,390	Intergenic	Disease progression
rs36082764	17	70330179	T	C	0.42	-0.62	0.11	$6.34e^{-8}$	<i>LINC00511</i>	0	ncRNA_intronic	Disease severity
rs4721411	7	2153071	T	C	0.40	0.53	0.10	$1.66e^{-7}$	<i>MAD1L1</i>	0	Intronic	Disease severity
rs10939702	4	10096692	T	G	0.45	0.57	0.12	$8.10e^{-7}$	<i>WDR1</i>	0	Intronic	Disease severity

statistic and directionality of each of these variants to be consistently replicated across cohorts (eTable 5, eFigure 8, B and C). Subsequent fine-mapping on both loci resolved rs3778978 in *MAD1L1* locus as the causal SNP and a list of 3 SNPs at the *LINC00511* locus (rs7213651, rs7218929, rs12950478) as the potential trait causing SNPs, narrowing down the spectrum of variants to be targeted in further in vivo and in vitro analyses (eTable 6). We found that the *MAD1L1* fine-mapped causal variant and the lead SNP overlapped with an active enhancer mark, and this region was predicted to interact with a TSS, supporting an effect of the GWAS-nominated variants in regulating the expression of *MAD1L1* (eFigure 9). For *LINC00511*, we found an anchored chromatin loop from the GWAS LD block in *LINC00511* to a region where the neuronal *SOX9* active promoter is found, suggesting that mutations in this distal regulatory region may alter *SOX9* expression in neurons specifically (eFigure 10).

We explored eQTL databases from FUMA. We found the lead variant in *MAD1L1* as well as the fine-mapped nominated causal variant to be a significant cis-eQTL in BIOS and eQTLGen (eTable 8, links.lww.com/NXG/A624). We then performed a colocalization analysis to evaluate whether there is a shared causal variant between the 2 traits (eMethods). We did not find direct colocalization evidence for any gene within ± 1 Mb from the GWAS lead SNPs. There was no cis-eQTL data available for *SOX9*. In the *MAD1L1* locus, the posterior probability H3 (PPH3) (association with both phenotypic and expression traits, but distinct causal variants) was the highest (PPH3 in *MAD1L1*: eQTLGen = 0.97, MetaBrain = 0.98, PsychENCODE = 0.75).

To understand how *ACP6*, *MAD1L1*, and *SOX9* might contribute to Parkinson disease motor function, we have described their biological function based on previous research (eResults, links.lww.com/NXG/A624).

Discussion

To understand the biology of motor progression in PD, we performed a large well-powered GWAS of PD motor progression. We have found 1 haplotype block at the *GJAS* locus that is significantly associated with axial PD motor progression. This association was consistently replicated across individual cohorts included in our motor progression GWAS meta-analysis and was replicated in an analysis of H/Y supporting our findings. Further exploration of the GWAS significant signals in eQTL databases suggests that the GWAS hits may control the expression of *ACP6*, an enzyme that regulates lipid metabolism in mitochondria.³⁰ Changes in *ACP6* concentrations are found in Gaucher disease (GD), although there is no clear link between *ACP6* levels in and GD progression. *ACP6* has a high astrocyte specificity,³¹ and mitochondrial dysfunction has been widely associated with PD etiology.³²

We used the MDS-UPDRS III (PD motor examination) scale, a sensitive measure of motor progression over time which has

been widely studied in observational and interventional studies of PD. A study of patients with untreated de novo PD in the PPMI study, followed up for 5 years to assess the progression of MDS-UPDRS, showed a linear increase of 2.4 points per year in MDS-UPDRS part III total score.²⁸ In this study, we observed a similar yearly rate of change for the total MDS-UPDRS score across the studies we included in our analysis (2.3 points/year on average) (Table 1). We have used linear mixed-effect models to investigate the common genetic variability associated with the severity and progression of distinct PD motor aspects. This concept may be consistent with PD subtype studies having a differential motor severity and progression.³³⁻³⁸ Another aspect of this differential approach to PD symptomatology is that limb and axial PD motor components may have a different cellular and pathophysiologic basis, with axial and limb motor symptoms related to cholinergic and dopaminergic dysfunction, respectively.^{39,40}

We corrected all models by AAO and sex and PCs as confounding variables. We performed a fixed effects meta-analysis as opposed to a pooled analysis to further account for between cohorts heterogeneity, as cohorts we included had different inclusion and exclusion criteria and were either genotyped with different microarrays or whole genome sequenced. Our results are not confounded by levodopa response, as defined in our sensitivity analysis (eFigure 1, eTable 1, links.lww.com/NXG/A624). In this data set, we have identified common genetic variability which determines axial, but not limb motor progression.

The lack of association between common genomic variation and the MDS-UPDRS limb subscale could be due to a combination of limited power and the levodopa effect in early disease. A study reported that measures of mobility, tremor, gait, and posture were consistent and reliable measures of PD progression.⁴¹ Because these measures are well represented in the axial score (except for tremor), this may be better powered to assess progression. Moreover, the limb signs may be more sensitive to levodopa use than the axial signs, making it possible that true genetic associations with limb motor progression were masked. Finally, we found the individual cohorts with the largest sample size had a higher axial rate of change compared with the limb rate of change (Table 1). A separate GWAS meta-analysis assessing the PD genetic contribution to the disease motor severity and subsequent functional annotation identified *MAD1L1* and *SOX9* as candidate genes associated with PD axial motor severity. Nevertheless, these potential associations did not reach genome-wide significance, and further analysis in distinct PD cohorts is needed for validation.

Strengths of our study include the large sample size and replication of our results across cohorts and across different measures of axial motor progression. Potential limitations of our identification of *ACP6* as the relevant gene at the *GJAS* locus include the lack of colocalization between the phenotype and expression GWAS, although these analyses are currently limited by the sample size of eQTL data sets and the lack of cell-specific gene expression data.

We hypothesize that expression of *ACP6* is important in the function in cell groups relevant to axial progression in PD including the pedunculopontine nucleus and that therapies directed toward mitochondrial lipid metabolism may be relevant to the disease modification. Further replication in independent cohorts genotyped in the global Parkinson's genetics program (GP2.org) will help to determine the importance of this region, and further analysis of this biochemical pathway may provide new insights into the pathogenesis of PD progression.

Acknowledgment

For the purpose of open access, the author has applied a CC BY public copyright licence to all Author Accepted Manuscripts arising from this submission. This research was supported by the National Institute for Health Research University College London Hospitals Biomedical Research Centre. The UCL Movement Disorders Centre is supported by the Edmond J. Safra Philanthropic Foundation. Data used in the preparation of this article were obtained from the AMP-PD Knowledge Platform (amp-pd.org). AMP-PD is a public-private partnership managed by the FNIH and funded by Celgene, GSK, Michael J. Fox Foundation for Parkinson's Research, the National Institute of Neurological Disorders and Stroke, Pfizer, and Verily. Clinical data and biosamples used in preparation of this article were obtained from the Parkinson's Progression Markers Initiative (PPMI) and the Parkinson's Disease Biomarkers Program (PDBP). PPMI—a public-private partnership—is funded by the Michael J. Fox Foundation for Parkinson's Research and funding partners, including [list the full names of all of the PPMI funding partners found at ppmi-info.org/about-ppmi/who-we-are/study-sponsors]. The PPMI Investigators have not participated in reviewing the data analysis or content of the manuscript. For up-to-date information on the study, visit ppmi-info.org. The Parkinson's Disease Biomarker Program (PDBP) consortium is supported by the National Institute of Neurological Disorders and Stroke (NINDS) at the National Institutes of Health. A full list of PDBP investigators can be found at pdbp.ninds.nih.gov/policy. The PDBP Investigators have not participated in reviewing the data analysis or content of the manuscript. The DIGPD cohort (ClinicalTrials.gov, NCT01564992) is a multicenter longitudinal cohort conducted in four Universities and four General Hospitals in France between 2009 and 2019, sponsored by Assistance Publique Hôpitaux de Paris, and funded by a grant from the French Ministry of Health (PHRC 2008, AOR0810) and a grant from the Agence Nationale de Sécurité et des Médicaments (ANSM-2013). The authors thank the DIGPD Study group which collected the data made available for this work. Both TPD and OPDC cohorts are primarily funded and supported by Parkinson's UK (parkinsons.org.uk/) and supported by the National Institute for Health and Care Research (NIHR) Clinical Research Network (CRN). The TPD study is also supported by NHS Greater Glasgow and Clyde. The OPDC cohort is also supported by the NIHR Oxford Biomedical Research Centre, based at the Oxford University Hospitals NHS Trust, and the University of Oxford. PD-STAT is funded and supported by grants from the Cure

Parkinson's Trust (cureparkinsons.org.uk/) and JP Moulton Charitable Foundation (perscitullp.com/moulton-charity-trust/), co-ordinated by the Peninsula Clinical Trials Unit, University of Plymouth, and sponsored by University Hospitals Plymouth NHS Trust. The Genotype-Tissue Expression (GTEx) Project was supported by the Common Fund of the Office of the Director of the National Institutes of Health and by NCI, NHGRI, NHLBI, NIDA, NIMH, and NINDS. The data used for the analyses described in this manuscript were obtained from console.cloud.google.com/storage/browser/gtex-resources on 01/26/2022. The PsychENCODE data were generated as part of the PsychENCODE Consortium. Visit 10.7303/syn26365932 for a complete list of grants and PIs. The data were obtained from resource.psychencode.org/.

Study Funding

This research was funded in whole or in part by Aligning Science Across Parkinson's [Grant number: ASAP-000478] through the Michael J. Fox Foundation for Parkinson's Research (MJFF).

Disclosure

H.R.M. reports paid consultancy from Roche; research grants from Parkinson's UK, Cure Parkinson's Trust, PSP Association, CBD Solutions, Drake Foundation, Medical Research Council (MRC), Michael J. Fox Foundation; and is a co-applicant on a patent application related to C9ORF72 - Method for diagnosing a neurodegenerative disease (PCT/GB2012/052140). D.G.G. has received grants from Michael's Movers, the Neurosciences Foundation, and Parkinson's UK; honoraria from AbbVie, BIAL Pharma, Britannia Pharmaceuticals, and GE Health care; and consultancy fees from Acorda Therapeutics and the Glasgow Memory Clinic. M.T.M.H. received funding/grant support from Parkinson's UK, Oxford NIHR BRC, University of Oxford, CPT, Lab10X, NIHR, Michael J. Fox Foundation, H2020 European Union, GE Health care, and the PSP Association. She also received payment for Advisory Board attendance/consultancy for Biogen, Roche, Sanofi, CuraSen Therapeutics, Evidera, Manus Neurodynamica, Lundbeck. Y.B.-S. has received grant funding from the MRC, NIHR, Parkinson's UK, NIH, and ESRC. C.C. receives salary from University Hospitals Plymouth NHS Trust and National Institute of Health and Care Research. She has received advisory, consulting, or lecture fees from AbbVie, Bial, Scient, Orkyn, Abidetex, UCB, Pfizer, EverPharma, Lundbeck, Global Kinetics, Kyowa Kirin, Britannia, and Medscape and research funding from Parkinson's UK, Edmond J. Safra Foundation, National Institute of Health and Care Research, and Cure Parkinson's. J.C.C. has served on advisory boards for Biogen, Denali, Idorsia, Prevail Therapeutic, Servier, Theranexus, and UCB and received grants from Sanofi and the Michael J. Fox Foundation outside of this work. A.E. received funding/grant support by Agence Nationale de la Recherche, France Parkinson, and the Michael J. Fox foundation. J.H. is supported by the UK Dementia Research Institute, which receives its funding from DRI Ltd, funded by the UK Medical Research Council, Alzheimer's Society, and Alzheimer's

Research UK. He is also supported by the MRC, Wellcome Trust, Dolby Family Fund, National Institute for Health Research University College London Hospitals Biomedical Research Centre. All other authors report no competing interests. Go to [Neurology.org/NG](https://www.neurology.org/NG) for full disclosures.

Publication History

Previously published in medRxiv doi: <https://doi.org/10.1101/2022.10.28.22281645>. Received by *Neurology: Genetics* January 17, 2023. Accepted in final form June 8, 2023. Submitted and externally peer reviewed. The handling editor was Stefan M. Pulst, MD, Dr med, FAAN.

Appendix Authors

Name	Location	Contribution
Alejandro Martínez Carrasco, MSc	Department of Clinical and Movement Neurosciences, UCL Queen Square Institute of Neurology; UCL Movement Disorders Centre, University College London, United Kingdom; Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy Chase, MD	Drafting/revision of the manuscript for content, including medical writing for content; study concept or design; analysis or interpretation of data
Raquel Real, PhD	Department of Clinical and Movement Neurosciences, UCL Queen Square Institute of Neurology; UCL Movement Disorders Centre, University College London, United Kingdom; Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy Chase, MD	Drafting/revision of the manuscript for content, including medical writing for content; analysis or interpretation of data
Michael Lawton, PhD	Population Health Sciences, Bristol Medical School, University of Bristol, United Kingdom	Drafting/revision of the manuscript for content, including medical writing for content; analysis or interpretation of data
Regina Hertfelder Reynolds, PhD	Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy Chase, MD; Genetics and Genomic Medicine, UCL Great Ormond Street Institute of Child Health, University College London, United Kingdom	Analysis or interpretation of data
Manuela Tan, PhD	Department of Neurology, Oslo University Hospital, Norway	Drafting/revision of the manuscript for content, including medical writing for content; analysis or interpretation of data
Lesley Wu, MSc	Department of Clinical and Movement Neurosciences, UCL Queen Square Institute of Neurology; UCL Movement Disorders Centre, University College London, United Kingdom; Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy Chase, MD	Major role in the acquisition of data

Appendix (continued)

Name	Location	Contribution
Nigel Williams, PhD	Institute of Psychological Medicine and Clinical Neurosciences, MRC Centre for Neuropsychiatric Genetics and Genomics, Cardiff University, United Kingdom	Analysis or interpretation of data
Camille Carroll, MD	Faculty of Health, University of Plymouth, Plymouth, United Kingdom	Drafting/revision of the manuscript for content, including medical writing for content; major role in the acquisition of data
Jean-Christophe Corvol, MD, PhD	Sorbonne Université, Institut du Cerveau - Paris Brain Institute - ICM, INSERM, CNRS; Assistance Publique Hôpitaux de Paris, Department of Neurology, Hôpital Pitié-Salpêtrière, Paris, France	Drafting/revision of the manuscript for content, including medical writing for content; major role in the acquisition of data
Michele Hu, PhD	Division of Clinical Neurology, Nuffield Department of Clinical Neurosciences; Oxford Parkinson's Disease Centre, University of Oxford, UK	Major role in the acquisition of data
Donald Grosset, MD	School of Neuroscience and Psychology, University of Glasgow, United Kingdom	Drafting/revision of the manuscript for content, including medical writing for content; major role in the acquisition of data
John Hardy, PhD	Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy Chase, MD; Department of Neurodegenerative Diseases, UCL Queen Square Institute of Neurology; UK Dementia Research Institute, University College London; Reta Lila Weston Institute, UCL Queen Square Institute of Neurology; National Institute for Health Research (NIHR) University College London Hospitals Biomedical Research Centre, United Kingdom; Institute for Advanced Study, The Hong Kong University of Science and Technology, Hong Kong SAR, China	Drafting/revision of the manuscript for content, including medical writing for content; study concept or design
Mina Ryten, PhD	Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy Chase, MD; Genetics and Genomic Medicine, UCL Great Ormond Street Institute of Child Health; NIHR Great Ormond Street Hospital Biomedical Research Centre, University College London, United Kingdom	Analysis or interpretation of data

Continued

Appendix (continued)

Name	Location	Contribution
Yoav Ben-Shlomo, PhD	Population Health Sciences, Bristol Medical School, University of Bristol, United Kingdom	Analysis or interpretation of data
Maryam Shoai, PhD	Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy Chase, MD; Department of Neurodegenerative Diseases, UCL Queen Square Institute of Neurology; UK Dementia Research Institute, University College London; Reta Lila Weston Institute, UCL Queen Square Institute of Neurology, London, United Kingdom	Drafting/revision of the manuscript for content, including medical writing for content; study concept or design; analysis or interpretation of data
Huw R. Morris, PhD	Department of Clinical and Movement Neurosciences, UCL Queen Square Institute of Neurology; UCL Movement Disorders Centre, University College London, United Kingdom; Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy Chase, MD	Drafting/revision of the manuscript for content, including medical writing for content; major role in the acquisition of data; study concept or design; analysis or interpretation of data

References

- Lees AJ, Hardy J, Revesz T. Parkinson's disease. *Lancet*. 2009;373(9680):2055-2066.
- Wickremaratchi MM, Knipe MDW, Sastry BSD, et al. The motor phenotype of Parkinson's disease in relation to age at onset. *Mov Disord*. 2011;26(3):457-463.
- Wickremaratchi MM, Ben-Shlomo Y, Morris HR. The effect of onset age on the clinical features of Parkinson's disease. *Eur J Neurol*. 2009;16(4):450-456.
- de la Fuente-Fernández R, Schulzer M, Kuramoto L, et al. Age-specific progression of nigrostriatal dysfunction in Parkinson's disease. *Ann Neurol*. 2011;69(5):803-810.
- Nalls MA, Blauwendraat C, Vallerga CL, et al. Identification of novel risk loci, causal insights, and heritable risk for Parkinson's disease: a meta-analysis of genome-wide association studies. *Lancet Neurol*. 2019;18(12):1091-1102.
- Blauwendraat C, Heilbron K, Vallerga CL, et al. Parkinson's disease age at onset genome-wide association study: defining heritability, genetic loci, and α -synuclein mechanisms. *Mov Disord*. 2019;34(6):866-875.
- Klein C, Westenberger A. Genetics of Parkinson's disease. *Cold Spring Harb Perspect Med*. 2012;2(1):a008888. doi:10.1101/cshperspect.a008888
- Deng H, Wang P, Jankovic J. The genetics of Parkinson disease. *Ageing Res Rev*. 2018;42:72-85.
- Iwaki H, Blauwendraat C, Leonard HL, et al. Genomewide association study of Parkinson's disease clinical biomarkers in 12 longitudinal patients' cohorts. *Mov Disord*. 2019;34(12):1839.
- Tan MMX, Lawton MA, Jabbari E, et al. Genome-wide association studies of cognitive and motor progression in Parkinson's disease. *Mov Disord*. 2021;36(2):424-433.
- Goetz CG, Tilley BC, Shaftman SR, et al. Movement disorder society-sponsored revision of the unified Parkinson's disease rating scale (MDS-UPDRS): scale presentation and clinimetric testing results. *Mov Disord*. 2008;23(15):2129-2170.
- Henderson EJ, Lord SR, Brodie MA, et al. Rivastigmine for gait stability in patients with Parkinson's disease (ReSPonD): a randomised, double-blind, placebo-controlled, phase 2 trial. *Lancet Neurol*. 2016;15(3):249-258.
- Zeng Z, Wang L, Shi W, et al. Effects of unilateral stimulation in Parkinson's disease: a randomized double-blind crossover trial. *Front Neurol*. 2021;12:812455.
- Goetz CG, Luo S, Wang L, Tilley BC, LaPelle NR, Stebbins GT. Handling missing values in the MDS-UPDRS. *Mov Disord*. 2015;30(12):1632-1638.
- Das S, Forer L, Schönerr S, et al. Next-generation genotype imputation service and methods. *Nat Genet*. 2016;48(10):1284-1287.
- Fahn S, Oakes D, Shoulson I, et al. Levodopa and the progression of Parkinson's disease. *N Engl J Med*. 2004;351(24):2498-2508.
- Luke SG. Evaluating significance in linear mixed-effects models in R. *Behav Res Methods*. 2017;49(4):1494-1502.
- Yuan M, Xu XS, Yang Y, et al. SCEBE: an efficient and scalable algorithm for genome-wide association studies on longitudinal outcomes with mixed-effects modeling. *Brief Bioinform*. 2021;22(3):bbaa130. doi:10.1093/bib/bbaa130
- Willer CJ, Li Y, Abecasis GR. METAL: fast and efficient meta-analysis of genomewide association scans. *Bioinformatics*. 2010;26(17):2190.
- Tseng GC, Ghosh D, Feingold E. Comprehensive literature review and statistical considerations for microarray meta-analysis. *Nucleic Acids Res*. 2012;40(9):3785-3799.
- Andersson R, Gebhard C, Miguel-Escalada I, et al. An atlas of active enhancers across human cell types and tissues. *Nature*. 2014;507(7493):455-461.
- Nott A, Holtman IR, Coufal NG, et al. Brain cell type-specific enhancer-promoter interactome maps and disease - risk association. *Science*. 2019;366(6469):1134-1139. doi:10.1126/science.aay0793
- PsychENCODE Consortium, Akbarian S, Liu C, et al. The PsychENCODE project. *Nat Neurosci*. 2015;18(12):1707-1712.
- Vösa U, Claringbould A, Westra HJ, et al. Large-scale cis- and trans-eQTL analyses identify thousands of genetic loci and polygenic scores that regulate blood gene expression. *Nat Genet*. 2021;53(9):1300.
- de Klein N, Tsai EA, Vochteloo M, et al. Brain expression quantitative trait locus and network analysis reveals downstream effects and putative drivers for brain-related diseases. *Nat Genet*. 2023;55(3):377-388.
- Watanabe K, Taskesen E, van Bochoven A, Posthuma D. Functional mapping and annotation of genetic associations with FUMA. *Nat Commun*. 2017;8(1):1826.
- Pruim RJ, Welch RP, Sanna S, et al. LocusZoom: regional visualization of genome-wide association scan results. *Bioinformatics*. 2010;26(18):2336-2337.
- Holden SK, Finseth T, Sillau SH, Berman BD. Progression of MDS-UPDRS scores over five years in de novo Parkinson disease from the Parkinson's progression Markers initiative cohort. *Mov Disord Clin Pract*. 2018;5(1):47-53.
- Malek N, Kanavou S, Lawton MA, et al. L-dopa responsiveness in early Parkinson's disease is associated with the rate of motor progression. *Parkinsonism Relat Disord*. 2019;65:55-61.
- Hiroshima M, Takenawa T. Isolation of a cDNA encoding human lysophosphatidic acid phosphatase that is involved in the regulation of mitochondrial lipid biosynthesis*. *J Biol Chem*. 1999;274(41):29172-29180.
- ACP6 Protein Expression Summary - the Human Protein Atlas. Accessed July 14, 2022. proteinatlas.org/ENSG00000162836-ACP6.
- Schapira AHV. Mitochondria in the aetiology and pathogenesis of Parkinson's disease. *Lancet Neurol*. 2008;7(1):97-109.
- Jankovic J, McDermott M, Carter J, et al. Variable expression of Parkinson's disease: a base-line analysis of the DATATOP cohort. The Parkinson Study Group. *Neurology*. 1990;40(10):1529-1534.
- Schiess MC, Zheng H, Soukup VM, Bonnen JG, Nauta HJ. Parkinson's disease subtypes: clinical classification and ventricular cerebrospinal fluid analysis. *Parkinsonism Relat Disord*. 2000;6(2):69-76.
- Sauerbier A, Jenner P, Todorova A, Chaudhuri KR. Non motor subtypes and Parkinson's disease. *Parkinsonism Relat Disord*. 2016;22(suppl 1):S41-S46.
- Erro R, Picillo M, Vitale C, et al. Clinical clusters and dopaminergic dysfunction in de novo Parkinson disease. *Parkinsonism Relat Disord*. 2016;28:137-140.
- Lewis SJG, Foltynie T, Blackwell AD, Robbins TW, Owen AM, Barker RA. Heterogeneity of Parkinson's disease in the early clinical stages using a data driven approach. *J Neurol Neurosurg Psychiatry*. 2005;76(3):343-348.
- van Rooden SM, Colas F, Martínez-Martín P, et al. Clinical subtypes of Parkinson's disease. *Mov Disord*. 2011;26(1):51-58.
- Müller MLTM, Bohnen NI. Cholinergic dysfunction in Parkinson's disease. *Curr Neurol Neurosci Rep*. 2013;13(9):377.
- Lee SH, Lee MJ, Lyoo CH, Cho H, Lee MS. Impaired finger dexterity and nigrostriatal dopamine loss in Parkinson's disease. *J Neural Transm (Vienna)*. 2018;125(9):1333-1339.
- Evers LJW, Krijthe JH, Meinders MJ, Bloem BR, Heskes TM. Measuring Parkinson's disease over time: the real-world within-subject reliability of the MDS-UPDRS. *Mov Disord*. 2019;34(10):1480-1487.