

2022-10-20

Summing Feynman diagrams in the worldline formalism

Edwards, J

<https://pearl.plymouth.ac.uk/handle/10026.1/21797>

10.22323/1.416.0052

Proceedings of Science

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

Summing Feynman diagrams in the worldline formalism

N. Ahmadiniaz,^a J.P. Edwards,^b C. Lopez-Arcos,^c M.A. Lopez-Lopez,^a C. Moctezuma Mata,^d J. Nicasio^d and C. Schubert^{e,*}

^a*Helmholtz-Zentrum Dresden-Rossendorf, Bautzner Landstraße 400, 01328 Dresden, Germany*

^b*Centre for Mathematical Sciences, University of Plymouth, Plymouth, PL4 8AA, UK*

^c*Escuela de Matemáticas, Universidad Nacional de Colombia Sede Medellín, Carrera 65 59A-110, Medellín, Colombia*

^d*Instituto de Física y Matemáticas Universidad Michoacana de San Nicolás de Hidalgo Edificio C-3, Apdo. Postal 2-82 C.P. 58040, Morelia, Michoacán, México*

^e*Centro Internacional de Ciencias A.C., Campus UNAM-UAEM, Cuernavaca, Morelos, México*

E-mail: christianschubert137@gmail.com

The worldline formalism shares with string theory the property that it allows one to write down master integrals that effectively combine the contributions of many Feynman diagrams. While at the one-loop level these diagrams differ only by the position of the external legs along a fixed line or loop, at multiloop they generally involve different topologies. Here we summarize various efforts that have been made over the years to exploit this property in a computationally meaningful way. As a first example, we show how to generalize the Landau-Khalatnikov-Fradkin formula for the non-perturbative gauge transformation of the fermion propagator in QED to the general $2n$ - point case by pure manipulations at the path-integral level. At the parameter-integral level, we show how to integrate out individual photons in the low-energy expansion, and then sketch a recently introduced general framework for the analytical evaluation of such worldline integrals involving a reduction to quantum mechanics on the circle and the relation between inverse derivatives and Bernoulli polynomials.

*Loops and Legs in Quantum Field Theory - LL2022,
25-30 April, 2022
Ettal, Germany*

*Speaker

1. QED in the worldline representation

1.1 Worldline representation of dressed scalar propagator

Let us start with Feynman's 1950 worldline path integral representation [1] of the Green's function for the interacting Klein-Gordon operator $-(\partial + ieA)^2 + m^2$,

$$\begin{aligned} D^{xx'}[A] &= \langle x' | \int_0^\infty dT \exp \left[-T(-(\partial + ieA)^2 + m^2) \right] | x \rangle \\ &= \int_0^\infty dT e^{-m^2 T} \int_{x(0)=x'}^{x(T)=x} \mathcal{D}x(\tau) e^{-\int_0^T d\tau \left(\frac{1}{4} \dot{x}^2 + ie\dot{x} \cdot A(x(\tau)) \right)}. \end{aligned} \quad (1)$$

Choosing the background field as N plane waves, $A^\mu(x(\tau)) = \sum_{i=1}^N \varepsilon_i^\mu e^{ik_i \cdot x(\tau)}$, and Fourier transforming the endpoints, we get a representation of the “photon-dressed propagator” shown in Fig. 1.




Figure 1: The photon-dressed scalar propagator.

Note that the worldline representation provides the complete amplitude in one piece: differently from the Feynman diagrammatic approach, there is neither a need to sum over “crossed” diagrams, nor to distinguish between the linear and the “seagull” vertex couplings of the scalar particle to the Maxwell field.

1.2 Worldline representation of scalar QED effective action

Similarly, the one-loop effective action can be presented in terms of a path integral over all closed loops in spacetime:

$$\begin{aligned} \Gamma[A] &= \int_0^\infty \frac{dT}{T} \text{Tr} \exp \left[-T(-(\partial + ieA)^2 + m^2) \right] \\ &= \int_0^\infty \frac{dT}{T} e^{-m^2 T} \int_{x(0)=x(T)} \mathcal{D}x(\tau) e^{-\int_0^T d\tau \left(\frac{1}{4} \dot{x}^2 + ie\dot{x} \cdot A(x(\tau)) \right)}. \end{aligned} \quad (2)$$

Expanding the field in N plane waves, one gets the full scalar QED one-loop N -photon amplitudes.

1.3 Worldline representation of spinor QED effective action

At the level of the N -photon amplitudes, the transition from scalar to spinor QED can (up to the normalization) simply be made by inserting, under the path integral (2), the “Feynman spin factor” $\text{Spin}[x, A]$,

$$\text{Spin}[x, A] = \text{tr}_\Gamma \mathcal{P} \exp \left[\frac{i}{4} e [\gamma^\mu, \gamma^\nu] \int_0^T d\tau F_{\mu\nu}(x(\tau)) \right]. \quad (3)$$

However, its use implies path ordering, forcing one to fix the ordering of the photons. For the purposes that I am going to discuss here, it will be essential to avoid this. Following Fradkin [2], this can be done replacing the spin factor by a Grassmann path integral,

$$\text{Spin}[x, A] \rightarrow \int \mathcal{D}\psi(\tau) \exp \left[- \int_0^T d\tau \left(\frac{1}{2} \psi \cdot \dot{\psi} - ie\psi^\mu F_{\mu\nu} \psi^\nu \right) \right] \quad (4)$$

where the Lorentz vectors $\psi^\mu(\tau)$ are anticommuting and antiperiodic functions of proper time: $\psi(\tau_1)\psi(\tau_2) = -\psi(\tau_2)\psi(\tau_1)$, $\psi(T) = -\psi(0)$. The main point of the Grassmann approach is to replace the path-ordered exponential by an ordinary exponential.

1.4 Worldline representation of dressed electron propagator

For the dressed fermion propagator, too, worldline representations have been around for decades [2–5]. However, only during the last few years a version has been developed that seems suitable for higher-order state-of-the-art calculations [6, 7]. The starting point is the second-order representation of the x -space Dirac propagator $S^{xx'}[A]$ in a Maxwell background:

$$\begin{aligned} S^{xx'}[A] &= [m + iD'] K^{xx'}[A] \\ K^{xx'}[A] &= \left\langle x' \left| \left[m^2 - D_\mu D^\mu + \frac{i}{2} e\gamma^\mu \gamma^\nu F_{\mu\nu} \right]^{-1} \right| x \right\rangle \\ &= \int_0^\infty dT e^{-m^2 T} e^{-\frac{1}{4} \frac{(x-x')^2}{T}} \int_{q(0)=0}^{q(T)=0} Dq e^{-\int_0^T d\tau \left(\frac{1}{4} \dot{q}^2 + ie\dot{q} \cdot A + ie \frac{x'-x}{T} \cdot A \right)} \\ &\quad \times 2^{-\frac{D}{2}} \text{symb}^{-1} \int_{\psi(0)+\psi(T)=0} D\psi e^{-\int_0^T d\tau \left[\frac{1}{2} \psi_\mu \dot{\psi}^\mu - ieF_{\mu\nu} (\psi+\eta)^\mu (\psi+\eta)^\nu \right]}. \end{aligned} \quad (5)$$

Here η^μ is an external Grassmann Lorentz vector, and the “symbol map” *symb* converts products of η s into fully antisymmetrised products of Dirac matrices:

$$\text{symb}(\gamma^{\alpha_1 \alpha_2 \dots \alpha_n}) \equiv (-i\sqrt{2})^n \eta^{\alpha_1} \eta^{\alpha_2} \dots \eta^{\alpha_n} \quad (6)$$

where $\gamma^{\alpha\beta\dots\rho}$ denotes the totally antisymmetrised product of gamma matrices:

$$\gamma^{\alpha_1 \alpha_2 \dots \alpha_n} \equiv \frac{1}{n!} \sum_{\pi \in S_n} \text{sign}(\pi) \gamma^{\alpha_{\pi(1)}} \gamma^{\alpha_{\pi(2)}} \dots \gamma^{\alpha_{\pi(n)}}. \quad (7)$$

1.5 Higher order QED processes

Since all the above formulas are valid off-shell, arbitrary QED processes can be constructed from these building blocks by sewing (Fig. 2).

Although the worldline representation is equivalent to Feynman diagrams, it is more global in the sense that it does not distinguish between diagrams differing only by the ordering of the photon legs along a line or loop.




Figure 2: A typical multiloop process in QED.

2. Generalization of the Landau-Khalatnikov-Fradkin transformation

For some purposes, this property becomes useful already at the path-integral level. In 1956, Landau and Khalatnikov [8] and independently Fradkin [9] showed, that the non-perturbative behaviour of the electron propagator $S(x; \xi)$ under a change $\xi \rightarrow \hat{\xi}$ of the covariant gauge parameter ξ can be written as (in exactly four dimensions)

$$S(x; \xi) = S(x; \hat{\xi}) \left[\frac{x^2}{x_0^2} \right]^{-\frac{\alpha}{4\pi}(\xi - \hat{\xi})} \quad (8)$$

where x_0 is an IR cutoff. Very recently, we have used the worldline formalism to extend this result to the general fermionic $2n$ -point correlator

$$\mathcal{A}(x_1, \dots, x_n; x'_1, \dots, x'_n | \xi) \equiv \langle \psi(x_1) \cdots \psi(x_n) \bar{\psi}(x'_1) \cdots \bar{\psi}(x'_n) \rangle \quad (9)$$

as follows [10]:

$$\mathcal{A}(x_1, \dots, x_n; x'_1, \dots, x'_n | \hat{\xi}) = \prod_{k,l=1}^n e^{(\hat{\xi} - \xi) S^{(k,l)}} \mathcal{A}(x_1, \dots, x_n; x'_1, \dots, x'_n | \xi), \quad (10)$$

$$S^{(k,l)} = \frac{e^2}{32\pi^{\frac{D}{2}}} \Gamma\left(\frac{D}{2} - 2\right) \left\{ \left[(x_k - x_l)^2 \right]^{2 - \frac{D}{2}} - \left[(x_k - x'_l)^2 \right]^{2 - \frac{D}{2}} - \left[(x'_k - x_l)^2 \right]^{2 - \frac{D}{2}} + \left[(x'_k - x'_l)^2 \right]^{2 - \frac{D}{2}} \right\}. \quad (11)$$

Here the sum k, l runs over all pairs of open fermion lines, and the exponential factor for fixed k and l implements the effect of the change of the covariant gauge parameter for photons inserted between them in all possible ways (for photons that on one or both ends hit a closed loop there is no effect). Note also that the dimension has been left general.

3. String-inspired treatment of the worldline path integral

Under the influence of string theory, in the nineties a perturbative approach to the evaluation of worldline path integrals using worldline Green's functions was developed [11, 12]. E.g. for the closed-loop case one has the basic correlators

$$\langle x^\mu(\tau_1) x^\nu(\tau_2) \rangle = -G(\tau_1, \tau_2) \delta^{\mu\nu}, \quad G(\tau_1, \tau_2) = |\tau_1 - \tau_2| - \frac{1}{T} \left(\tau_1 - \tau_2 \right)^2, \quad (12)$$

$$\langle \psi^\mu(\tau_1) \psi^\nu(\tau_2) \rangle = G_F(\tau_1, \tau_2) \delta^{\mu\nu}, \quad G_F(\tau_1, \tau_2) = \text{sign}(\tau_1 - \tau_2). \quad (13)$$

This allows one to derive compact master formulas for the photon-dressed propagators and the one-loop N -photon amplitudes in scalar and spinor QED, as well as for many other types of amplitudes (for reviews, see [13, 14]).

3.1 Master formula for the N -photon amplitudes in scalar and spinor QED

The in many ways prototypical one of these master formulas is the one for the one-loop N -photon amplitudes in scalar QED:

$$\begin{aligned} \Gamma[\{k_i, \varepsilon_i\}] &= (-ie)^N \int_0^\infty \frac{dT}{T} (4\pi T)^{-\frac{D}{2}} e^{-m^2 T} \prod_{i=1}^N \int_0^T d\tau_i \\ &\times \exp \left\{ \sum_{i,j=1}^N \left[\frac{1}{2} G_{ij} k_i \cdot k_j + i \dot{G}_{ij} k_i \cdot \varepsilon_j + \frac{1}{2} \ddot{G}_{ij} \varepsilon_i \cdot \varepsilon_j \right] \right\} |_{\text{lin}(\varepsilon_1, \dots, \varepsilon_N)} . \end{aligned} \quad (14)$$

Here T is the loop proper-time and τ_i parametrizes the position of photon i along the loop. A projection on the terms linear in each of the polarization vectors $\varepsilon_1, \dots, \varepsilon_N$ is understood. Besides the Green's function $G_{ij} \equiv G(\tau_i, \tau_j)$ also its first and second derivatives appear,

$$\dot{G}(\tau_1, \tau_2) = \text{sign}(\tau_1 - \tau_2) - 2 \frac{(\tau_1 - \tau_2)}{T} , \quad (15)$$

$$\ddot{G}(\tau_1, \tau_2) = 2\delta(\tau_1 - \tau_2) - \frac{2}{T} . \quad (16)$$

A similar master formula can be written down for the spinor loop using worldline supersymmetry.. However, in practice it is usually preferable to use an integration-by-parts that removes all the \ddot{G}_{ij} , and the following ‘‘Bern-Kosower replacement rule’’ [11]:

$$\dot{G}_{i_1 i_2} \dot{G}_{i_2 i_3} \cdots \dot{G}_{i_n i_1} \rightarrow \dot{G}_{i_1 i_2} \dot{G}_{i_2 i_3} \cdots \dot{G}_{i_n i_1} - G_{F i_1 i_2} G_{F i_2 i_3} \cdots G_{F i_n i_1} . \quad (17)$$

3.2 The four-photon amplitudes

Let us write down the resulting representation for the familiar four-photon amplitude, usually given in terms of the six Feynman diagrams displayed in Fig. 3.




Figure 3: Six Feynman diagrams for photon-photon scattering.

After a large number of integrations by parts, one finds the following tensor decomposition [15] (still off-shell):

$$\hat{\Gamma} = \hat{\Gamma}^{(1)} + \hat{\Gamma}^{(2)} + \hat{\Gamma}^{(3)} + \hat{\Gamma}^{(4)} + \hat{\Gamma}^{(5)} , \quad (18)$$

$$\begin{aligned}
\hat{\Gamma}^{(1)} &= \hat{\Gamma}_{(1234)}^{(1)} T_{(1234)}^{(1)} + \hat{\Gamma}_{(1243)}^{(1)} T_{(1243)}^{(1)} + \hat{\Gamma}_{(1324)}^{(1)} T_{(1324)}^{(1)}, \\
\hat{\Gamma}^{(2)} &= \hat{\Gamma}_{(12)(34)}^{(2)} T_{(12)(34)}^{(2)} + \hat{\Gamma}_{(13)(24)}^{(2)} T_{(13)(24)}^{(2)} + \hat{\Gamma}_{(14)(23)}^{(2)} T_{(14)(23)}^{(2)}, \\
\hat{\Gamma}^{(3)} &= \sum_{i=1,2,3} \hat{\Gamma}_{(123)i}^{(3)} T_{(123)i}^{(3)r_4} + \sum_{i=2,3,4} \hat{\Gamma}_{(234)i}^{(3)} T_{(234)i}^{(3)r_1} + \sum_{i=3,4,1} \hat{\Gamma}_{(341)i}^{(3)} T_{(341)i}^{(3)r_2} + \sum_{i=4,1,2} \hat{\Gamma}_{(412)i}^{(3)} T_{(412)i}^{(3)r_3}, \\
\hat{\Gamma}^{(4)} &= \sum_{i<j} \hat{\Gamma}_{(ij)ii}^{(4)} T_{(ij)ii}^{(4)} + \sum_{i<j} \hat{\Gamma}_{(ij)jj}^{(4)} T_{(ij)jj}^{(4)}, \\
\hat{\Gamma}^{(5)} &= \sum_{i<j} \hat{\Gamma}_{(ij)ij}^{(5)} T_{(ij)ij}^{(5)} + \sum_{i<j} \hat{\Gamma}_{(ij)ji}^{(5)} T_{(ij)ji}^{(5)}.
\end{aligned}$$

Remarkably, the basis of five tensors $T^{(i)}$ is identical with the one found in 1971 by Costantini, De Tollis and Pistoni [16] using the QED Ward identity:

$$T_{(1234)}^{(1)} \equiv Z(1234), \quad (19)$$

$$T_{(12)(34)}^{(2)} \equiv Z(12)Z(34), \quad (20)$$

$$T_{(123)i}^{(3)r_4} \equiv Z(123) \frac{r_4 \cdot f_4 \cdot k_i}{r_4 \cdot k_4} \quad (i = 1, 2, 3), \quad (21)$$

$$T_{(12)11}^{(4)} \equiv Z(12) \frac{k_1 \cdot f_3 \cdot f_4 \cdot k_1}{k_3 \cdot k_4}, \quad (22)$$

$$T_{(12)12}^{(5)} \equiv Z(12) \frac{k_1 \cdot f_3 \cdot f_4 \cdot k_2}{k_3 \cdot k_4}. \quad (23)$$

Here we have further introduced

$$f_i^{\mu\nu} \equiv k_i^\mu \varepsilon_i^\nu - \varepsilon_i^\mu k_i^\nu, \quad (24)$$

$$Z(ij) \equiv \frac{1}{2} \text{tr}(f_i f_j) = \varepsilon_i \cdot k_j \varepsilon_j \cdot k_i - \varepsilon_i \cdot \varepsilon_j k_i \cdot k_j, \quad (25)$$

$$Z(i_1 i_2 \dots i_n) \equiv \text{tr} \left(\prod_{j=1}^n f_{i_j} \right) \quad (n \geq 3). \quad (26)$$

The corresponding parameter integrals are extremely compact:

$$\hat{\Gamma}^{(k)}_{\dots} = \int_0^\infty \frac{dT}{T} T^{4-\frac{D}{2}} e^{-m^2 T} \int_0^1 \prod_{i=1}^4 du_i \hat{\gamma}^{(k)}_{\dots}(\dot{G}_{ij}) e^{T \sum_{i<j=1}^4 G_{ij} k_i \cdot k_j} \quad (27)$$

where, for spinor QED,

$$\hat{\gamma}_{(1234)}^{(1)} = \dot{G}_{12} \dot{G}_{23} \dot{G}_{34} \dot{G}_{41} - G_{F12} G_{F23} G_{F34} G_{F41}, \quad (28)$$

$$\hat{\gamma}_{(12)(34)}^{(2)} = (\dot{G}_{12} \dot{G}_{21} - G_{F12} G_{F21}) (\dot{G}_{34} \dot{G}_{43} - G_{F34} G_{F43}), \quad (29)$$

$$\hat{\gamma}_{(123)i}^{(3)} = (\dot{G}_{12} \dot{G}_{23} \dot{G}_{31} - G_{F12} G_{F23} G_{F31}) \dot{G}_{4i}, \quad (30)$$

$$\hat{\gamma}_{(12)11}^{(4)} = (\dot{G}_{12} \dot{G}_{21} - G_{F12} G_{F21}) \dot{G}_{13} \dot{G}_{41}, \quad (31)$$

$$\hat{\gamma}_{(12)12}^{(5)} = (\dot{G}_{12} \dot{G}_{21} - G_{F12} G_{F21}) \dot{G}_{13} \dot{G}_{42} \quad (32)$$

(plus permutations thereof), and the coefficient functions for scalar QED are obtained from these simply by deleting all the G_{Fij} .

3.3 Master formula for the N-photon dressed scalar propagator

A master formula for the scalar propagator dressed with N photons was obtained by Daikouji et al. in 1996 [17]:

$$D^{pp'}(k_1, \varepsilon_1; \dots; k_N, \varepsilon_N) = (-ie)^N \int_0^\infty dT e^{-m^2 T} \times \prod_{i=1}^N \int_0^T d\tau_i e^{-Tb^2 + \sum_{i,j=1}^N [\Delta_{ij} k_i \cdot k_j - 2i \bullet \Delta_{ij} \varepsilon_i \cdot k_j - \bullet \Delta_{ij}^* \varepsilon_i \cdot \varepsilon_j]} \Big|_{\varepsilon_1 \varepsilon_2 \dots \varepsilon_N}. \quad (33)$$

Here we have introduced the vector $b \equiv p' + \frac{1}{T} \sum_{i=1}^N (k_i \tau_i - i \varepsilon_i)$ and a different worldline Green's function $\Delta(\tau, \tau')$,

$$\langle q^\mu(\tau) q^\nu(\tau') \rangle = -2\Delta(\tau, \tau') \delta^{\mu\nu}, \quad \Delta(\tau, \tau') = \frac{|\tau - \tau'|}{2} - \frac{\tau + \tau'}{2} + \frac{\tau \tau'}{T}. \quad (34)$$

3.4 Master formula for the photon-dressed electron propagator

To the contrary, a master formula for the photon-dressed electron propagator was obtained only very recently [6, 7]:

$$K_N^{pp'}(k_1, \varepsilon_1; \dots; k_N, \varepsilon_N) = (-ie)^N \text{symb}^{-1} \int_0^\infty dT e^{-m^2 T} \int_0^T d\tau_1 \dots \int d\theta_N \times e^{-\sqrt{2}\eta \cdot \sum_{i=1}^N (\varepsilon_i + i\theta k_i) + \sum_{i,j=0}^{N+1} [\hat{g}_{ij} k_i \cdot k_j + 2i D_i \hat{g}_{ij} \varepsilon_i \cdot k_j + D_i D_j \hat{g}_{ij} \varepsilon_i \cdot \varepsilon_j]} \Big|_{\varepsilon_1 \dots \varepsilon_N}. \quad (35)$$

This now involves Grassmann variables $\theta_1, \dots, \theta_N$, the super derivative $D = \frac{\partial}{\partial \theta} - \theta \frac{\partial}{\partial \tau}$ and the super worldline Green's function $\hat{g}(\tau, \theta; \tau', \theta') = \frac{1}{2}(|\tau - \tau'| + \theta \theta' \text{sign}(\tau - \tau'))$. The main advantages of this approach are that (i) the use of the symbol map leads to an early projection on the Clifford basis, effectively avoiding the appearance of long Dirac traces (ii) the spin-averaging can be done without fixing the number or helicity assignments of the photons. As a check, in [7] the formalism was used to recalculate the (polarized and unpolarized) Compton scattering amplitude, and complete agreement was found with [18].

3.5 On to multiloop

Dealing with the amplitude as a whole becomes important when one uses the one-loop amplitudes to construct higher-loop amplitudes by sewing. For example, from the one-loop six-photon amplitude we can construct the three-loop quenched propagator (Fig. 4) etc.




Figure 4: The quenched three-loop photon propagator.

This type of sums of diagrams is known to suffer from extensive cancellations.

4. The fundamental problem of worldline integration

Returning to the one-loop level, considering that G_{Fij} s can always be eliminated by

$$G_{Fij}G_{Fjk}G_{Fki} = -(\dot{G}_{ij} + \dot{G}_{jk} + \dot{G}_{ki}) \quad (36)$$

the most general integral that one will ever have to compute in the worldline approach to QED (or any abelian theory) is of the form

$$\int_0^1 du_1 du_2 \cdots du_N \text{Pol}(\dot{G}_{ij}) e^{\sum_{i < j=1}^N G_{ij} k_i \cdot k_j} \quad (37)$$

with arbitrary N and polynomial $\text{Pol}(\dot{G}_{ij})$. Here we have rescaled $\tau_i = Tu_i$, so that now

$$G_{ij} = |u_i - u_j| - (u_i - u_j)^2, \quad \dot{G}_{ij} = \text{sign}(u_i - u_j) - 2(u_i - u_j). \quad (38)$$

Ideally, we would like to compute this integral without decomposing the integrand into ordered sectors. This leads to a very non-standard integration problem. An easily solvable special case are “cycle integrals”:

$$b_n \equiv \int_0^1 du_1 du_2 \cdots du_n \dot{G}_{12} \dot{G}_{23} \cdots \dot{G}_{n1} = \begin{cases} -2^n \frac{B_n}{n!} & n \text{ even} \\ 0 & n \text{ odd} \end{cases} \quad (39)$$

where B_n denotes the n th Bernoulli number, and “super cycle integrals”,


$$b_n - f_n \equiv \int_0^1 du_1 du_2 \cdots du_n \left(\dot{G}_{B12} \dot{G}_{B23} \cdots \dot{G}_{Bn1} - G_{F12} G_{F23} \cdots G_{Fn1} \right) = (2 - 2^n) b_n. \quad (40)$$

In the worldline formalism, those are all that is needed to calculate the one-loop N -photon amplitudes in the low-energy approximation [19].

General polynomial integrals can be done recursively by the application of the following formula [20]

$$\begin{aligned} \int_0^1 du \dot{G}(u, u_1)^{k_1} \dot{G}(u, u_2)^{k_2} \cdots \dot{G}(u, u_n)^{k_n} &= \frac{1}{2n} \sum_{i=1}^n \prod_{j \neq i} \sum_{l_j=0}^{k_j} \binom{k_j}{l_j} \dot{G}_{ij}^{k_j - l_j} \sum_{l_i=0}^{k_i} \binom{k_i}{l_i} \\ &\times \frac{(-1)^{\sum_{j=1}^n l_j}}{(1 + \sum_{j=1}^n l_j) n^{\sum_{j=1}^n l_j}} \left\{ \left(\sum_{j \neq i} \dot{G}_{ij} + 1 \right)^{1 + \sum_{j=1}^n l_j} - (-1)^{k_i - l_i} \left(\sum_{j \neq i} \dot{G}_{ij} - 1 \right)^{1 + \sum_{j=1}^n l_j} \right\}. \end{aligned} \quad (41)$$

This can be used, for example, to integrate out a low-energy photon in a multiphoton amplitude *without fixing the order of the remaining photons*. In a forthcoming publication [21] this is applied to the four-photon amplitude, creating a building block for higher-loop calculations that will make it possible, for example, to unify the calculation of the various 3-loop g -2 contributions shown in Fig. 5 and the calculation of the 4-loop β -function contributions shown in Fig. 6.

**Figure 5:** LBL contributions to the three-loop $g - 2$.**Figure 6:** LBL contributions to the four-loop β function. The external photons are at low-energy.

5. Inverse derivative expansion

Much more difficult is the inclusion of the universal factors $e^{G_{ij}k_i \cdot k_j}$. The following strategy was proposed in [20]: the worldline path integral is performed in the Hilbert space H'_P of periodic functions orthogonal to the constant functions (because the zero mode must be fixed). In this space the ordinary n th derivative ∂_P is invertible, and the integral kernel of the inverse is given essentially by the n th Bernoulli polynomial $B_n(x)$ [22, 23]:

$$\langle u_1 | \partial_P^{-n} | u_{n+1} \rangle = -\frac{1}{n!} B_n(|u_1 - u_{n+1}|) \text{sign}^n(u_1 - u_{n+1}) \quad (n \geq 1), \quad (42)$$

$$\langle u_i | \partial^0 | u_j \rangle = \delta(u_i - u_j) - 1. \quad (43)$$

Thus worldline integration naturally relates to the theory of Bernoulli numbers and polynomials. Further, in [20] it was shown how to expand the universal exponential factor in terms of the matrix elements of the inverse derivatives:

$$e^{G_{ij}k_i \cdot k_j} = 1 + 2 \sum_{n=1}^{\infty} (k_i \cdot k_j)^{n-1/2} H_{2n-1} \left(\frac{\sqrt{k_i \cdot k_j}}{2} \right) (\langle u_i | \partial^{-2n} | u_j \rangle - \langle u_i | \partial^{-2n} | u_i \rangle) \quad (44)$$

where the $H_n(x)$ are Hermite polynomials. Let us show how to use this expansion for the simplest case of the N -point function in ϕ^3 theory, where the master formula simply becomes ($k_{ij} \equiv k_i \cdot k_j$)

$$\hat{I}_N(k_1, \dots, k_N) = \int_0^\infty \frac{dT}{T} T^{N-D/2} e^{-m^2 T} \int_0^1 du_1 \dots du_N \exp \left[T \sum_{i < j=1}^N G_{ij} k_{ij} \right]. \quad (45)$$

In the three-point case one can use this to write ($\hat{B}_n \equiv \frac{B_n}{n!}$)

$$\begin{aligned}
e^{k_{12}G_{12}+k_{13}G_{13}+k_{23}G_{23}} &= \left\{ 1 + 2 \sum_{i=1}^{\infty} k_{12}^{i-\frac{1}{2}} H_{2i-1} \left(\frac{\sqrt{k_{12}}}{2} \right) \left[\langle u_1 | \partial^{-2i} | u_2 \rangle + \hat{B}_{2i} \right] \right\} \\
&\times \left\{ 1 + 2 \sum_{j=1}^{\infty} k_{13}^{j-\frac{1}{2}} H_{2j-1} \left(\frac{\sqrt{k_{13}}}{2} \right) \left[\langle u_1 | \partial^{-2j} | u_3 \rangle + \hat{B}_{2j} \right] \right\} \\
&\times \left\{ 1 + 2 \sum_{k=1}^{\infty} k_{23}^{k-\frac{1}{2}} H_{2k-1} \left(\frac{\sqrt{k_{23}}}{2} \right) \left[\langle u_2 | \partial^{-2k} | u_3 \rangle + \hat{B}_{2k} \right] \right\}.
\end{aligned} \tag{46}$$

Since $\int_0^1 du_{i,j} \langle u_i | \partial^{-2i} | u_j \rangle = 0$, the three $\langle u_i | \partial^{-2n} | u_j \rangle$ can produce a non-vanishing integral only together, and then by $\int_0^1 du |u\rangle \langle u| = \mathbb{1}$,

$$\int_{123} \langle u_1 | \partial^{-2i} | u_2 \rangle \langle u_2 | \partial^{-2k} | u_3 \rangle \langle u_3 | \partial^{-2j} | u_1 \rangle = \text{Tr}(\partial^{-2(i+j+k)}) = -\hat{B}_{2(i+j+k)}. \tag{47}$$

In this way we get a closed form-expression for the $N = 3$ momentum expansion coefficients,

$$\begin{aligned}
I_3(a, b, c) \equiv \int_{123} G_{12}^a G_{13}^b G_{23}^c &= a!b!c! \sum_{i=\lfloor 1+a/2 \rfloor}^a \sum_{j=\lfloor 1+b/2 \rfloor}^b \sum_{k=\lfloor 1+c/2 \rfloor}^c h_i^a h_j^b h_k^c \\
&\times \left(\hat{B}_{2i} \hat{B}_{2j} \hat{B}_{2k} - \hat{B}_{2(i+j+k)} \right).
\end{aligned} \tag{48}$$

Here we have assumed that a, b, c are all different from zero, and the coefficients h_i^a are (from the explicit formula for the Hermite polynomials)

$$h_i^a = (-1)^{a+1} \frac{2(2i-1)!}{(2i-a-1)!(2a-2i+1)!}. \tag{49}$$

At the four-point level, we encounter more general integrals involving the “cubic worldline vertex”

$$V_3^{ijk} \equiv \int_0^1 du \langle u | \partial^{-i} | u_1 \rangle \langle u | \partial^{-j} | u_2 \rangle \langle u | \partial^{-k} | u_3 \rangle \tag{50}$$

but they can be reduced to chain integrals by a systematic IBP procedure. This remains true at higher points.

6. Summary and Outlook

1. In the worldline formalism, we can integrate out photons in the low-energy limit, or to any finite order in the external momentum, without fixing the ordering of the remaining legs.
2. At full momentum, we can use the inverse derivative expansion, and try a resummation. For this purpose, eventually we will need formulas relating the Bernoulli numbers to hypergeometric functions.
3. This provides also a powerful new approach to the calculation of the ϕ^3 and QED heat-kernel expansions.

References

- [1] R. P. Feynman, Phys. Rev. **80** (1950) 440.
- [2] E. S. Fradkin, Nucl. Phys. **76** (1966) 588.
- [3] R. P. Feynman, Phys. Rev. **84** (1951) 108.
- [4] E.S. Fradkin, D.M. Gitman, Phys. Rev. **D 44** (1991) 3230.
- [5] M. Reuter, M. G. Schmidt and C. Schubert, Ann. Phys. (N.Y.) **259** 313 (1997), hep-th/9610191.
- [6] N. Ahmadinia, V.M. Banda Guzmán, F. Bastianelli, O. Corradini, J.P. Edwards and C. Schubert, JHEP **2008** (2020) 049, arXiv:2004.01391 [hep-th].
- [7] N. Ahmadinia, V.M. Banda Guzmán, F. Bastianelli, O. Corradini, J.P. Edwards and C. Schubert, JHEP **01** (2022) 050, arXiv:2107.00199 [hep-th].
- [8] L. Landau, K. I. Khalatnikov., Sov. Phys. JETP **2**, 69 (1956).
- [9] E. S. Fradkin, Zh. Eksp. Teor. Fiz. **29**, 258 (1955).
- [10] N. Ahmadinia, J. P. Edwards, J. Nicasio and C. Schubert, Phys. Rev. D **104** (2021) 2, 025014, e-Print: 2012.10536 [hep-th].
- [11] Z. Bern and D.A. Kosower, Nucl. Phys. **B 379** (1992) 451.
- [12] M. J. Strassler, Nucl. Phys. **B 385** (1992) 145, hep-ph/9205205.
- [13] C. Schubert, Phys. Rept. **355**, 73 (2001), arXiv:hep-th/0101036.
- [14] J. P. Edwards and C. Schubert, arXiv:1912.10004 [hep-th].
- [15] N. Ahmadinia, C. Lopez-Arcos, M. A. Lopez-Lopez and C. Schubert, arXiv:2012.11791 [hep-th].
- [16] V. Costantini, B. de Tollis and G. Pistoni, *Nuov. Cim. A2* (1971) no. 3, 733.
- [17] K. Daikouji, M. Shino, Y. Sumino, Phys. Rev. **D 53** (1996) 4598, hep-ph/9508377.
- [18] A. Denner and S. Dittmaier, Nucl. Phys. B **540** (1999) 58, hep-ph/9805443.
- [19] G. V. Dunne and C. Schubert, JHEP **0208**, 053 (2002), hep-th/0205004.
- [20] J. P. Edwards, C. M. Mata, U. Müller and C. Schubert, SIGMA **17** (2021), 065, 2106.1207.
- [21] N. Ahmadinia, C. Lopez-Arcos, M. A. Lopez-Lopez and C. Schubert, “The QED four – photon amplitudes off-shell: part 3”, in preparation.
- [22] M.G. Schmidt and C. Schubert, Phys. Lett. B **318**, 438 (1993), hep-th/9309055.
- [23] M.G. Schmidt and C. Schubert, Phys. Rev. D **53**, 2150 (1996), hep-th/9410100.