

2023

# Testing the potential use of UK wetland plant species in paludiculture using examples from the Somerset Levels

Bentley, O.

Bentley, O. (2023) 'Testing the potential use of UK wetland plant species in paludiculture using examples from the Somerset Levels', *The Plymouth Student Scientist*, 16(2), pp. 175-198.

<https://pearl.plymouth.ac.uk/handle/10026.1/21840>

---

The Plymouth Student Scientist

University of Plymouth

---

*All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.*

## Appendices

### Appendix 1




Image: Olivia Bentley

Figure 7: Final vegetation sample pellets produced using the XRF press machine.

### Appendix 2




Figure 8: Box plot showing the mean carbon storage capacity per t ha<sup>-1</sup>-yr<sup>-1</sup> of each wetland plant species harvested on the Somerset levels.

### Appendix 3




Figure 9: Grouping information of the seven plant species dry biomass production per t ha<sup>-1</sup>-yr<sup>-1</sup> obtained through a post hoc test using the Fisher LSD method. Pairings which do not a zero within the intervals are statistically significantly different.

### Appendix 4

Table 8: Output of Kruskal Wallis non-parametric test conducted on the phosphorus content (%) of the seven wetland plant species harvested on the Somerset levels.

#### Test

Null hypothesis H<sub>0</sub>: All medians are equal  
 Alternative hypothesis H<sub>1</sub>: At least one median is different

Method	DF	H-Value	P-Value
Not adjusted for ties	6	19.64	0.003
Adjusted for ties	6	19.66	0.003

*The chi-square approximation may not be accurate when some sample sizes are less than 5.*

## Appendix 5

Table 9: Output of Kruskal Wallis non-parametric test conducted on the phosphorus removal capacity ( $t\ ha^{-1}\text{-yr}^{-1}$ ) of the seven wetland plant species harvested on the Somerset levels.

### Test

Null hypothesis  $H_0$ : All medians are equal  
 Alternative hypothesis  $H_1$ : At least one median is different

Method	DF	H-Value	P-Value
Not adjusted for ties	6	12.80	0.046
Adjusted for ties	6	12.83	0.046

*The chi-square approximation may not be accurate when some sample sizes are less than 5.*

## Appendix 6

Table 10: Analysis of variance from a one-way ANOVA conducted on carbon content (%) between six plant species excluding *Typha latifolia*.

### Analysis of Variance

Source	DF	Adj SS	Adj MS	F-Value	P-Value
Sample_1	5	795.881	159.176	426.52	0.00000000000004468
Error	12	4.478	0.373		
Total	17	800.360			

## Appendix 7




Figure 10: Grouping information of the six plant species carbon content (%), excluding *T.latifolia*, obtained through a post hoc test Tukey test. Pairings which do not a zero within the intervals are statistically significantly different.

## Appendix 8




Figure 11: Grouping information of five plant species carbon storage capacity (t ha<sup>-1</sup>-yr<sup>-1</sup>), excluding *T.latifolia* & *J. effusus*, obtained through a post hoc test Tukey test. Pairings which do not a zero within the intervals are statistically significantly different.