

2024-03-07

# mNCEA policy brief - Mind the Gap The need to continue long-term plankton monitoring

Holland, Matthew

Holland, M. et al. (2024). *mNCEA policy brief - Mind the Gap The need to continue long-term plankton monitoring*. PEARL Research Repository <https://doi.org/10.24382/bs9p-6e92>  
<https://pearl.plymouth.ac.uk/handle/10026.1/22151>

---


<https://doi.org/10.24382/bs9p-6e92>  
University of Plymouth

---

*All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.*

## Mind the Gap – The need to continue long-term plankton monitoring

**Briefing report:** While it is beneficial to explore novel plankton survey technology, it is essential that we also continue to maintain traditional long-term monitoring programmes to generate the necessary information to inform policy.




### Highlights

- Changes in plankton have important impacts on ecosystem services and health.
- Such changes can only be detected by studying long-term, consistent plankton datasets.
- Traditional long-term monitoring programmes use light microscopy and provide high quality trusted data.
- We must maintain traditional plankton monitoring to detect and understand the causes and consequences of important changes in pelagic habitats, many of which are occurring on decadal scales.
- Novel technologies offer efficiency benefits and fill some knowledge gaps left by traditional monitoring.
- Novel technologies should continue to develop in parallel with traditional monitoring, as traditional methods are necessary for validation.
- We cannot replace traditional monitoring with novel technology any time soon since we require long-term time-series and detailed taxonomic accounting to detect important changes and links to pressures.

## Summary

Changes in plankton have important implications for the continued provision of ecosystem services, including supporting commercial fish stocks, carbon sequestration, and oxygen production. Such changes can only be detected by studying long-term, consistent plankton datasets which are needed to understand the pressures driving these changes and how we can manage them. Traditional long-term plankton monitoring relies on light microscopy to identify and count plankton taxa, with methods fully supported by national / international QA/QC standards and providing high quality trusted data. Novel technologies, including imaging and molecular methods, offer more efficient means of collecting some types of plankton data, filling targeted knowledge gaps left by traditional monitoring. However, these data are often semi-quantitative, lacking in QA/QC standards, and/or in taxonomic resolution. While these technologies are developed it remains critical to maintain the continuity of traditional plankton monitoring to inform policy assessments of important changes in biodiversity. Losing these time-series, many of which span multiple decades, would impair our ability to detect important change in pelagic habitats, as most changes cannot be detected from short-term data. This would also accelerate the loss of taxonomic expertise, already under threat globally, diminishing our UK skill-base. Novel technologies should be explored in parallel to traditional monitoring, as they can provide complementary data to support policy assessments and research, however, it is important that we do not attempt to replace traditional monitoring with new technology before it has been thoroughly integrated into long-term monitoring programmes.



Traditional methods such as Continuous Plankton Recorder (CPR), nets and Niskin bottles have been used to monitor plankton for decades with many important research outputs. Combining traditional methods with newer methods including in situ imaging, molecular methods, advanced sensors, and satellites can improve spatial and temporal coverage and further our understanding of changes resulting from pressures on plankton communities. Figure designed by Dr Stacey McCormack (Visual Knowledge) and published in {ADDIN EN.CITE



*The Marine Biological Association coordinates the Continuous Plankton Recorder (CPR) survey, employing specialised towed instruments deployed routinely from commercial ships and ferries as they travel their regular routes. These recorders are equipped with a mechanism that allows them to filter and collect plankton samples from seawater as they are towed. The CPR device features a silk filtering mesh that captures plankton as water passes through it, preserving a record of plankton abundance and distribution, biologically fixed in-situ. Once in the laboratory, each roll of CPR silk is carefully unrolled, cut into sections, subsampled, and counted along transects by trained analysts with the help of light microscopes. This innovative mechanical device has remained unchanged since it was first used in 1931. The high frequency and broad spatial coverage of the CPR survey enable scientists to study and analyse changes in plankton communities over time, detecting important changes in pelagic habitats biodiversity and providing crucial insights into the dynamics of marine ecosystems.*

## Plankton monitoring in UK waters

Traditional plankton monitoring data have revealed important large-scale declines in plankton abundance in UK waters and beyond { ADDIN EN.CITE { ADDIN EN.CITE.DATA }}. Here we use the term “traditional plankton monitoring” to refer to micro-phytoplankton and zooplankton samples collected by net, bottle, bucket, or Continuous Plankton Recorder and counted by trained taxonomists using light microscopes. Traditional plankton monitoring provides detailed abundance data which can be applied to address a wide range of questions and applications, including assessments of Good Environmental Status for OSPAR and UK Marine Strategy to inform policy decisions { ADDIN EN.CITE { ADDIN EN.CITE.DATA }}, Marine Climate Change Impacts Partnership (MCCIP) report cards to inform scientific understanding of climate change impacts on UK coasts and seas, and monitoring water quality for the Water Framework Directive { ADDIN EN.CITE

<EndNote><Cite><Author>Devlin</Author><Year>2009</Year><RecNum>741</RecNum><DisplayText>(Devlin et al., 2009)</DisplayText><record><rec-number>741</rec-number><foreign-keys><key app="EN" db-id="ft5rwd95gpftsquad2a55exgesvz2tr0x2t" timestamp="1700666735">741</key></foreign-keys><ref-type name="Journal Article">17</ref-type><contributors><authors><author>Devlin, Michelle</author><author>Barry, Jon</author><author>Painting, Suzanne</author><author>Best, Mike</author></authors></contributors><titles><title>Extending the phytoplankton tool kit for the UK Water Framework Directive: indicators of phytoplankton community structure</title><secondary-title>Hydrobiologia</secondary-title></titles><periodical><full-title>Hydrobiologia</full-title></periodical><pages>151-168</pages><volume>633</volume><dates><year>2009</year></dates><isbn>0018-8158</isbn></urls></record></Cite></EndNote>.

An excellent UK example is the Continuous Plankton Recorder (CPR) survey, the most geographically extensive marine monitoring programme in the world, with over 7 million nautical miles of tows over 90+ years, routinely counting 650+ taxa and facilitating the production of over 1000 peer-reviewed scientific publications { ADDIN EN.CITE

<EndNote><Cite><Author>Richardson</Author><Year>2006</Year><RecNum>676</RecNum><DisplayText>(Richardson et al., 2006)</DisplayText><record><rec-number>676</rec-number><foreign-keys><key app="EN" db-id="ft5rwd95gpftsquad2a55exgesvz2tr0x2t" timestamp="1683025600">676</key></foreign-keys><ref-type name="Journal Article">17</ref-type><contributors><authors><author>Richardson, AJ</author><author>Walne, AW</author><author>John, AWG</author><author>Jonas, TD</author><author>Lindley, JA</author><author>Sims, DW</author><author>Stevens, D</author><author>Witt, M</author></authors></contributors><titles><title>Using continuous plankton recorder data</title><secondary-title>Progress in Oceanography</secondary-title></titles><periodical><full-title>Progress in oceanography</full-title></periodical><pages>27-74</pages><volume>68</volume><number>1</number><dates><year>2006</year></dates><isbn>0079-

6611</isbn><urls></urls></record></Cite></EndNote>}. Similarly, Plymouth Marine Laboratory's Western Channel Observatory L4 station is a biodiversity reference site with identification of over 500 plankton taxa alongside eDNA and benthic sampling. The sampling intensity (over 3000 net hauls since 1988) and number of variables sampled make it a testbed site for understanding how ecosystems operate, with over 350 scientific publications produced from this data { ADDIN EN.CITE <EndNote><Cite><Author>McEvoy</Author><Year>2023</Year><RecNum>726</RecNum><DisplayText>(McEvoy et al., 2023)</DisplayText><record><rec-number>726</rec-number><foreign-keys><key app="EN" db-id="ft5rwd95gpftsquad2a55exgesvz2trz0x2t" timestamp="1699544812">726</key></foreign-keys><ref-type name="Journal Article">17</ref-type><contributors><authors><author>McEvoy, Andrea J.</author><author>Atkinson, Angus</author><author>Airs, Ruth L.</author><author>Brittain, Rachel</author><author>Brown, Ian</author><author>Fileman, Elaine S.</author><author>Findlay, Helen S.</author><author>McNeill, Caroline L.</author><author>Ostle, Clare</author><author>Smyth, Tim J.</author><author>Sommerfield, Paul J.</author><author>Tait, Karen</author><author>Tarran, Glen A.</author><author>Thomas, Simon</author><author>Widdicombe, Claire</author><author>Woodward, M.</author><author>Beesley, Amanda</author><author>Conway, David V.P.</author><author>Fishwick, James</author><author>Haines, Hannah</author><author>Harris, Carolyn</author><author>Harris, Roger</author><author>Hélaouët, Pierre</author><author>Johns, David</author><author>Lindeque, Penelope K.</author><author>Mesher, Thomas</author><author>McQuatters-Gollop, Abigail</author><author>Nunes, Joana</author><author>Perry, Frances</author><author>Queiros, Ana M.</author><author>Rees, Andrew</author><author>Rühl, Saskia</author><author>Sims, David</author><author>Torres, Ricardo</author><author>Widdicombe, Stephen</author></authors></contributors><titles><title>The Western Channel Observatory: a century of physical, chemical and biological data compiled from pelagic and benthic habitats in the Western English Channel</title><secondary-title>Earth System Science Data</secondary-title></titles><periodical><full-title>Earth System Science Data</full-title></periodical><pages>1-42</pages><volume>2023</volume><dates><year>2023</year></dates><isbn>1866-3591</isbn><urls></urls></record></Cite></EndNote>}.</p>
</div>
<div data-bbox="111 531 889 730" data-label="Text">
<p>Novel technologies, including automated imaging and molecular methods, are being explored as cost-effective alternatives to traditional monitoring. Automated imaging uses high-speed photography and machine learning to identify, count and measure plankton in real-time or near real-time. Molecular methods involve collection and analysis of genetic material from plankton specimens or from the pelagic environment. While there are objectives to integrate novel technologies into routine monitoring, there are currently significant limitations to how data collected via novel technologies can be applied and a lack of long-term, consistent time-series to support assessments of pelagic habitats biodiversity. We should continue to explore these technologies, but it is essential that we do not attempt to do so at the expense of funding traditional long-term monitoring programmes, since they remain necessary to support biodiversity assessments. In addition, the taxonomic and ecological expertise to validate these novel methods needs to be maintained. To promote their integration into routine monitoring, novel technologies should be explored in parallel to traditional monitoring to better understand how the data they generate compares to the detailed taxonomic accounting of abundance generated via traditional methods.</p>
</div>
<div data-bbox="111 747 326 768" data-label="Section-Header">
<h2>Traditional monitoring</h2>
</div>
<div data-bbox="111 769 889 852" data-label="Text">
<p>Traditional plankton monitoring involves collecting samples directly from the ocean and subsequently preserving them so they can be later analysed via light microscopy. The CPR survey is a unique example, using a mechanical device to automatically collect and preserve samples while it is towed behind commercial ships and ferries as they travel their regular routes. The CPR is an exceptional case because it does not incur costs for research vessel time, unlike most methods, including those using novel methods.</p>
</div>
<div data-bbox="111 861 889 895" data-label="Text">
<p>Once on land, samples are stored and plankton entities are identified and counted in a laboratory under light microscope. Highly trained taxonomists identify and count organisms in each sample, following a consistent and</p>
</div>



documented method. Depending on the institute-specific procedure and density of organisms, sample processing time ranges from 3 hours to two days. This approach provides a high level of taxonomic detail, with semi-quantitative categories of taxa abundance. The traditional approach also allows a very rapid “sanity check” for unusual, new, and suspect taxa or results. This highly specialised method of plankton identification and enumeration is, understandably, resource intensive.

The high-quality and consistency of data collected in this manner facilitates comparisons over long time periods and between laboratories. The North-East Atlantic Marine Biological Analytical Quality Control (NMBAQC) Scheme provides a source of external Quality Assurance (QA) for laboratories engaged in the production of such marine biological data. Through the NMBAQC, laboratories engage in annual intercomparisons to ensure the phytoplankton and zooplankton data they generate are comparable to other laboratories and over time.



*The Plankton Imager, recently developed by Cefas and Plankton Analytics in the UK, is an imaging tool for sampling zooplankton without the need of human intervention. It consists of a high-speed camera that images all passing particles in a flow of pumped seawater. Images are identified in real-time and uploaded via satellite. This system requires no on-board expertise, harmful chemicals, or deployment of gear from the ship. It can collect data at a significantly finer spatial resolution than traditional methods. It also measures zooplankton, supporting carbon accounting. In its current state of development, its main capability is counting copepods.*

## Novel methods

Limited resources and budgets for monitoring have been a major driver for technological advances made for the development of more efficient cost-effective methods to gather plankton data { ADDIN EN.CITE

<EndNote><Cite><Author>Danovaro</Author><Year>2016</Year><RecNum>707</RecNum><DisplayText>(Danovaro et al., 2016)</DisplayText><record><rec-number>707</rec-number><foreign-keys><key app="EN" db-id="ft5rwd95gpftsquad2a55exgesvz2trz0x2t" timestamp="1699011517">707</key></foreign-keys><ref-type name="Journal Article">17</ref-type><contributors><authors><author>Danovaro, Roberto</author><author>Carugati, Laura</author><author>Berzano, Marco</author><author>Cahill, Abigail E</author><author>Carvalho, Susana</author><author>Chenuil, Anne</author><author>Corinaldesi, Cinzia</author><author>Cristina, Sonia</author><author>David, Romain</author><author>Dell&apos;Anno, Antonio</author></authors></contributors><titles><title>Implementing and innovating marine monitoring approaches for assessing marine environmental status</title><secondary-title>Frontiers in Marine

Science

Frontiers in Marine Science

213

3

2016

2296-7745

}. Currently there is an array of novel technologies available to identify and count plankton, including molecular

Yates et al., 2019

723

Journal Article

17

Yates, Matthew C

Fraser, Dylan J

Derry, Alison M

Meta-analysis supports further refinement of eDNA for monitoring aquatic species-specific abundance in nature

Environmental DNA

5-13

2019

2637-4943

}. and automated imaging methods

Pitois et al., 2018

709

Journal Article

17

Pitois, Sophie G

Tilbury, Julian

Bouch, Paul

Close, Hayden

Barnett, Samantha

Culverhouse, Phil F

Comparison of a cost-effective integrated plankton sampling and imaging instrument with traditional systems for mesozooplankton sampling in the Celtic Sea

Frontiers in Marine Science

5

2018

2296-7745

}. These techniques take advantage of the latest computing and genetic sequencing technology, allowing for a far greater throughput of samples than could ever be achieved manually.

Imaging instruments, combined with machine learning to automatically classify the collected images, have received a high level of interest, due to their ability to provide rapid and unbiased data that can be stored digitally and quickly made available for use

Giering et al., 2022

718

Journal Article

17

Giering, Sarah LC

Culverhouse, Phil F

Johns, David G

McQuatters-Gollop, Abigail

Pitois, Sophie G

Are plankton nets a thing of the past? An assessment of in situ imaging of zooplankton for large-scale ecosystem assessment and policy decision-making

Frontiers in Marine Science

986206

9

2022

2296-7745

}. Thus, they can overcome many of the limitations characteristic of traditional methods of collecting and analysing plankton samples. For example, the Plankton Imager (PI, Cefas) has been used aboard the RV Cefas Endeavour since 2016. While the ship is underway, seawater is pumped through a flow cell and a high-speed camera captures images of all passing particles. Each image is automatically classified as either copepod or non-copepod by a machine learning algorithm as well as recording size information in real-time. The system works continuously for the duration of a survey

}. Similarly, instruments such as the Imaging FlowCytobot (IFCB; McLane Labs) uses artificial intelligence to analyse phytoplankton images and can obtain quantitative results and accuracies comparable to

human analysts. However, these methods generally only classify individual organisms into coarse groups, missing the taxonomic detail that can be achieved by human analysts, and what is often needed to understand important changes in pelagic biodiversity.

	In-situ imaging	Traditional sampling	Molecular methods
<b>Cost</b>	High initial investment cost (labour + purchase), and moderate routine cost to run and maintain equipment.	Low initial cost, with low continuous costs of sample collection and processing	Medium initial investment in instruments, primer and high cost of analysis
<b>Accuracy and bias</b>	High accuracy of sample volume measurement through flow cell, low subsampling, reduced human bias since process is automated	Volume for net samples estimated or calculated from flowmeters, or fully quantifiable from bottle sampling. Some subsampling induced bias during processing.	Small volume collected and subsampling-induced biases, need to consider effect of dispersion since these methods detect molecules separate to organism
<b>Generation of quantitative data</b>	Fully quantitative for targeted groups (e.g. copepods) with high spatial resolution, but low taxonomic resolution	Fully quantitative with high taxonomic resolution, but generally low spatial resolution	Low spatial resolution, but very high taxonomic resolution (genetic), although false positive results are common (detection of species which do not occur in sampling location)
<b>Processing efficiency</b>	High computer power for processing in real or near real-time	Labour intense processing with delay between collection and data availability	High throughput sequencing, with delay between collection and data availability
<b>Storage and sharing</b>	Digital sample only - easily shared but requires large storage space	Physical samples which can be reanalysed and shared, preserved in chemicals and with large physical storage requirements, making long-term storage potentially expensive	Produces semi-quantitative data, difficult to translate to abundance or biomass
<b>Suitability for monitoring</b>	Primarily for research. New time-series being established in some cases.	Newly collected data add to long existing time-series.	Primarily for research
<b>Comparability among laboratories</b>	Typically developed independently using different instruments and classification algorithms	Simple and consistent methods which are easy to interpret, comparable between labs and over time	Typically using different instruments and targeting different genetic sequences, generating different results

Molecular methods, such as eDNA and metabarcoding use genetic information to study and identify the various species in an ecosystem. All molecular methods require the collection of physical samples prior to analysis in a laboratory. To derive abundance (and biomass) data from eDNA, promising links have been shown between eDNA and cell biovolumes { ADDIN EN.CITE <EndNote><Cite><Author>Song</Author><Year>2023</Year><RecNum>724</RecNum><DisplayText>(Song & Liang, 2023)</DisplayText><record><rec-number>724</rec-number><foreign-keys><key app="EN" db-id="ft5rwd95gpftsquad2a55exgesvz2trz0x2t" timestamp="1699536513">724</key></foreign-keys><ref-type name="Journal Article">17</ref-type><contributors><authors><author>Song, Jinxi</author><author>Liang,



Dong

**Community structure of zooplankton and its response to aquatic environmental changes based on eDNA metabarcoding** *Journal of Hydrology* 2023, 622:129692-1694. The main advantage of molecular tools is their ability to detect a broad array of taxa, including rare species, and in targeted tests provides the highest level of taxonomic detail. They can also be valuable for studying delicate organisms which can be easily damaged or destroyed by net sampling.

Govindarajan, Annette F., Francolini, Rene D., Jech, J Michael, Lavery, Andone C., Llopiz, Joel K., Wiebe, Peter H., Zhang, Weifeng. **Exploring the use of environmental DNA (eDNA) to detect animal taxa in the mesopelagic zone** *Frontiers in Ecology and Evolution* 2021, 9:574877-701X. However, since molecular methods rely on trace amounts of genetic material there is always a risk of generating false positives, or detecting taxa which are not actually present in a sample. Additional information on sex, life stage and individual conditions are impossible to gather using eDNA approaches. Currently, eDNA techniques are also limited in their ability to estimate abundance of taxa.

Yates, Matthew C., Fraser, Dylan J., Derry, Alison M. **Meta-analysis supports further refinement of eDNA for monitoring aquatic species-specific abundance in nature** *Environmental DNA* 2019, 1:4943-4943. and are best suited for the provision of presence/absence data.

An important difference between traditional and emerging plankton sampling approaches is the issue of scale.

Scott, James, Pitois, Sophie, Creach, Veronique, Malin, Gill, Culverhouse, Phil, Tilbury, Julian. **Resolution changes relationships: Optimizing sampling design using small scale zooplankton data** *Progress in Oceanography* 2023, 210:102946-6611. Traditional approaches frequently integrate temporally over a week to a month and spatially, to 10 nautical miles in the case of the CPR. For the latest autonomous imaging approaches the integration scales are much finer, typically resolved to meters vertically and horizontally, and over

minutes to hours for glider-, ship- or buoy-mounted instruments. This enables fundamentally different areas of science to be explored. For example, thermocline, diel cycle, or tidal cycle dynamics, impacts of extreme events such as storms or floodwater discharge or predator prey patch interactions. Obviously, however, these instruments and surveys applying them have not been running long enough to have the statistical power to resolve climate change scales. Moreover, since they span such different scales as well as sampling in a different manner, intercalibration with traditional monitoring is highly challenging. This fine level of detail is also typically not necessary for addressing policy needs.

## The risks associated with losing long-term monitoring time-series

While novel methods may appear more efficient than traditional methods, this may not necessarily be the case. There are significant hurdles involved with operationalising any novel method into practical routine monitoring. It will likely take decades before novel methods can demonstrate the consistent high-quality results of the traditional methods in use today. Currently, most imaging and molecular methods are research oriented or specialised towards a small suite of target species that are insufficient for assessing biodiversity status for policy.

Many of the UK's traditional plankton monitoring programmes have been ongoing since the 1990s and early 2000s, with the CPR commencing much earlier in 1931. Most programmes have employed the same equipment and methods since they were initiated to maintain the comparability of a time-series, despite the emergence of improved technology and methods. Due to the highly variable and patchy distribution of plankton in both space and time, long-term monitoring is mandatory for detecting important changes. Hydrological processes, such as the Atlantic Multidecadal Oscillation (AMO) and North Atlantic Oscillation (NAO), generate pressures on plankton communities lasting several decades. Changes in plankton communities resulting from human impacts can often be overshadowed by the variability attributed to these natural cyclical processes { ADDIN EN.CITE <EndNote><Cite><Author>Harris</Author><Year>2014</Year><RecNum>627</RecNum><DisplayText>(Harris et al., 2014)</DisplayText><record><rec-number>627</rec-number><foreign-keys><key app="EN" db-id="ft5rwd95gpftsquad2a55exgesvz2tr0x2t" timestamp="1666093820">627</key></foreign-keys><ref-type name="Journal Article">17</ref-type><contributors><authors><author>Harris, Victoria</author><author>Edwards, Martin</author><author>Olhede, Sofia</author></authors></contributors><titles><title>Multidecadal Atlantic climate variability and its impact on marine pelagic communities</title><secondary-title>Journal of Marine Systems</secondary-title></titles><periodical><full-title>Journal of Marine Systems</full-title></periodical><pages>55-69</pages><volume>133</volume><dates><year>2014</year></dates><isbn>0924-7963</isbn></urls></record></Cite></EndNote>}. Thus, long-term contiguous data is necessary for detecting the effects of climate change, shifting circulation patterns, and rising sea levels. We also need to continuously update long-term time-series to understand ecological change since we can only detect change by comparing current conditions to previous conditions. Therefore, the value of a time-series for addressing ecological questions is inextricably linked to its consistency and duration { ADDIN EN.CITE <EndNote><Cite><Author>Brander</Author><Year>2003</Year><RecNum>712</RecNum><DisplayText>(Brander & Drinkwater, 2003)</DisplayText><record><rec-number>712</rec-number><foreign-keys><key app="EN" db-id="ft5rwd95gpftsquad2a55exgesvz2tr0x2t" timestamp="1699021526">712</key></foreign-keys><ref-type name="Journal Article">17</ref-type><contributors><authors><author>Brander, Keith</author><author>Drinkwater, Ken</author></authors></contributors><titles><title>The relationship between scientific understanding and the length of time series: the CPR example</title><secondary-title>Content/Table des matières</secondary-title></titles><periodical><full-title>Content/Table des matières</full-title></periodical><pages>9</pages><dates><year>2003</year></dates></urls></record></Cite></EndNote >}.</p></div>

Research funding is increasingly allocated to exploring novel and innovative methods and technologies and it is becoming more difficult to obtain funding for routine monitoring or biodiversity assessments. In the 1980s the CPR survey was almost lost due to funding constraints. If the survey had not been saved, our current understanding of climate change and its impacts on marine biodiversity would be severely hampered { ADDIN EN.CITE <EndNote><Cite><Author>Reid</Author><Year>2003</Year><RecNum>526</RecNum><DisplayText>(Reid et al., 2003; Richardson et al., 2006)</DisplayText><record><rec-number>526</rec-number><foreign-keys><key app="EN" db-id="ft5rwd95gpftsquad2a55exgesvz2trz0x2t" timestamp="1653305788">526</key></foreign-keys><ref-type name="Journal Article">17</ref-type><contributors><authors><author>Reid, Philip C</author><author>Colebrook, JM</author><author>Matthews, JBL</author><author>Aiken, JCPR</author><author>Team, Continuous Plankton Recorder</author></authors></contributors><titles><title>The Continuous Plankton Recorder: concepts and history, from Plankton Indicator to undulating recorders</title><secondary-title>Progress in Oceanography</secondary-title></titles><periodical><full-title>Progress in oceanography</full-title></periodical><pages>117-173</pages><volume>58</volume><number>2-4</number><dates><year>2003</year></dates><isbn>0079-6611</isbn></urls></urls></record></Cite></EndNote>}. Continued cuts to monitoring budgets have resulted in the loss of several CPR routes, temporarily reinstated through the mNCEA programme, however, future funding remains uncertain. Multiple fixed-point stations are currently in a precarious status, including the Western Channel Observatory, subject to significant funding reductions. In Scotland, SEPA have discontinued plankton monitoring, the LPO has no dedicated funding to continue, and the Scottish Coastal Observatory has experienced significant reductions in taxonomic resource, impacting the volume of samples that can be analysed.

The erosion of monitoring programme funding has also contributed to reducing taxonomic capability within the UK research community and beyond. With the current inability to recruit junior taxonomists, often driven by a lack of resources, those who remain have little time to expand their skill set to focus on emerging species of concern. Critical skills are also lost when experienced taxonomists leave or retire { ADDIN EN.CITE <EndNote><Cite><Author>McQuatters-Gollop</Author><Year>2017</Year><RecNum>725</RecNum><Prefix>see</Prefix><DisplayText>(see McQuatters-Gollop et al., 2017)</DisplayText><record><rec-number>725</rec-number><foreign-keys><key app="EN" db-id="ft5rwd95gpftsquad2a55exgesvz2trz0x2t" timestamp="1699537845">725</key></foreign-keys><ref-type name="Journal Article">17</ref-type><contributors><authors><author>McQuatters-Gollop, Abigail</author><author>Johns, David G</author><author>Bresnan, Eileen</author><author>Skinner, Jennifer</author><author>Rombouts, Isabelle</author><author>Stern, Rowena</author><author>Aubert, Anais</author><author>Johansen, Marie</author><author>Bedford, Jacob</author><author>Knights, Antony</author></authors></contributors><titles><title>From microscope to management: the critical value of plankton taxonomy to marine policy and biodiversity conservation</title><secondary-title>Marine Policy</secondary-title></titles><periodical><full-title>Marine Policy</full-title></periodical><pages>1-10</pages><volume>83</volume><dates><year>2017</year></dates><isbn>0308-597X</isbn></urls></urls></record></Cite></EndNote>}. Research roles of the future will require taxonomic skills

to, for example, train Artificial Intelligence classification models, build genetic databases, and perform validation between traditional and novel methods. There needs to be a stronger recognition of the essential role that long-term time-series and taxonomic skills play in the development and incorporation of new technologies into ecological assessments. Without the traditional taxonomic skill set, these novel methods cannot be validated. Maintaining traditional long-term monitoring programmes will provide opportunities and incentives to promote training in taxonomy and will foster accelerated development of novel technology.

## How novel technologies can complement traditional monitoring

While imaging { ADDIN EN.CITE <EndNote><Cite><Author>Ostle</Author><Year>2023</Year><RecNum>720</RecNum><DisplayText>(Ostle & Hélaouët, 2023)</DisplayText><record><rec-number>720</rec-number><foreign-keys><key app="EN" db-id="ft5rwd95gpftsquad2a55exgesvz2tr0x2t" timestamp="1699283900">720</key></foreign-keys><ref-type name="Journal Article">17</ref-type><contributors><authors><author>Ostle, Clare</author><author>Hélaouët, Pierre</author></authors></contributors><titles><title>The Continuous Plankton Recorder as a platform for sensor development</title><secondary-title>PICES Press</secondary-title></titles><periodical><full-title>PICES Press</full-title></periodical><pages>64-65</pages><volume>31</volume><number>2</number><dates><year>2023</year></dates><isbn>1195-2512</isbn><urls></urls></record></Cite></EndNote>} and molecular { ADDIN EN.CITE <EndNote><Cite><Author>Suter</Author><Year>2021</Year><RecNum>719</RecNum><DisplayText>(Suter et al., 2021)</DisplayText><record><rec-number>719</rec-number><foreign-keys><key app="EN" db-id="ft5rwd95gpftsquad2a55exgesvz2tr0x2t" timestamp="1699283763">719</key></foreign-keys><ref-type name="Journal Article">17</ref-type><contributors><authors><author>Suter, Leonie</author><author>Polanowski, Andrea Maree</author><author>Clarke, Laurence John</author><author>Kitchener, John Andrew</author><author>Deagle, Bruce Emerson</author></authors></contributors><titles><title>Capturing open ocean biodiversity: comparing environmental DNA metabarcoding to the continuous plankton recorder</title><secondary-title>Molecular ecology</secondary-title></titles><periodical><full-title>Molecular ecology</full-title></periodical><pages>3140-3157</pages><volume>30</volume><number>13</number><dates><year>2021</year></dates><isbn>0962-1083</isbn><urls></urls></record></Cite></EndNote>} technologies lack the duration of use of traditional plankton monitoring time-series, the information they provide can be complementary to traditional approaches { ADDIN EN.CITE <EndNote><Cite><Author>Ratnarajah</Author><Year>2023</Year><RecNum>728</RecNum><DisplayText>(Ratnarajah et al., 2023)</DisplayText><record><rec-number>728</rec-number><foreign-keys><key app="EN" db-id="ft5rwd95gpftsquad2a55exgesvz2tr0x2t" timestamp="1700216954">728</key></foreign-keys><ref-type name="Journal Article">17</ref-type><contributors><authors><author>Ratnarajah, Lavenia</author><author>Abu-Alhaija, Rana</author><author>Atkinson, Angus</author><author>Batten, Sonia</author><author>Bax, Nicholas J</author><author>Bernard, Kim S</author><author>Canonico, Gabrielle</author><author>Cornils, Astrid</author><author>Everett, Jason D</author><author>Grigoratou, Maria</author></authors></contributors><titles><title>Monitoring and modelling marine zooplankton in a changing climate</title><secondary-title>Nature Communications</secondary-title></titles><periodical><full-title>Nature Communications</full-title></periodical><pages>564</pages><volume>14</volume><number>1</number><dates><year>2023</year></dates><isbn>2041-1723</isbn><urls></urls></record></Cite></EndNote>}. Due to their diverse range of sizes and patterns of distribution, no single method can effectively sample the full plankton community, leading researchers to select the most appropriate method to fulfil particular research or policy aims { ADDIN EN.CITE <EndNote><Cite><Author>Owens</Author><Year>2013</Year><RecNum>721</RecNum><DisplayText>(Owens et al., 2013; Skjoldal et al., 2013)</DisplayText><record><rec-number>721</rec-number><foreign-keys><key app="EN" db-id="ft5rwd95gpftsquad2a55exgesvz2tr0x2t" timestamp="1699284433">721</key></foreign-

keys<<ref-type name="Journal Article">17</ref-type><contributors><authors><author>Owens, NJP</author><author>Hosie, GW</author><author>Batten, SD</author><author>Edwards, M</author><author>Johns, DG</author><author>Beaugrand, Gregory</author></authors></contributors><titles><title>All plankton sampling systems underestimate abundance: response to "Continuous plankton recorder underestimates zooplankton abundance" by JW Dippner and M. Krause</title><secondary-title>Journal of Marine Systems</secondary-title></titles><periodical><full-title>Journal of Marine Systems</full-title></periodical><pages>240-242</pages><volume>128</volume><dates><year>2013</year></dates><isbn>0924-7963</isbn><urls></urls></record></Cite><Cite><Author>Skjoldal</Author><Year>2013</Year><RecNum>722</RecNum><record><rec-number>722</rec-number><foreign-keys><key app="EN" db-id="ft5rwd95gpftsquad2a55exgesvz2tr0x2t" timestamp="1699284482">722</key></foreign-keys><ref-type name="Journal Article">17</ref-type><contributors><authors><author>Skjoldal, Hein Rune</author><author>Wiebe, Peter H</author><author>Postel, Lutz</author><author>Knutsen, Tor</author><author>Kaartvedt, Stein</author><author>Sameoto, Douglas D</author></authors></contributors><titles><title>Intercomparison of zooplankton (net) sampling systems: Results from the ICES/GLOBEC sea-going workshop</title><secondary-title>Progress in oceanography</secondary-title></titles><periodical><full-title>Progress in oceanography</full-title></periodical><pages>1-42</pages><volume>108</volume><dates><year>2013</year></dates><isbn>0079-6611</isbn><urls></urls></record></Cite></EndNote>}. Novel technologies can provide efficient alternatives for addressing questions related to plankton distribution in space and time { ADDIN EN.CITE <EndNote><Cite><Author>Scott</Author><Year>2021</Year><RecNum>710</RecNum><DisplayText>(Scott et al., 2021)</DisplayText><record><rec-number>710</rec-number><foreign-keys><key app="EN" db-id="ft5rwd95gpftsquad2a55exgesvz2tr0x2t" timestamp="1699011879">710</key></foreign-keys><ref-type name="Journal Article">17</ref-type><contributors><authors><author>Scott, James</author><author>Pitois, Sophie</author><author>Close, Hayden</author><author>Almeida, Nevena</author><author>Culverhouse, Phil</author><author>Tilbury, Julian</author><author>Malin, Gill</author></authors></contributors><titles><title>In situ automated imaging, using the Plankton Imager, captures temporal variations in mesozooplankton using the Celtic Sea as a case study</title><secondary-title>Journal of Plankton Research</secondary-title></titles><periodical><full-title>Journal of plankton research</full-title></periodical><pages>300-313</pages><volume>43</volume><number>2</number><dates><year>2021</year></dates><isbn>0142-7873</isbn><urls></urls></record></Cite></EndNote>}, targeted detection of harmful algal bloom species { ADDIN EN.CITE <EndNote><Cite><Author>Medlin</Author><Year>2017</Year><RecNum>745</RecNum><DisplayText>(Medlin & Orozco, 2017)</DisplayText><record><rec-number>745</rec-number><foreign-keys><key app="EN" db-id="ft5rwd95gpftsquad2a55exgesvz2tr0x2t" timestamp="1702376246">745</key></foreign-keys><ref-type name="Journal Article">17</ref-type><contributors><authors><author>Medlin, Linda K</author><author>Orozco, Jahir</author></authors></contributors><titles><title>Molecular techniques for the detection of organisms in aquatic environments, with emphasis on harmful algal bloom species</title><secondary-title>Sensors</secondary-title></titles><periodical><full-title>Sensors</full-title></periodical><pages>1184</pages><volume>17</volume><number>5</number><dates><year>2017</year></dates><isbn>1424-8220</isbn><urls></urls></record></Cite></EndNote>}, and new migrants or alien species { ADDIN EN.CITE <EndNote><Cite><Author>Créach</Author><Year>2021</Year><RecNum>715</RecNum><DisplayText>(Créach et al., 2021)</DisplayText><record><rec-number>715</rec-number><foreign-keys><key app="EN" db-id="ft5rwd95gpftsquad2a55exgesvz2tr0x2t" timestamp="1699024536">715</key></foreign-keys><ref-type name="Journal Article">17</ref-type><contributors><authors><author>Créach, Véronique</author><author>Derveaux, Sabine</author><author>Owen, Katy R</author><author>Pitois, Sophie</author><author>Antajan, Elvire</author></authors></contributors><titles><title><style face="normal"

Use of environmental DNA in early detection of *Mnemiopsis leidyi* in UK coastal waters **Biological Invasions** 1-10 (2021) 1387-3547, however, traditional methods and associated skills remain critical to obtaining a detailed taxonomic accounting of the plankton community and validating novel methods. Most importantly, only traditional methods have decades of associated historical data required to detect long-term ecological changes.

We should be embracing novel technology, while also ensuring the continuity of traditional monitoring time-series. We cannot simply switch from traditional to novel methods since the continuity of long time-series is critical to supporting biodiversity assessments { ADDIN EN.CITE <EndNote><Cite><Author>Brander</Author><Year>2003</Year><RecNum>712</RecNum><DisplayText>(Brander & Drinkwater, 2003)</DisplayText><record><rec-number>712</rec-number><foreign-keys><key app="EN" db-id="ft5rwd95gpftsquad2a55exgesvz2trz0x2t" timestamp="1699021526">712</key></foreign-keys><ref-type name="Journal Article">17</ref-type><contributors><authors><author>Brander, Keith</author><author>Drinkwater, Ken</author></authors></contributors><titles><title>The relationship between scientific understanding and the length of time series: the CPR example</title><secondary-title>Content/Table des matières</secondary-title></titles><periodical><full-title>Content/Table des matières</full-title></periodical><pages>9</pages><dates><year>2003</year></dates><urls></urls></record></Cite></EndNote >}. As technology continues to improve, it is possible that traditional net sampling will become less important { ADDIN EN.CITE <EndNote><Cite><Author>Giering</Author><Year>2022</Year><RecNum>718</RecNum><DisplayText>(Giering et al., 2022)</DisplayText><record><rec-number>718</rec-number><foreign-keys><key app="EN" db-id="ft5rwd95gpftsquad2a55exgesvz2trz0x2t" timestamp="1699026081">718</key></foreign-keys><ref-type name="Journal Article">17</ref-type><contributors><authors><author>Giering, Sarah LC</author><author>Culverhouse, Phil F</author><author>Johns, David G</author><author>McQuatters-Gollop, Abigail</author><author>Pitois, Sophie G</author></authors></contributors><titles><title>Are plankton nets a thing of the past? An assessment of in situ imaging of zooplankton for large-scale ecosystem assessment and policy decision-making</title><secondary-title>Frontiers in Marine Science</secondary-title></titles><periodical><full-title>Frontiers in Marine Science</full-title></periodical><pages>986206</pages><volume>9</volume><dates><year>2022</year></dates><isbn>2296-7745</isbn><urls></urls></record></Cite></EndNote>}, however, until this occurs we must find ways to maintain taxonomic skills and apply traditional and novel methods in a complementary manner to facilitate ongoing scientific progress.

## References

{ ADDIN EN.REFLIST }