

1998

# The influence of nitrification in determining the supply, distribution and fate of nitrogen in grassland soils

Hatch , David John

<http://hdl.handle.net/10026.1/2305>

---

<http://dx.doi.org/10.24382/4052>

University of Plymouth

---

*All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.*

**THE INFLUENCE OF NITRIFICATION IN DETERMINING THE SUPPLY,  
DISTRIBUTION AND FATE OF NITROGEN IN GRASSLAND SOILS**

by

**DAVID JOHN HATCH**

A thesis submitted to the University of Plymouth in partial fulfillment for the degree of

**DOCTOR OF PHILOSOPHY**

Department of Agriculture and Food Studies,  
Seale-Hayne Faculty of Agriculture, Food and Land Use

in collaboration with  
The Institute of Grassland and Environmental Research



July 1998

## ABSTRACT

### The Influence of Nitrification in Determining the Supply, Distribution and Fate of Nitrogen in Grassland Soils

David John Hatch

The microbiology of nitrification has been extensively studied, but the ecology and environmental impact of the process has received less attention. The reason for this has more to do with the difficulties of conducting field experiments to examine the links with other processes which cause losses of nitrogen, than a failure to appreciate its importance in agricultural systems. This project was designed to overcome some of the limitations of existing field techniques to enable simultaneous measurements of nitrification and the major processes of N supply (mineralization) and N loss (denitrification and leaching) to be examined.

The study proceeded in three distinct phases: firstly, soils with contrasting N management histories were examined, using laboratory assays for *potential* activities. Clear differences which resulted from higher N inputs were established, with correspondingly higher nitrifying activities. For example, in a fertilized soil, ammonia-oxidizers produced 48.4 compared with 1.3 nM  $\text{NO}_2^- \text{g}^{-1} \text{soil h}^{-1}$  in an unfertilized soil. Potential nitrite-oxidizing rates were 93.4 and 62.5 nM  $\text{g}^{-1} \text{h}^{-1}$ , respectively. Assays of enzyme kinetics, therefore confirmed the higher nitrifying activity in the fertilized soil, but demonstrated a lower affinity of the enzyme for  $\text{NO}_2^-$  substrate, with  $K_m$  values of 436 and 310  $\mu\text{M NO}_2^- \text{-N}$ , respectively. Nitrifying rates in soils from grass-clover swards were intermediate between the fertilized and unfertilized soils.

Secondly, a new field incubation technique was developed and used to obtain *actual* rates by concurrent measurements of the major N cycling processes. A strong correlation was established between nitrification and denitrification ( $r^2 = 0.98$ ). The measurements were verified by comparison with other independent methods. Net rates of nitrification in the same soil type ranged from 0.55 - 1.17 kg N  $\text{ha}^{-1} \text{d}^{-1}$ , with the highest rates in the fertilized soil. Over 70% of the mineralized N was nitrified, of which 80% was subsequently lost (i.e. either denitrified or leached).

Thirdly, the practical implications of these findings were examined in greater detail using  $^{15}\text{N}$  labelling techniques which enabled process rates (net and gross) to be established in a model of the N cycle. When nitrification was inhibited, there were no significant differences between gross or net mineralization rates in the soils from the three swards, which indicated that N-immobilization could be directly influenced by the level of nitrifying activity in these soils.

The influence of nitrification in determining the pathways of N loss from grassland soils was quantified in this study. From a detailed investigation of the processes involved in N cycling, it was deduced that nitrification was also one of the major factors in determining the outcome of competition for inorganic N between plant and microbial biomasses.

## CONTENTS

Page No.

### CHAPTER 1 Introduction and literature review

1.1	The nitrogen cycle in grassland soils	1
1.2	Nitrification by autotrophic organisms	3
1.3	Heterotrophic nitrification	7
1.4	Other nitrification processes	10
1.5	Controls on nitrification	10
	1.5.1 Temperature	11
	1.5.2 Moisture and aeration	13
	1.5.3 pH effects	15
	1.5.4 Substrate	16
1.6	Coupling of processes	19

### CHAPTER 2 Development of a soil incubation system for the field measurement of nitrification

2.1	Project aims	22
2.2	Background to existing methodology	22
2.3	Description of equipment used in the novel incubation system	25
	2.3.1 Summary	28
2.4	Laboratory tests on component parts of the incubation system	28
	2.4.1 I-E Resins	28
	2.4.2 Tests on I-E bags	29
	2.4.3 Summary	34
2.5	Loading/flow rate effects on absorption by I-E bags	34
2.6	Inert porous blocks	38
	2.6.1 Oasis blocks	38
	2.6.2 Gypsum blocks	39
2.7	Acetylene inhibition of nitrification	41
	2.7.1 Formulations for controlled release capsules	44
	2.7.2 Laboratory tests on controlled release capsules	45
	2.7.3 Conclusions	48

### CHAPTER 3 Initial field testing of the soil incubation system and protocols for analysis of soil samples

3.1	Initial field tests of incubation system	49
	3.1.1 Temperature within the incubated soil core	49
3.2	Experimental sites	52
	3.2.1 Field measurements of mineralization/nitrification	53
	3.2.2 Results from initial field measurements	55
	3.2.3 Assessment of inorganic N in soils	57
	3.2.4 Summary	58
3.3	Protocols for analysing soil samples	59

## CONTENTS cont'd

	Page No.
3.3.1 <i>Effect of extraction shaking time</i>	60
3.3.2 <i>Effectiveness of KCl extraction on recovery of soil N</i>	61
3.4 Calculation of N cycling processes	62
 <b>CHAPTER 4 The effect of field management systems on the processes of decomposition and nitrification</b>	
4.1 Experimental layout	65
4.2 Mineralization measurements	67
4.2.1 <i>Field incubation</i>	67
4.2.2 <i>Anaerobic incubation</i>	68
4.2.3 <i>Chemical extraction</i>	68
4.2.4 <i>Results of mineralization measurements</i>	69
4.3 Microbial biomass	70
4.3.1 <i>Basal respiration</i>	70
4.3.2 <i>Substrate-induced respiration</i>	71
4.3.3 <i>SMB activity</i>	71
4.4 Laboratory tests of nitrifying activity	72
4.4.1 <i>Nitrification potential using a perfusion system</i>	72
4.4.2 <i>Results of measurements of nitrifying activity using perfusion tubes</i>	76
4.4.3 <i>Phosphate buffers</i>	77
4.5 Potential activity of nitrifiers in buffered soil slurries	77
4.5.1 <i>Potential activity of ammonia-oxidizers in soil slurries</i>	78
4.5.2 <i>Potential activity of nitrite-oxidizers in soil slurries</i>	80
4.5.3 <i>Results of ammonium-oxidation measurements using incubated soil slurries</i>	80
4.5.4 <i>Kinetics of nitrite-oxidation from incubated soil slurries</i>	82
4.6 Conclusions	86
 <b>CHAPTER 5 Concurrent measurements of net mineralization, nitrification, denitrification and leaching from field incubated soil cores</b>	
5.1 Experimental strategy	87
5.2 Investigation of microbial activity following spring application of N	88
5.2.1 <i>Net ammonification/nitrification/denitrification</i>	89
5.2.2 <i>Soil microbial biomass (SMB)</i>	89
5.2.3 <i>Results from the spring studies</i>	92
5.3 Investigations of N transformations in autumn/winter	97
5.3.1 <i>Methods of soil analyses</i>	101
5.3.2 <i>Results from the autumn studies</i>	103
5.3.3 <i>Conclusions from the autumn/winter studies</i>	109

## CONTENTS cont'd

Page No.

### CHAPTER 6 Development of a multi-point soil injector to measure gross and net rates of the major N cycling processes

6.1	Introduction: concepts and constraints	113
6.2	Equipment for $^{15}\text{N}$ enrichment of undisturbed soil	116
	6.2.1 <i>Injection of <math>^{15}\text{N}</math></i>	119
6.3	Experimental details	119
	6.3.1 <i>Testing uniformity of soil <math>^{15}\text{N}</math> injection</i>	123
6.4	Soil/plant processing	123
6.5	Diffusion of $^{15}\text{N}$ labelled solutions	123
	6.5.1 <i>Standards and controls</i>	126
6.6	Other measurements	127
6.7	Results and discussion	128
	6.7.1 <i>Summary</i>	138

### CHAPTER 7 Overview: project synthesis and conclusions

7.1	Introduction	139
7.2	Synthesis of the data	139
7.3	Re-appraisal of the main aims	142
	7.3.1 <i>Development of a new field incubation technique</i>	143
	7.3.2 <i>Linking in-situ nitrification measurements with other processes</i>	145
	7.3.3 <i>Examining the practical implications</i>	146
7.4	Sinks within the soil for mineral N	147
	7.4.1 <i>Microbial biomass</i>	149
	7.4.2 <i>Plant biomass</i>	149
7.5	Limitations/drawbacks identified	150
7.6	Future priorities for investigation	153
7.7	Options for control	154
7.8	Synopsis: acceptance/rejection of hypothesis	154
7.9	Conclusions	156

### CHAPTER 8 References 157

### APPENDICES

Appendix 1	172
Appendix 2	175
Appendix 3	176
Appendix 4	177

### LIST OF ABBREVIATIONS 180

## FIGURES

		Page No.
Fig. 1.1	The Nitrogen cycle	2
Fig. 1.2	Pathways for heterotrophic nitrification	8
Fig. 1.3	Acetylene and chlorate inhibitors	9
Fig. 1.4	Coupling links between processes	20
Fig. 2.1	Field incubation system	26
Fig. 2.2	Recovery of nitrate from I-E bags	30
Fig. 2.3	Effect of extractant pH	33
Fig. 2.4	Perfusion system	34
Fig. 2.5	Effect of loading/flow rate on recovery of nitrate by resin bags	37
Fig. 2.6	Effect of oasis on the SWC of soil cores	39
Fig. 2.7	Effect of gypsum blocks on the SWC of soil cores	40
Fig. 2.8	Diagram of gypsum blocks	41
Fig. 2.9	Standard curve for acetylene	45
Fig. 3.1	Temperature of incubated soil cores compared to surrounding soil	50
Fig. 3.2	Map showing experimental layout of treatments	54
Fig. 3.3	Net mineralization and nitrification	56
Fig. 3.4	Inorganic N in unfertilized and fertilized swards	58
Fig. 3.5	Diagram showing paired incubations	63
Fig. 4.1	Map showing experimental layout and subplots of treatments	66
Fig. 4.2	Soil perfusion tube with modified outlet	73
Fig. 4.3	Nitrification potential from soil perfusion columns	79
Fig. 4.4	Nitrification potential from incubated soil slurries	81
Fig. 4.5	Enzyme kinetics of nitrite-oxidizers in unfertilized soils	83
Fig. 4.6	Enzyme kinetics of nitrite-oxidizers in fertilized soils	84
Fig. 5.1	Net ammonification following N applications	93
Fig. 5.2	Denitrification following 2 <sup>nd</sup> spring application of N	94
Fig. 5.3	Correlation between nitrification/denitrification	95
Fig. 5.4	Mean daily rates of net mineralization/nitrification	105
Fig. 5.5	Mean daily rates of denitrification	106
Fig. 5.6	Correlations between nitrification and denitrification	107
Fig. 6.1	(a) CAD/CAM drawing of the injector manifold	118
	(b) Template patterns for TA, TB and TC	118
Fig. 6.2	Diagram of <sup>15</sup> N soil injector	120
Fig. 6.3	Schematic diagram of CN analyser linked to mass. spec.	125
Fig. 6.4	Gross mineralization/immobilization and net mineralization	137
Fig. 7.1	Diagram of N cycle processes and N pools	148

## TABLES

Table 1.1	The Family <i>Nitrobacteraceae</i>	6
Table 2.1	Acetylene (%) in soil atmosphere	47
Table 3.1	Effect of shaking time on extraction of inorganic N	60
Table 3.2	Recovery of N added to soil	62
Table 4.1	Some important properties of the soils	67
Table 4.2	Mineralization estimates by 3 methods	69
Table 4.3	Soil microbial biomass respiration rates	72
Table 4.4	Nitrifying activity measured with perfusion tubes	76
Table 5.1	<i>SMB</i> total C and total N	95
Table 5.2	Soil particle size distribution	98
Table 5.3	Percentage by volume of soil pore spaces	98
Table 5.4	Some properties of the freely and poorly drained soils	99
Table 5.5	Settling time before decantation of silt	102
Table 5.6	Comparison between the tube incubation and alternative methods	103
Table 6.1	Inorganic N contents	129
Table 6.2	Mean percentage recovery of $^{15}\text{N}$	131
Table 6.3	Gross rates of processes	132
Table 6.4	Nitrification and consumption of nitrate	133
Table 6.5	Net mineralization of nitrogen	134
Table 6.6	Soil microbial biomass N and C	135
Table 7.1	Inorganic N contents of the unlabelled N pools	152

## PLATES

Plate 2.1	Perfusion system and pump	35
Plate 6.1	$^{15}\text{N}$ soil injector	117

## ACKNOWLEDGEMENTS

I am pleased to record my gratitude to those who have helped and encouraged me during the course of this project. My particular thanks go to my Supervisors: Drs Steve Jarvis, (IGER), Rob Parkinson (University of Plymouth) and David Whitehead (University of Reading) for their guidance and the benefit of their considerable experience. I am grateful to Professor Roger Wilkins for providing me with the opportunity and facilities to undertake this work and to IGER for allowing me to undertake these studies on a part-time basis.

Others have helped in many ways: Jon Shaw, Phil Murray, Jane Hawkins and Robert Hatch with problems in computing and graphics; Stewart Ledgard, Ross Monaghan, Andrew Bristow, Daniel Murphy, Nick Ostle and John Stark with advice on  $^{15}\text{N}$  techniques; Roger Lovell, Pam Owen, Michael Davies and Liz Dixon with other technical aspects. I thank Andrew Rook for his statistical advice.

Special thanks should go to my wife Rosemary and son Robert who have supported and encouraged me throughout and to whom I now promise some quality time.

I should like this work to be dedicated to the memory of my father:

Frederick John Hatch (1920-1986)

## AUTHOR'S DECLARATION

At no time during the registration for the degree of Doctor of Philosophy has the author been registered for any other University award.

This study was carried out at the Institute of Grassland and Environmental Research, North Wyke, Okehampton, Devon: the work formed part of a MAFF commission with IGER.

In 1996, the author was awarded a Stapleton Fellowship and a British Grassland Society Bursary in order to undertake a period of study at Ruakura Agricultural Research Centre, Hamilton, New Zealand as part of a British Council-funded exchange.

Relevant scientific seminars and conferences were attended, at which the work has been presented. These are listed below.

**Publications:** (%Initiated; %Executed; %Wrote)  
Hatch, Jarvis and Parkinson 1996 (90; 100; 90)  
Lovell and Hatch 1997 (90; 60; 60)  
Hatch, Jarvis and Parkinson, 1998 (80; 100; 70)  
Ledgard, Jarvis and Hatch 1998 (0; 20; 5)\*  
Clough, Jarvis and Hatch 1998 (0; 30; 10)\*  
Bhogal, Hatch, Shepherd and Jarvis (in press)\*  
Patra, Jarvis and Hatch (in press) (10; 20; 30)\*  
(\*associated studies)

Presentations given, and Conferences attended:

**Posters:**


Seale Hayne Faculty of Agriculture & Land Use, September 1995  
British Soil Science Society Meeting, Belfast, September 1998

**Conferences:**

9th Nitrogen Workshop, Braunschweig, Germany, September 1996  
British Grassland Society, 5th Research Meeting, September 1997

**Presentations (seminars/talks):**

IGER, North Wyke, November 1995  
AgResearch, Invermay, New Zealand, December 1996  
British Grassland Society, Field Meeting at IGER, September 1997  
Seale Hayne Faculty of Agriculture & Land Use, June 1998  
University of Oslo, Norway, September 1998

Signed  .....

Date 31<sup>st</sup> July 1998 .....

## CHAPTER 1 Introduction and literature review

### 1.1 The Nitrogen cycle in grassland soils

Nitrogen (N) is a ubiquitous element which makes up 78.1% (by volume) of the atmosphere and is a component of all soils, comprising less than 0.1% of desert soils to over 2% of highly organic soils (Haynes, 1986). These vast sources of N, however, are unavailable to most forms of terrestrial life, which must rely on the activities of specialized microorganisms in order to obtain the N needed for the vital building blocks of life (e.g. amino acids, proteins). Only a few soil organisms (e.g. *Rhizobium*, *Azotobacter*, *Azospirillum*) are able to fix N from the atmosphere by conversion of dinitrogen (N<sub>2</sub>) to ammonium (NH<sub>4</sub><sup>+</sup>) using the nitrogenase-enzyme system (Sprent and Sprent, 1990).

In contrast, a host of heterotrophic organisms are able to assimilate organically-bound soil N, through the decomposition of soil organic matter (Dickinson and Pugh, 1974). The process of hydrolysis and catabolic breakdown of complex compounds into inorganic forms is termed mineralization and culminates in the release of NH<sub>4</sub><sup>+</sup> (ammonification). The energy made available to the decomposer organisms by this process enables the uptake of NH<sub>4</sub><sup>+</sup> for incorporation into cell components (Jansson and Persson, 1982). This anabolic process is termed immobilization and is concomitant with mineralization, although the net result of the opposing processes is, to some extent, dependent on the C:N ratio of the substrate; the critical C:N ratio for net mineralization is considered to be <25 - 30 (Harmsen and van Schreven, 1955), but is likely to vary between different ecosystems. Thus, net mineralization results in release of NH<sub>4</sub><sup>+</sup>, whereas net immobilization results in a decrease in available soil inorganic N.



This chapter describes the nitrification processes and the factors controlling the activity of the nitrifying organisms, which relate to the soil ecosystem. Our current understanding and areas where information is inadequate, or inconclusive are highlighted.


## 1.2 Nitrification by autotrophic organisms

Nitrification is generally defined as the biological oxidation of ammonia ( $\text{NH}_3$ ) to  $\text{NO}_3^-$ , with nitrite ( $\text{NO}_2^-$ ) as an intermediary product. Much of our present understanding of the nature of the process of nitrification stems from the early pioneering work of the Russian microbiologist, Sergei Nikolaevitch Winogradsky (1856-1946). In 1922, after an absence of nearly 35 years from scientific research, he returned to continue his investigations and reviewed the literature of the intervening years; he concluded that little new knowledge had been acquired to add to that of his earlier work. For an account of the life of Winogradsky, see Waksman (1946).

Almost half a century after Winogradsky's death, a wealth of literature has now been amassed, with contributions covering a wide range of factors concerned in the process of nitrification. These have been extensively reviewed under different environmental headings, including marine and estuarine waters (Kaplan, 1983; Ward, 1978), water and sewage treatment plants (Focht and Chang, 1975; Wanielista and Eckenfelder, 1978; Winkler, 1981; Painter, 1970) and soils ( Alexander, 1965; Focht and Verstraete, 1977; Belser, 1979; Schmidt, 1982; Haynes, 1986). However, the ecology of nitrifying bacteria and the factors controlling the growth and development of the organisms in their natural environments are still poorly defined, not least in agricultural soils, where edaphic factors (i.e. the soil chemical and physical properties which influence growing organisms) are usually disturbed

in some way when the measurements are made.

Winogradsky's major contribution (Winogradsky, 1890) was to identify two distinct genera of chemoautotrophic bacteria which are responsible for oxidizing nitrogen (N) in its most reduced form ( $\text{NH}_3$ ), through to the most oxidized form ( $\text{NO}_3^-$ ). He called the ammonia-oxidizing and nitrite-producing bacteria *Nitrosomonas*, and the nitrite-oxidizing and nitrate-producing bacteria were named *Nitrobacter*. The family *Nitrobacteraceae* (in the order *Pseudomonadales*) are slow growing, gram-negative aerobes with optimum activity at temperatures of between 30 and 37°C and a neutral to alkaline pH, with no growth at pH < 4.0 (Focht and Verstraete, 1977). They were shown to be true autotrophs, i.e. having no need of organic substrates because all their requirements for carbon are obtained from carbon dioxide ( $\text{CO}_2$ ), and their energy needs are met by the oxidation of ammonia and nitrite, as shown by the following equations:



Where  $\Delta F$  = Change in free energy

[from Russell, 1973]

The first oxidation stage involves the transfer of six electrons from the N in ammonia, with

a valency of 3-, through to the N in nitrite at a valency of 3+. The final step involves a further pair of electrons to produce nitrate, which has an oxidation state of 5+. The energy yield from these oxidizing reactions is very low, which makes the nitrifiers poor competitors against heterotrophic organisms. Fenchel and Blackburn (1979) calculated that *Nitrosomonas* generates only 1 to 4% and *Nitrobacter* from 3 to 10% of the cell material that might be expected from a heterotroph, from equivalent sources of available energy.

There are five genera of ammonia-oxidizers known to exist in soil: *Nitrosomonas*, *Nitrosolobus*, *Nitrosovibrio*, *Nitrospira* and *Nitrosococcus* and one nitrite-oxidizer, *Nitrobacter* (Belser, 1979). Since *Nitrosomonas* and *Nitrobacter* are the two most commonly isolated genera in terrestrial habitats, they have been assumed to be the most abundant. This is based on the principle of 'competitive exclusion', which proposes that a single species excludes all others from a generalized  $\text{NH}_4^+$  oxidizing niche (McLaren and Ardakani, 1972). This view has been challenged (Belser and Schmidt, 1978a), at least in the case of the ammonia-oxidizers, and may in reality only reflect their ability to out-compete other organisms in liquid culture. These authors found *Nitrosomonas* and *Nitrospira* to be more common than *Nitrosolobus*, but in another study on a Rothamsted soil, *Nitrosolobus* was found to predominate (MacDonald, 1979). Assessments of nitrifier populations by standard techniques of serial dilution and isolation, e.g. the Most Probable Number (MPN) method (Alexander, 1982), can give misleading results. This was demonstrated by Rennie and Schmidt (1974), who found counts of soil nitrifiers to be four orders of magnitude less by the MPN method, than determinations based on direct microscopic counting, using the fluorescent antibody (FA) stain (Bohlool and Schmidt, 1980). In another study, Johnson and Sieburth (1976) established the presence of

*Nitrosomonas* and *Nitrobacter* populations in a marine surface slime, by *in situ* electron microscopy, but were unable to culture the organisms by conventional methods. Belser and Schmidt ( 1978b) used FA staining and a modified isolation technique to show the existence of considerable serological diversity, particularly within the *Nitrospira* genus.

All the nitrifiers, with the exception of some strains of *Nitrobacter winogradsky*, are obligate chemolithotrophs (obtaining their energy from simple inorganic reactions). The currently recognized genera of chemolithotrophic nitrifiers is shown in Table 1.1.

**Table 1.1** The Family Nitrobacteraceae (From Belser, 1979)

Oxidation	Genus	Species	Habitat
Ammonium to nitrite	<i>Nitrosomonas</i>	<i>europaea</i>	Soil, water, sewage
	<i>Nitrosolobus</i>	<i>multiformis</i>	Soil
	<i>Nitrosovibrio</i>	<i>tenuis</i>	Soil
	<i>Nitrospira</i>	<i>briensis</i>	Soil
	<i>Nitrosococcus</i>	<i>nitrosus</i>	Soil
Nitrite to nitrate	<i>Nitrobacter</i>	<i>oceanus</i>	Marine
		<i>mobilis</i>	Marine
		<i>winogradsky</i>	Soil, water
Nitrite to nitrate	<i>Nitrospina</i>	<i>gracilis</i>	Marine
	<i>Nitrococcus</i>	<i>mobilis</i>	Marine


### 1.3 Heterotrophic nitrification

For many years after Winogradsky's discovery, nitrification was thought to be the sole preserve of autotrophic bacteria, and it was another 50 years or so, before the first evidence of N oxidation by heterotrophic organisms was presented (Nelson, 1929). It has now been shown that heterotrophic bacteria (e.g. *Arthrobacter*), as well as fungi (e.g. *Aspergillus*) and also some *Actinomyces*, are able to nitrify in pure culture under laboratory conditions (Focht and Verstraete, 1977) and are potential nitrifiers in soil (Kilham, 1986). The occurrence of heterotrophic nitrification, however, may be masked in natural systems since the process is not obligatory for the growth of the organisms. Higher yields of energy can be gained by heterotrophs from the oxidation of carbon (C) in organic substrates, than might be obtained from the small amounts of reduced N which appear to be oxidized during nitrification (Kilham, 1986). In view of this, the standard definition of nitrification as a strictly inorganic reaction is generally considered too restrictive, and an alternative definition of nitrification as:

**"The biological conversion of N in organic or inorganic compounds from a reduced to a more oxidized state",**

which was previously proposed by Alexander *et. al.* (1960), has now become more widely adopted. This definition is to be preferred as the precise biochemical pathway for heterotrophic nitrification is still uncertain and it may involve either an organic or inorganic pathway or a combination of both options, as shown in Fig. 1.2.


There are still doubts surrounding the role of heterotrophic nitrification in natural ecosystems (Schmidt, 1982) which may only become important in very acidic soils. For example, when  $\text{NH}_4^+$  was added to an acid soil, the basal rate of nitrification was not



**Figure 1.2** Possible pathways of heterotrophic nitrification (after Focht and Verstraete, 1977).

stimulated by the additional substrate (Weber and Gainey, 1962), suggesting the absence of any autotrophic component. In another study, evidence of nitrifying activity was detected even though nitrifiers could not be isolated; the addition of lime also failed to increase the measured rate of nitrification (Ishaque and Cornfield, 1972). The fact that nitrification occurs in situations where autotrophic organisms would not be expected to be active, or indeed in environments from where autotrophs cannot be isolated, provides only circumstantial evidence that the source of nitrification is heterotrophic (Focht and Verstraete, 1977). It is possible that unidentified species of autotrophs, which are adapted to such conditions, may yet exist. In any case, nitrification rates obtained from cultures grown axenically (i.e. in pure isolated cultures) are  $10^3$  to  $10^4$  times lower for heterotrophs than for autotrophs (Focht and Verstraete, 1977) and their contribution in natural systems may, therefore, be of little importance and only significant where autotrophs do not thrive.

The use of selective inhibitors may present the best opportunity for unravelling the relative contributions of different nitrifying processes in mixed populations. Acetylene ( $\text{C}_2\text{H}_2$ ) has



**Figure 1.3** Acetylene and Chlorate inhibition of autotrophic and heterotrophic nitrification (from Schimel *et al.*, 1984).

been shown to be an effective physiological block of autotrophic  $\text{NH}_4^+$  oxidation to  $\text{NO}_2^-$  (Berg *et al.*, 1982), as a non-competitive inhibitor of  $\text{NH}_3$  oxygenase which does not inhibit heterotrophic nitrification (Hynes and Knowles, 1982). Chlorate will inhibit the autotrophic oxidation of  $\text{NO}_2^-$  to  $\text{NO}_3^-$  (Hynes and Knowles, 1983) and both chlorate and  $\text{C}_2\text{H}_2$  have been used in a soil perfusion column to demonstrate the dominance of autotrophic nitrification in an agricultural soil maintained at pH values ranging from 4.5 to 7.5 (Kilham, 1987). In the same study, nitrification in an acid soil under coniferous plantation was shown to be predominantly heterotrophic. The blocking effect of these two inhibitors is illustrated in Fig. 1.3. More recently, the stable isotope  $^{15}\text{N}$  has been used in a modified application of the mean pool dilution technique to enable the separate pathways of autotrophic and heterotrophic nitrification to be evaluated (Barraclough and Puri, 1995). In this study only 8% of the observed nitrification in an acid woodland soil was attributed to heterotrophic activity, so that whilst biochemical blocks are effective in separating heterotrophic from autotrophic nitrification, their specificity is difficult to prove.

#### 1.4 Other nitrification processes

Two other nitrifying processes have been described in soils, but their significance is unclear. Methylophilic bacteria, capable of oxidizing methane and other C-containing compounds have been shown to also oxidize  $\text{NH}_4^+$  (Dalton, 1977) and a strictly non-microbial chemical oxidation can occur in soils exhibiting high manganese oxide content (Bartlett, 1981).

#### 1.5 Controls on nitrification

The term 'climax vegetation' is used to describe the end-point in the successional development of plant communities, which approach a steady-state when the vegetation has come into approximate equilibrium with the local conditions. For various successional stages leading up to climax vegetation, characteristically low  $\text{NO}_3^-$  levels were demonstrated for different soils, the lowest level being generally associated with the climax stage itself (Rice and Pancholy, 1972, 1973). These workers suggested that nitrification was inhibited under these conditions by increasing levels of tannins. However, Bohlool *et al.* (1977) could find no evidence of inhibition using pure cultures of nitrifiers with high concentrations of added tannins.

Similarly, unfertilized grassland soils generally have very little  $\text{NO}_3^-$  (Richardson, 1938) and small, but variable,  $\text{NH}_4^+$  contents. This observation led to speculation of allelopathy (the adverse influence that one living organism has on another by the secretion of a chemical inhibitor) associated with grass roots which were presumed to release substances into the rhizosphere that could inhibit nitrification (Theron, 1951). This hypothesis provided a plausible explanation of the controls which might govern low-input systems;

the conservation of the limited supply of nutrients under these circumstances, would be an essential stratagem to maintain the equilibrium. However, the difficulty in separating apparent allelopathic inhibition from other naturally occurring controls remains, and the latter constraints should first be eliminated before allelopathy can be invoked. Hence, Robinson (1963) found that low rates of nitrification in a grassland soil could be increased when soil pH and substrate in the form of urea were amended. Clark and Paul (1970) reviewed the literature on allelopathy in grassland, but the outcome remained inconclusive. More recently, Bremner and McCarty (1993) have looked again at the evidence, both for and against allelopathic interactions, and suggested that even where allelopathy is suspected, such effects are negligible.

An alternative explanation for low nitrification rates is that of nutrient-deprivation. For example, an inadequate phosphate supply was shown to limit  $\text{NO}_2^-$  oxidizing bacteria in a savanna grassland soil (Purchase, 1974a,b). The role of mycorrhizae may be of particular interest in this respect as they are known to be effective scavengers of phosphate and also preferentially take up N in the  $\text{NH}_4^+$  form. In this way mycorrhizae could act as direct competitors with nitrifiers for substrate and could, therefore exert a controlling influence in climax systems (Verstraete, 1981).

### 1.5.1 Temperature

The two physical factors which most affect nitrifier populations are temperature and moisture (Belser, 1979). As might be expected from the effects on microbes in general, a rise in temperature results in increased activity up to an optimum point. If the optimum temperature is exceeded, further increases cause a decline in activity until a stage is reached

from which the ultimate recovery to previous rates is permanently impaired. Evidence from the literature would suggest that autotrophic nitrification is more sensitive to high temperatures than the process of ammonification (Etinger-Tulczynska, 1969; Ishaque and Cornfield, 1974; Myers, 1975). Consequently, the supply of substrate ( $\text{NH}_4^+$ ) at elevated temperatures may exceed the oxidizing capacity of the nitrifiers and  $\text{NO}_2^-$  can accumulate (Mahandrappa, *et al.*, 1966). Nitrification activity is usually optimal between 25 and 35 °C (Justice and Smith, 1962) and was found to be almost completely inhibited at 40 °C by Keeney and Bremner (1967). However, these authors found that ammonification continued to be stimulated up to 40 °C, so that nearly all the  $\text{NH}_4^+$  produced remained in this form, whereas at lower temperatures, virtually all of the mineralized N was converted to  $\text{NO}_3^-$ . *Nitrosomonas* has a high activation energy equivalent to a  $Q_{10}$  of 3.0 ( $Q_{10}$  = the ratio of the rate of progress of a reaction, at a given temperature, to the rate at a temperature 10 °C lower), but *Nitrobacter* has a  $Q_{10}$  of only 1.7 (Belser, 1979). Increased temperature, therefore, stimulates  $\text{NH}_4^+$  oxidation to a greater extent than  $\text{NO}_2^-$  oxidation. Model-based predictions by Laudelout and co-workers suggested that when populations of these two nitrifiers were equal, high temperatures would lead to an accumulation of  $\text{NO}_2^-$ , and they were able to demonstrate that this could occur in practice (Laudelout *et al.*, 1974,75,76).

Low temperatures affect both ammonification and nitrification rates, but have a greater influence on the latter process (Tyler *et al.*, 1959). Fluctuating, low temperatures are more inhibitory than constant low temperatures, and it has been suggested that the effect on the microbes may be bacteriocidal (Campbell and Bierderbeck, 1972). Indigenous nitrifiers can alter their temperature optima in response to climatic variations (Mahendrappa *et al.*, 1966).

For example, in a tropical soil the optimum was found to be 35 °C with nitrification continuing up to 50 °C (Myers, 1975), whereas in a soil from Alberta (Canada) the optimum was only 20 °C and almost completely ceased at 30 °C (Mahli and McGill, 1982). Nitrification activity was still detected in Alberta soils even when the ground was frozen (Mahli and Nyborg, 1979), but nitrification is more generally inhibited until the soil has warmed to at least 4 or 5 °C (Anderson and Boswell, 1964).

### 1.5.2 Moisture and Aeration

The rate of diffusion of oxygen (O<sub>2</sub>) through air is ten thousand times faster than through water (Smith and Arah, 1990) so that changes in soil moisture content will directly affect the aerobicity of soils. The oxidation of NO<sub>2</sub><sup>-</sup> to NO<sub>3</sub><sup>-</sup> by *Nitrobacter* was shown by Aleem *et al.*, (1965) using <sup>18</sup>O, to involve molecular O<sub>2</sub> generated from water and not obtained directly from atmospheric O<sub>2</sub>. Although the soil liquid phase is the source of both the O<sub>2</sub> and HCO<sub>3</sub><sup>-</sup>, which are utilized in nitrification, Schmidt (1982) suggests that utilizable C is unlikely ever to be limiting to the autotrophic soil nitrifiers, but O<sub>2</sub> frequently is. For example, the solubility of O<sub>2</sub> would allow a concentration of 0.315 mM to develop in soil solution (at 15 °C) in equilibrium with air at atmospheric pressure, but Schmidt (1982) describes certain conditions that can promote O<sub>2</sub> depletion, *viz.* (i) high soil moisture content, which fills soil pores and restricts recharge of O<sub>2</sub> from the gaseous phase; (ii) high soil temperatures, which reduce the solubility of O<sub>2</sub> and increases O<sub>2</sub> demand by heterotrophic microorganisms; and (iii) readily-oxidizable organic matter, which also increases heterotrophic O<sub>2</sub> demand. A differential effect of limited O<sub>2</sub> supply on the two main nitrifier populations may be deduced from the work of Laudelout *et al.*, (1976) who found the  $K_m$  (the concentration of substrate at which the rate of reaction proceeds at half

the maximum rate) for  $O_2$  to be smaller for *Nitrosomonas* (0.016 mM) than for *Nitrobacter* (0.062 mM). Under conditions of low aeration and high temperature, their model predicted complete temporal separation of  $NH_4^+$  oxidation and  $NO_2^-$  oxidation; the  $NH_4^+$  oxidizers could out-compete the  $NO_2^-$  oxidizers for  $O_2$  and growth of the latter would be inhibited. This differs from the purely temperature-driven effect described above, where high temperatures alone did not inhibit the activity of *Nitrobacter*. In natural habitats, both groups of nitrifiers would compete poorly with heterotrophs, which have much lower  $K_m$  values (0.001 mM) for  $O_2$  (Focht and Verstraete, 1977).

When soil moisture tension approaches saturation point (0 kPa) nitrification ceases completely (Miller and Johnson, 1964; Malhi and McGill, 1982) or occurs only very slowly (Sabey, 1969). At moderately high moisture levels (1-10 kPa) nitrification is enhanced so long as aeration is adequate (Dommergues, 1966). At the "permanent wilting point" (-1500 kPa) nitrification can still occur (Miller and Johnson, 1964; Dubey, 1968; Sabey, 1969) and below this point nitrifiers are more inhibited than ammonifiers (Dommergues, 1966). Even severe desiccation probably only decreases the size of the population (Belser, 1979) and a proportion of it can remain viable in air-dried soil for as much as 14 years (Alexander, 1965). When soils were re-wetted after prolonged adverse conditions of high temperatures, high solute concentrations and desiccation during a four month dry period in a Moroccan coastal soil, activity quickly recovered following the onset of the first rainfall (Chiang *et al.*, 1972). Thus, where increasing soil moisture content tends to enhance nitrification, provided  $O_2$  is available, the nitrifying activity in dry soils will vary with soil texture along with those properties which govern osmotic potential (Russell, 1973).

### 1.5.3 pH effects

The sensitivity of nitrifiers to pH has been recognized for some time, but was probably the main reason for the general misconception of earlier workers that successful isolation and culture could only be achieved by the inclusion of solid particles (usually  $\text{CaCO}_3$ ) in the culture medium. The likely benefit of adding  $\text{CaCO}_3$  was to neutralize acid produced by the organisms, but when the pH was controlled the requirement for particulate matter was shown to be unnecessary, either for *Nitrobacter* (Goldberg and Gainey, 1955) or for *Nitrosomonas* (Engel and Alexander, 1958).

The pH optimum for the growth of nitrifiers is the alkaline side of neutral (Painter, 1970) which, in soils, encourages autotrophs to become the dominant species. Below pH 4.5 (e.g. in acid forest soils) autotrophs are inhibited (Sarawat, 1982) and nitrification is more likely to be heterotrophic. However, it is possible that some strains of autotrophic nitrifiers also exist which have adapted to acid conditions and in soils of pH 4 to 5, nitrification is often still detected (Weber and Gainey, 1962; Walker and Wickramasinghe, 1979; Federer, 1983). Conclusive proof of adaptation is hard to find since microsites may develop in acid soils which are at a higher pH than is indicated by bulk soil measurements. Liming can increase the numbers of autotrophs by several orders of magnitude (Soriano and Walker, 1973), but in soils with pH >7.5 the equilibrium between  $\text{NH}_4^+$  and free ammonia ( $\text{NH}_3$ ) moves in favour of the latter, which is particularly toxic to *Nitrobacter* (Morrill and Dawson, 1967).

At low soil pH the inhibition curve is a function of the equilibrium between  $\text{NO}_2^-$  and nitrous acid ( $\text{HNO}_2$ ); whilst acidity may be directly inhibitory, mobilization of aluminium is also implicated as a limiting factor (Brar and Giddens, 1968).

#### 1.5.4 Substrate

The major limitation to autotrophic nitrification is likely to be the availability of substrate itself (Alexander, 1965) and, therefore, the whole process is under the control of ammonification (mineralization). The inadequacy of this supply can be demonstrated in the soil by the addition of  $\text{NH}_4^+$ , which can greatly stimulate the growth of the nitrifier population (McLaren, 1971; Ardakani *et al.*, 1974). Thus, additional inputs of  $\text{NH}_4^+$  from other sources such as fertilizers, atmospheric deposition and hydrolysis of organic compounds in urine, will also exert an influence on nitrifying activity. The processes generally proceed in the order of the fastest, which is  $\text{NO}_2^-$  oxidation (a fact borne out by the negligible amounts of  $\text{NO}_2^-$  generally found in soil), followed by  $\text{NH}_4^+$  oxidation, with the slowest process being ammonification. Measurements of nitrification rates based entirely on  $\text{NO}_3^-$  production could be misleading; nitrifying activity can only proceed at the rate dictated by  $\text{NH}_4^+$  production.

The nitrifying organisms involved in the two stages of nitrification are sensitive to their own substrate, but are more sensitive to the substrate of the other organism (Painter, 1970). 'Substrate repression' by  $\text{NH}_4^+$  is in the range 400-800  $\mu\text{g N g}^{-1}$  dry soil (Haynes, 1986), but the inhibitory effect of high  $\text{NH}_4^+$  concentrations has also been attributed to the toxicity of  $\text{NH}_3$  (Broadbent *et al.*, 1957; Stojanovic and Alexander, 1958) or to the lowered pH caused by adding  $(\text{NH}_4)_2\text{SO}_4$  (Justice and Smith, 1962; Mahli and McGill, 1982). 'End product repression' can also occur, with  $\text{NO}_2^-$  concentrations ranging from 1400 to 4200  $\text{mg N l}^{-1}$  adversely affecting the growth of *Nitrosomonas* (Painter, 1970). Similarly, high concentrations of  $\text{NO}_3^-$  can inhibit  $\text{NO}_2^-$  oxidation by *Nitrobacter* (Boon and Laudelout, 1962).

*Nitrosomonas* and *Nitrobacter* oxidize about 35 and 100 atoms of N, respectively, for the fixation of one molecule of CO<sub>2</sub> (Alexander, 1965). About three times as much nitrogenous substrate, therefore, is required for the growth of NO<sub>2</sub><sup>-</sup> oxidizing bacteria than for NH<sub>4</sub><sup>+</sup> oxidizing bacteria. Cell yields calculated from this relationship and from cell counts suggest that from 1 to 4 x 10<sup>4</sup> cells of *Nitrobacter* μg<sup>-1</sup>N are produced from the oxidation of NO<sub>2</sub><sup>-</sup>, and about three times this number of cells of *Nitrosomonas* are produced from the oxidation of an equivalent amount of NH<sub>4</sub><sup>+</sup>-N (Focht and Verstraete, 1977). The 'generation time' (time taken for the population to double itself) for *Nitrobacter* has been reported to be as short as 7 hours and for *Nitrosomonas* to be about 11 hours by Alexander (1965) using culture solutions. In natural systems, Morrill and Dawson (1967) showed the number of *Nitrosomonas* bacteria did slightly exceed those of *Nitrobacter*, but were about the same when the soil was slightly acid (pH 6.4).

The affinity of the autotrophs for their substrates is indicated by  $K_m$  values (assessed in the temperature range of 20 to 30 °C) of between 1 and 10 mg N l<sup>-1</sup> and 5 and 9 mg N l<sup>-1</sup>, for NH<sub>4</sub><sup>+</sup> and NO<sub>2</sub><sup>-</sup> oxidation, respectively (Focht and Verstraete, 1977). Substrate concentrations in soil are generally likely to be equal to, or less than the  $K_m$  values and, therefore, oxidation rates will be dependent on the substrate concentration.

For this reason and those outlined above, it is clear that over-reliance on NO<sub>3</sub><sup>-</sup> levels as indicators of nitrifying activity is inadvisable; the determination of soil NO<sub>3</sub><sup>-</sup> in isolation has contributed little to an understanding of the dynamics of the process (Schmidt, 1982). Indeed, Jansson (1958) relegates the nitrate ion to a minor role in the N cycle in soils in favour of NH<sub>4</sub><sup>+</sup>, which is represented as the key form, undergoing continual consumption


and renewal by the microbial biomass. Competition for  $\text{NH}_4^+$  is placed in the order heterotrophs>nitrifiers>plants;  $\text{NO}_3^-$  then represents the inorganic N which is surplus to the requirements of the general soil biomass. Jackson *et al.* (1989) also found that plants were less successful competitors for  $\text{NH}_4^+$  than microbes. Besides microbial immobilization, other competing processes may be responsible for lowering the  $\text{NO}_3^-$  level in soil, e.g. plant uptake, denitrification and losses through leaching. Furthermore, even if the complication of  $\text{NO}_3^-$  uptake by plants is avoided, Russell (1973) lists four conditions which must be met before  $\text{NO}_3^-$  will accumulate in a fallow soil: (1) there must be an adequate supply of readily decomposable organic matter; (2) the soil must be kept free of weeds; (3) the soil must be kept moist, but not necessarily continuously; (4) there must not be too much rain, or  $\text{NO}_3^-$  will be removed by leaching. In attempting to accurately measure nitrification in the field, therefore, all of these naturally occurring factors need to be either eliminated, accounted for, or controlled.

Changes under carefully defined conditions could give important indicators which could help to characterize soil inorganic N status and behaviour. For example, Jarvis and Barraclough (1991) found that the ratio of  $\text{NO}_3^- : \text{NH}_4^+$  widened in a grassland soil with increasing N fertilizer rates. In the same study, when a common application rate of N was subsequently applied, differences in this ratio relating to the previous treatment remained, which suggested a change in nitrification had taken place. Increases in nitrifying potential, however, can be an undesirable development in stable ecosystems. The end-product of the process ( $\text{NO}_3^-$ ) is a more mobile ion than the  $\text{NH}_4^+$  ion (which is retained on cation exchange sites of soil particles,) and is, therefore, more vulnerable to loss.

## 1.6 Coupling of processes

Knowles (1978) has proposed that a close coupling may exist between the processes of nitrification, denitrification and possibly also N fixation, through intermediates which are common to different parts of the N cycle (Fig. 1.1). Thus,  $N_2$ ,  $NO_3^-$ ,  $NO_2^-$  and  $N_2O$  may all be represented as both substrates and products providing coupling links between the various processes, as illustrated in Fig.1.4. Knowles (1978), suggested that  $NO_3^-$  could provide a coupling between nitrification and denitrification in two ways. First, by 'successive coupling', whereby conditions conducive to nitrification (*viz.* dry/aerobic soils) produce an accumulation of  $NO_3^-$ . Conditions then develop which are more appropriate to denitrification (*viz.* wet/anaerobic soils), during which  $NO_3^-$  is reduced to  $N_2O$  and  $N_2$ . The second way is by 'simultaneous coupling', which proceeds from adjacent microsites. Nitrifying activity in an aerobic site near the surface of a saturated soil generates  $NO_3^-$ , which is transported downwards to an anaerobic microsite in the mobile water-phase, where the process of denitrification predominates. Wetting and drying cycles which can stimulate mineralization in soils (Birch, 1964) may, therefore, also promote further losses of N from the soil (Patrick and Wyatt, 1964).

The identification of  $N_2O$  as an intermediate product of  $NH_4^+$  oxidation (Yoshida and Alexander, 1970) can also be envisaged as providing a possible coupling between nitrification and denitrification (see Fig. 1.4) and in addition, has implications for the environment in respect of ozone destruction (Crutzen, 1970). Under strictly aerobic conditions,  $N_2O$  emission was demonstrated during nitrification of fertilizer N applied to several different soils (Bremner and Blackmer, 1978). However, the existence of isolated anaerobic microsites in otherwise aerobic soil conditions, can be brought about by the rapid



**Figure 1.4** Diagram of the processes of nitrification (upper), denitrification (lower) and  $\text{N}_2$  fixation (bottom), with possible coupling links shown by dashed lines (from Knowles, 1978).

decomposition of plant litter (with the associated consumption of  $\text{O}_2$ ), producing 'hotspots' of denitrification (Tiedje, 1987). This would tend to obscure the true source of  $\text{N}_2\text{O}$ . However, there is now increasing evidence to show that the processes of nitrification and denitrification can be combined in conditions which do not necessarily favour either process, and that the organisms responsible are less sensitive to  $\text{O}_2$  supply than was previously thought (Kuenen and Robertson, 1994). More recently, the relative contributions of the two processes to the flux of  $\text{N}_2\text{O}$  have been resolved using differential labelling of the respective N pools and examination of the  $^{15}\text{N}$ -distribution in the accumulated  $\text{N}_2\text{O}$  (Stevens *et al.*, 1997). The possibility of a further coupling between N fixation and processes which involve  $\text{N}_2$  and  $\text{N}_2\text{O}$  (Fig. 1.4) has been suggested, since nitrogenase can reduce both of the end-products of denitrification (Knowles, 1978).

A need for "simultaneous investigations of the interactions between nitrifying and reducing (particularly denitrifying) microorganisms", was identified as the dominant theme of the conference on: "Nitrification and Reduction of Nitrogen Oxides" (1976), by the chairman (Payne, 1978). The fact that his plea for more integrated ecological studies failed to produce an adequate response in succeeding years, is due more to the high degree of difficulty encountered in combining these studies and the complex physiology of the soil nitrifiers and their microhabitat, than to any diminution of interest in the topic. A similar request for studies to be undertaken in which natural conditions remained largely unaltered, was also made by Schmidt (1982). He stated that "microbiological data on the nitrifiers in relation to the processes as carried out in nature, would contribute considerably to an understanding of the basis for a particular nitrification rate, and to the reliability of models to predict rates at which nitrification takes place under various environmental conditions," but also cautioned that "a major difficulty in studying nitrification, especially in soil, is that the process interfaces with other components of the N cycle."

**It is the general aim of the present work to attempt to measure nitrification rates in soils under different grassland management regimes in the natural environment, with the contemporaneous assessment of N losses (through the processes of denitrification and leaching), to try to relate changes to possible coupling and/or stimulation of the processes involved.**

The specific aims of the project are stated at the start of the next chapter and the means by which they are to be achieved are also described and elaborated further in succeeding chapters.

## CHAPTER 2 Development of a soil incubation system for the field measurement of nitrification

### 2.1 Project aims

a) To **develop** a new technique for the field measurement of nitrification, with simultaneous estimates of losses of nitrate by denitrification and leaching. *The approach is firstly, to identify the limitations associated with existing methods and secondly, to attempt to overcome those aspects which most directly interfere with natural fluctuations in environmental conditions (eg. temperature, rainfall, evaporation).*

b) To **link** the *in situ* measurements of nitrification with other processes, to gain an understanding of the controlling factors determining the fate of nitrate. *Nitrification activity will be assessed in actual and potential terms and related to measured losses of nitrate by denitrification and leaching.*

c) To **examine** the practical implications of these findings under different grassland management regimes. *Many aspects of N cycling in grassland soils are investigated in isolation from related processes; the aim of this work is to combine different measurements in a microcosm of the field environment.*

### 2.2 Background to existing methodology

Since any limitation in the process of mineralization, which supplies substrate ( $\text{NH}_4^+$ ) to the process of nitrification, will necessarily determine the maximum rate at which the latter

proceeds (see Section 1.5.4), evaluation of nitrifying rates will be more meaningful if concomitant assessments of both processes can be made.

Recent advances in the use of the stable isotope  $^{15}\text{N}$  (Barraclough, 1991; Barraclough and Puri, 1995) have enabled measurements to be made of **gross** rates of several of the processes involved in N cycling in soil (including nitrification), but because of the expense and complexity of the analytical equipment and materials required, these methods are best suited to short-term evaluations that make direct comparisons between treatments. The  $^{15}\text{N}$  technique provides a powerful tool for the researcher, to add to the other methods available for field measurements, but is less well-suited to long-term seasonal measurements, which are required in order to construct balances of the flows of N in agricultural systems. The  $^{15}\text{N}$  technique will be discussed in detail in later sections and will be employed to assess **specific** rates of processes, under well-defined conditions, to supplement measurements of **potential** rates obtained from laboratory incubations.

In order, therefore, to obtain **net** measurements of the processes connected with nitrification, that may be related in seasonal terms to losses of N from the soil, an improved field incubation technique was developed to meet these requirements. The existing methods rely generally on an assessment of the incremental (net mineralization) or decremental (net immobilization) change in inorganic N ( $\text{NH}_4^+ + \text{NO}_3^-$ ). Soil samples taken at the start of an incubation are compared with a similar sample incubated in conditions that are intended to resemble closely the environmental fluctuations in the field. The methods adopted by workers to measure mineralization can roughly be divided into two soil sampling approaches: 1) disturbed soil and 2) undisturbed soil cores. Disturbed soil samples, either

crumbled by hand or sieved, may be incubated in the field by burying the sample in a polyethylene bag (Eno, 1960; Westerman and Crothers, 1980; Raison *et al.*, 1987). The advantage of this approach is that losses of  $\text{NO}_3^-$  by leaching are prevented, which might otherwise be overlooked in the evaluation. Losses of N by denitrification may, however, still result in an under estimate of mineralization and further inaccuracies can be caused by alterations to the soil structure. The use of relatively undisturbed soil samples in the form of intact soil cores will help to overcome some of these disadvantages (Cabrera and Kissel, 1988; Sierra, 1992), but losses can still occur as a result of denitrification. For this reason, Hatch *et al.* (1990, 1991) used acetylene ( $\text{C}_2\text{H}_2$ ) in the headspace of the incubating vessel to prevent nitrification and, therefore, avoid the production of  $\text{NO}_3^-$ , which might otherwise be denitrified to gaseous forms ( $\text{N}_2\text{O}$  and  $\text{N}_2$ ) which are more difficult to quantify. The assessment of net mineralization in this case, relied upon the measured accumulation of  $\text{NH}_4^+$  during an incubation of 14 days.

All these methods suffer from the shared problem of using samples that are effectively isolated, to varying extents, from the normal fluctuations in soil water due to evaporation and precipitation during the period of incubation. Gill *et al.* (1995) attempted to minimise these effects by using a shortened incubation period of 7 days, but they were unable to find any significant effect of soil water content (SWC) on mineralization rates, although the literature would suggest that wetting and drying of soil has a major influence on the process (Birch, 1964; Stanford and Epstein, 1974). In addition, the sides of the soil cores are exposed during incubation so that the increase in aeration may falsely elevate rates of mineralization.

### 2.3 Description of equipment used in the novel incubation system

The novel incubation system (Fig. 2.1) incorporates a PVC incubation tube (length 200 mm, i.d. 38 mm, o.d. 49 mm) which acts both as the soil coring tool (when fitted with a detachable cutting ring) and as a permanent sheath in which the soil core is held during incubation (Fig. 2.1a). A soil core is obtained by driving the PVC tube into the ground, using a club hammer and metal striking cap. The correct depth of core (150 mm length) is determined when the soil surface reaches a set of four holes (3 mm diameter) drilled around the tube (50 mm from the top and 150 mm from the base of the tube) which are equally spaced to provide outlets for surplus water to drain by lateral flow from the soil core surface. An O-ring ( neoprene, 5 mm diameter) is positioned immediately below the drainage holes, in a groove cut on the outside of the tube.

The PVC incubation tube containing the soil core is extracted from the ground by means of a larger metal soil corer (80 mm diameter i.d., 300 mm length). The metal corer is placed over the top of the protruding incubation tube and driven into the ground to a marked position equivalent to the depth of the soil core (Fig. 2.1b). In this way, the metal corer is used to extract the tube, together with an additional outer soil core, consisting of a cylindrical section which forms an annular ring (30 mm width) around the incubation tube (Fig. 2.1c). The soil from the outer core is carefully peeled away from the PVC tube and retained as a closely representative sample of the inner soil core sample. The inorganic N content of the outer core provides a baseline 'starting' value for subsequent comparisons with the field incubated soil core.




Figure 2.1 a - d. Field incubation system

An incubation vessel, consisting of a PVC tube (length 175 mm, 50 mm i.d., 60 mm o.d.), is driven into a suitably sized hole in the ground, so that the top of the tube is level with the soil surface, to provide a receptacle for the incubation tube (Fig. 2.1d). The inner edge of this vessel is machined to provide a groove into which the O-ring on the incubation tube locates to make a gas-tight connection between the two tubes. When the incubation tube is held in this position, it is suspended 25 mm above the soil at the bottom of the incubation vessel. Capillarity between the base of the soil core and the undisturbed soil layer is re-established and maintained through two circular layers of fine nylon (48 mm diam., 18.1 cm<sup>2</sup>, 355 µm mesh, Lockertex, Warrington, Cheshire) sewn together with nylon thread in the shape of a 'tea-bag', leaving a 6 mm gap in the perimeter. Using a small funnel, 4 g of wet ion-exchange (I-E) resin beads (20-50 mesh) are placed into the bag through this opening (see Section 2.4.1), which is then sealed using cyanoacrylate adhesive. The I-E bag is sandwiched between two filter paper discs (47 mm diameter, grade 40, Whatman Ltd.) to protect the resin and improve contact between surfaces. When the I-E bag is fully wetted it occupies a space 10 mm deep and the remaining space is filled with a gypsum block (47 mm diameter, 15 mm deep: see Section 2.6). The gypsum block makes good contact with the undisturbed soil layer at a depth of 175 mm from the soil surface and acts as a passive and inert medium for the movement of soil water between the incubated core and the surrounding soil.

Plant growth in the soil core is suppressed by means of a specially constructed device, referred to here as a *phytostat*, which prevents light from entering the incubation tube. It consists of a pair of PVC discs (37 mm diameter), centrally mounted on a metal shaft (6 mm diameter, 50 mm length) each of which has four holes (3 mm diameter) drilled through

the discs and spaced equally around the perimeter, 5 mm from the outer edge. The discs on the shaft are rotated so as to place them eccentrically to each other, with no vertical alignment between the holes. In this way, the *phytostat* permits water movement both in and out of the incubation tube, but light is excluded and plant growth prevented.

### 2.3.1 Summary

The novel incubation system has the following potential advantages over existing methods of field incubation:

- 1) Assessment of 'starting' values is based on soil which is closely associated with the soil core sample taken for incubation.
- 2) Natural wetting and drying cycles continue during incubation.
- 3) Aerobicity of incubated soil cores remains largely unaltered.
- 4) Losses of  $\text{NO}_3^-$  by leaching and denitrification can be assessed.

### 2.4 Laboratory tests on component parts of the incubation system

The incubation system comprises a means of assessing losses of  $\text{NO}_3^-$ -N by leaching using ion-exchange (I-E) resin bags and denitrification by a nitrification block method using acetylene ( $\text{C}_2\text{H}_2$ ). Each of these aspects will be examined in this section.

#### 2.4.1 I-E resins



Ion-exchange materials are insoluble matrices containing labile ions, which are capable of exchanging with other ions in the surrounding medium, but remain physically unchanged throughout. Ion-exchange is a stoichiometric process (the ion equivalents entering the resin are equal in charge to those leaving it), but resins exhibit a preference for certain ions (the

'selectivity coefficient'), so that the affinity of the resin for a particular ion needs to be considered carefully. Efficient ion-exchange is achieved when the ion which is to be absorbed has a considerably greater affinity for the resin than the ions already occupying the exchange sites.

Anion exchange resins derive their properties from the amino groups in the resin structure. The resin chosen to intercept  $\text{NO}_3^-$  in soil solutions was Dowex 1-x 8 (a strongly basic resin supplied in the  $\text{Cl}^-$  form by Fluka Chemicals, Gillingham, Dorset). The exchange capacity of the resin is  $1.33 \text{ mol dm}^{-3}$  (or  $1.33 \text{ meq. ml}^{-1}$  wet resin), operating over the full pH range and up to a maximum temperature of  $150^\circ\text{C}$ . The mesh size of the material used in the construction of the bags is constrained by the need to prevent leakage of resin whilst at the same time allowing for efficient desorption through the free movement of the extractant between the resin beads. Earlier tests on nylon materials showed that mesh sizes of  $<200 \mu\text{m}$  caused air bubbles to become entrapped, which prevented the bags from becoming fully immersed in the extractant, leading to incomplete desorption.

#### 2.4.2 Tests on I-E bags


The method of estimating  $\text{NO}_3^-$  leaching by I-E resin bags described by Lajtha (1988) was examined in greater detail to test its suitability to meet the requirements of the novel incubation system. Briefly, anion-exchange resin (Dowex 1-x 8) was converted to the bicarbonate form by three successive rinses in  $0.5 \text{ M NaHCO}_3$ . The I-E bags were rinsed



**Figure 2.2** Mean recoveries of  $\text{NO}_3^-$ -N from I-E resin bags using a) HCl ( $n = 3$ ) and b)  $\text{H}_2\text{SO}_4$  ( $n = 2$ ) extractants.

in deionised water and spun dry in a hand-operated centrifuge. For the purpose of the present study a domestic salad shaker (*Moulinex*) was found to be satisfactory. The I-E bags were each immersed in 200 ml of 3.6 mM NaNO<sub>3</sub> containing 50 mg l<sup>-1</sup> NO<sub>3</sub><sup>-</sup>-N, for 2 h with occasional stirring, then rinsed in deionised water and spun dry, as before. Samples of the original NO<sub>3</sub><sup>-</sup> solution, along with samples of the solution that had been exposed to the I-E bags, were kept for analysis. The I-E bags were extracted using three 50 ml rinses of 0.5 M HCl (Lajtha, 1988) and made to standard volume (200 ml) with deionised water. The recommended concentration of 0.5 M was extended to include 1.0, 1.5, 2.0 and 2.5 M HCl (3 reps. to each concentration) in the range of extractants. The efficiency of these extractants was expressed in terms of the mean recoveries of the known NO<sub>3</sub><sup>-</sup> loadings and calculated for each I-E bag from the difference between the extractant concentrations, analysed before and after exposure to the resin (Fig. 2.2a). Only poor recoveries of NO<sub>3</sub><sup>-</sup> were found at the lowest concentration of extractant 0.5 M (22.4%) and recoveries increased exponentially to the highest concentration of 2.5 M (61.9%), suggesting impractically high concentrations of HCl would be required for near complete recovery.


The choice of HCl seemed to be of doubtful value in meeting the requirements of the present study and was probably adopted by the previous author as a compromise to enable both NO<sub>3</sub><sup>-</sup> and PO<sub>4</sub><sup>3-</sup> to be extracted from soil solutions using the same resin. Also, replacement of the Cl<sup>-</sup> functional group with HCO<sub>3</sub><sup>-</sup> introduced an additional complicating step, with no apparent benefit in terms of affinity for NO<sub>3</sub><sup>-</sup> and the further disadvantage that the volume of the resin increased by some 20%. The order of ion affinities for anion exchange resins established by Kunin and Myers (1947) is:



It would appear, therefore, that  $\text{SO}_4^{2-}$  is the more effective ion to displace  $\text{NO}_3^-$ , which in turn, should readily displace  $\text{Cl}^-$  from the resin. Indeed,  $\text{Cl}^-$  was adopted as the functional group to be exchanged with  $\text{NO}_3^-$  by Wyland and Jackson (1993), who also found that recovery was enhanced when  $\text{SO}_4^{2-}$  was present in the solution and 2 M KCl was used as the extractant.

Recovery tests were repeated with  $\text{Cl}^-$ -charged resins, combining a longer exposure to  $\text{NO}_3^-$  solutions (24 h), so that additional information on the capacity of the I-E bags could be obtained. Extractant solutions containing  $\text{SO}_4^{2-}$  were considered in either the potassium or sodium form of the salt, but neither had the required solubility to achieve sufficiently high concentrations of the anion. Using duplicate I-E bags for each concentration to establish the trend, a range of  $\text{H}_2\text{SO}_4$  extractants was tried and included 0.5, 0.75, 1.0, 1.5 and 2.0 M solutions. Recovery of  $\text{NO}_3^-$  from the I-E bags was increased compared with HCl extractant and the maximum recovery improved to 75.7% (Fig. 2.2b).

The possibility that the increasing acidity of the extractant solutions was interfering with the  $\text{NO}_3^-$  analysis was checked by re-analysing the whole set of extractions. Each sample was divided into two sub-samples: one from each pair remained unchanged and the other was neutralized by titration with 0.375 M  $\text{Na}_2\text{CO}_3$ . No significant difference was found between the two sets of samples, which were closely correlated ( $r^2 = 0.864$ ), so that the increased recovery values can properly be attributed to the greater affinity of the resin for  $\text{SO}_4^{2-}$  over  $\text{NO}_3^-$  (Fig. 2.3).



**Figure 2.3** Effect of extractant pH on analytical values for nitrate. Data points are means ( $n = 2$ ).


A comparison was also made of the effect of increasing the extraction time to see if any improvement could be obtained in recovery. Four replicates each of the  $\text{NO}_3^-$  pre-loaded I-E bags were extracted in 1.5 M  $\text{H}_2\text{SO}_4$ , by either leaving to soak for 24 h with occasional stirring, or shaken vigorously for 1 h. Mean recoveries were 77.7% ( $\pm 5.16$ ) and 77.2% ( $\pm 1.40$ ) for 24 h and 1 h, respectively, and were not significantly different ( $P = 0.05$ ). The exchange capacity of the resin was found to be  $>2 \text{ mg NO}_3^- \text{-N g}^{-1}$  (wet resin), which meant that  $< 10\%$  of the theoretically available exchange sites were occupied. Average weekly winter rainfall (January, South West England) is about 30 mm and the average  $\text{NO}_3^- \text{-N}$  concentration in leached soil water (for grazed fields receiving  $200 \text{ kg N ha}^{-1} \text{ y}^{-1}$ ) is about  $10 \text{ mg l}^{-1}$  (Scholefield *et al.*, 1993). Each soil core (with a surface area of  $11.4 \text{ cm}^2$ ) could, therefore, potentially leach  $342 \mu\text{g N}$  during an incubation lasting one week, i.e. well within the exchange capacity of the I-E resin.

### 2.4.3 Summary

The I-E bags were used in the  $\text{Cl}^-$  form and the recommended regenerant was taken from a technical handbook (Ion-exchange resins, BDH Chemicals Ltd. Poole, England) specifying  $\text{HCl}$  (5% w/v): 350 ml for 100 ml moist resin (5.7 ml wet resin weighs 4 g). The method adopted to recover absorbed  $\text{NO}_3^-$  was to immerse the I-E bag in 50 ml of 1.5 M  $\text{H}_2\text{SO}_4$  extractant for 1 h with shaking, and then to rinse the bag with two further 50 ml batches of the extractant, finally making to standard volume (200 ml) with rinses of deionised water.

### 2.5 Loading / flowrate effects on absorption by I-E bags

The efficiency of absorption of  $\text{NO}_3^-$  by I-E bags was evaluated further using a perfusion system (see Fig. 2.4; Plate 2.1).




**Figure 2.4** Perfusion system



Plate 2.1 Perfusion tubes and pump

Twelve perspex leaching tubes (325 mm long, 50 mm i.d.) were supported in upright positions and each was packed to a depth of about 150 mm with 200 g acid washed, dry horticultural sand (screened to <1 mm). The lower ends of the tubes were stoppered with rubber bungs (size 49), centrally drilled to hold a length of glass tubing (70 mm long, 5 mm o.d.). The glass tubing was positioned so that the upper end was flush with the surface of the bung to provide a free-draining outlet from the perspex tube. The outlet was covered by a circular piece of fine nylon gauze (48 mm diameter, 140  $\mu\text{m}$  mesh) to retain the sand. Perfusate (100 ml) containing either 100, 50, 25 or 10  $\text{mg l}^{-1}$   $\text{NO}_3^-$ -N was used to wet the sand, with each concentration replicated three times. The perfusion tubes were allowed to drain freely overnight and then anion-exchange resin bags (in the  $\text{Cl}^-$  form, as previously described), sandwiched between two filter paper discs (47 mm diameter, grade 40, Whatman Ltd.), were placed beneath the sand and above the nylon mesh. Positioning of the I-E bags was achieved by briefly inverting the perfusion tube. A plunger was employed to retain the sand and care was taken to avoid displacing perfusate, so that the sand remained primed at the point where drainage would resume when more perfusate was added. In this way, precise timing of flowrates through the I-E bags was achieved and commenced from when eluent flowed from the outlets. Perfusate was supplied to the top of the tube from a peristaltic pump at either 50 or 100  $\text{ml 6 h}^{-1}$ . Previous tests on alternative means of controlling the rate of perfusion, consisted of connecting the bases of the perfusion tubes to a peristaltic pump with a pull-through flow system (Killham, 1987). However, this arrangement was less satisfactory for the purposes of the present tests, because 'fingering' (by-pass flow) occurred between the sand and the wall of the tubes and high concentrations of  $\text{NO}_3^-$  were detected in the eluents.

Good recoveries were sustained over the range of  $\text{NO}_3^-$  concentrations applied in perfusates at  $50 \text{ ml } 6 \text{ h}^{-1}$ , which simulated a rainfall event of *c.*  $7 \text{ mm h}^{-1}$  (Fig. 2.5). At the lowest  $\text{NO}_3^-$  loading ( $10 \text{ mg N l}^{-1} \times 50 \text{ ml } 6 \text{ h}^{-1}$ ), which represented an increase of nearly 50% in the (anticipated) N-loading in eluents from field incubated soil cores, over 90% was recovered in the I-E bags. At twice this flowrate, simulating extreme rainfall events of *c.*  $15 \text{ mm h}^{-1}$  (sustained for 6 h), the recoveries were less uniform and ranged from 77 - 94%. The low recovery of 77% from the highest loading ( $100 \text{ mg N l}^{-1} \times 100 \text{ ml}$ ) was to be expected, since it exceeded the known capacity of the resins by >20%. Performance of the I-E bags was, therefore, well within the necessary limits required to extract  $\text{NO}_3^-$  from soil solutions, but measurements of potential leaching losses from incubated soil cores will be adjusted using the established recovery factor of 90%.




**Figure 2.5** Effect of loading / flowrates on recovery of nitrate by resin bags (means of three replicates).

## 2.6 Inert porous blocks

The incubation system described in Section 2.3 specifies the requirement for a supporting structure which is both inert and porous to separate the I-E bags from contact with the soil layer beneath the incubated soil cores. The 'porous block' must ensure that the I-E bags are only exposed to  $\text{NO}_3^-$  in the mobile water phase emanating from the incubated soil core and not affected to any large extent by the undisturbed soil layer beneath the incubation vessel (Fig. 2.1d). In addition, water tension in the soil core had to be maintained and various materials were considered and tested for the necessary properties. The results from field tests on the two most promising materials are described below.

### 2.6.1 Oasis blocks


Cylindrical blocks (47 mm diameter, 15 mm deep) made from *Oasis* (an open cell expanded foam used by florists to provide water to plants under capillary tension), were tested for their suitability. Soil cores (4 replicates) were field incubated for 7 d periods using the new system lasting for a total of 35 d (total precipitation of 32.1 mm). Soil water contents (SWC) of the soil cores were compared with the surrounding soil (Fig. 2.6) using a Student t-test. Mean SWC of soil cores was significantly increased (by an average of 22%) compared with soil which was not in contact with the *Oasis* ( $P < 0.05$ ). The water-holding capacity of *Oasis* is almost  $1 \text{ ml cm}^{-3}$ , which amounts to a total of 28 ml water per block. The water content of the soil core at field capacity (Hallsworth series) was about 20 ml, representing a potential increase of 140% if all of the available water were taken up by the soil core. The likelihood of water-logging, particularly at the base of the soil core was high and the choice of *Oasis* was rejected.



**Figure 2.6** Effect of *Oasis* on the SWC of incubated soil cores under field conditions. Vertical bars = SEMs (n = 4)


### 2.6.2 Gypsum blocks

Calcium sulphate ( $\text{CaSO}_4$ ) occurs naturally and as the dihydrate,  $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$ , is known as gypsum. When gypsum is heated to  $97^\circ\text{C}$ , loss of water occurs to form the hemihydrate:




The product of the forward reaction is 'Plaster of Paris'. When water is added, the reaction is reversed and a solid mass of interlocking crystals of gypsum are formed, accompanied

by a slight expansion and generation of heat. Gypsum blocks (47 mm diameter, 15 mm deep) were individually moulded using 25 g of 'Plaster of Paris' in a 1:5 ratio with water (w/v) to achieve a thick, but still fluid mixture. The blocks were allowed to cure for 12 h and then a 6 mm hole was drilled through the centre to prevent any pools of water from forming on the top of the block. The water holding capacity ( $\pm$  SEM, 11 d.f.) of the gypsum block was 9.0 ml ( $\pm$  0.34) which was decreased slightly to 8.7 ml ( $\pm$  0.33) when a 30 cm tension was applied. Incubated soil cores (4 replicates) were compared over 7 d periods for a total of 28 d (total precipitation of 152.3 mm) and were not found to be significantly different from non-incubated soil (Fig. 2.7).



**Figure 2.7** Effect of gypsum blocks on SWC of incubated soil cores under field conditions. Vertical bars are SEMs ( $n = 4$ )

Data points have not been joined for the soil cores on either graph, because they represent



**Figure 2.8** Diagram of gypsum blocks

non-contiguous measurements from successive individual samples, whereas the data from the surrounding soil are repeated measurements from the same sample set. Gypsum blocks were incorporated into the field incubation system and were modified to accommodate controlled-release capsules (see following sections describing  $C_2H_2$  infusion of soil cores) by drilling four further holes around the periphery of the block (Fig. 2.8).


### 2.7 Acetylene inhibition of nitrification

The recognition of the inhibitory effect of  $C_2H_2$  on autotrophic nitrification (Walter *et al.*, 1979), also presented a means by which denitrifying activity could be suppressed through restricted substrate ( $NO_3^-$ ) supply. Warren (1988) employed  $C_2H_2$  in laboratory incubations to inhibit nitrification, thereby denying denitrification; the amount of accumulated  $NH_4^+$  was taken as an estimate of net mineralization. Ryden (1982) showed that only low concentrations of  $C_2H_2$  were needed to inhibit nitrification and mineralization was not affected. Hatch *et al.* (1990) further developed the method for field application by including 2%  $C_2H_2$  in the headspace of incubation vessels. Subsequent refinements (Gill

*et al.*, 1995) ensured that acetone (which is a contaminant of commercially supplied C<sub>2</sub>H<sub>2</sub>) was removed by two sequential water-traps to avoid introducing additional sources of energy. Another consideration is that degradation may occur in the soil, after a variable lag phase of up to 8 d, through the development of a C<sub>2</sub>H<sub>2</sub>-metabolizing population (Terry and Duxbury, 1985), which can then increase rapidly (Topp and Germon, 1986). Further difficulties associated with C<sub>2</sub>H<sub>2</sub>-inhibition have been identified by Smith and Arah (1990), who pointed out that soil types that would be expected to have the highest denitrifying capacity (e.g. heavy clay/anaerobic soils), often gave lower than expected rates, due to poor infusion of the gas through water-filled pores. Jackson *et al.* (1989) attempted to overcome this problem by infusing soil cores with C<sub>2</sub>H<sub>2</sub> dissolved in water, but this caused increases in soil water content.

Calcium carbide (CaC<sub>2</sub>) was used to generate acetylene gas for lighting purposes in the early days of the motor industry (1 kg CaC<sub>2</sub> yields 300 l of gas); a controlled release of gas was achieved by altering the rate of addition of water onto the carbide granules.

The reaction of carbide with water is:



Controlled release technology has been pioneered in the pharmaceutical industry using coatings of sugars or water soluble celluloses, but the difficulties of applying water soluble coatings to a water-activated ingredient (*viz.* CaC<sub>2</sub>), made them unsuitable for the present application. Previously, the majority of (medicinal) enteric coatings were applied as

solutions in organic solvents. However, because of the hazards associated with such solvents, e.g. cost, safeguards and disposal problems, these have been largely discarded in favour of aqueous solutions. Proprietary mixtures of water soluble celluloses (e.g. hydroxypropylmethylcellulose, HPMC) with water-insoluble celluloses (e.g. ethyl cellulose, EC) were tested in the present study. Combinations of mixtures (HPMC : EC) from 9:1 (soluble) to 5:5 (nearly insoluble) as 10% mixtures (w/v) in an organic solvent (50:50 methylene chloride/ethanol) were examined. Unfortunately, the difficulty of obtaining an even coating over irregular lumps of carbide, ultimately meant that this technique had to be rejected. An alternative approach of coating carbide with waxes was used by Banerjee and Mosier (1989) to achieve long-term persistence in soil. However, the method described by these authors was better suited to coating large particles of carbide and the encapsulating method gave variable and prolonged delays in the release of  $C_2H_2$  which did not meet the requirements of the present short-term incubations.

Gelatine capsules are employed to package medicinal formulations and in the past have also been coated to achieve delayed dissolution. Coatings such as beeswax, carnuba wax and shellac were used and, for some applications, the gelatine itself was modified chemically. Reactive side-chain groups in gelatine consist of amino, carboxyl and hydroxyl groups, of which the amino groups are the most amenable to chemical modification and the hydroxyl groups the least so. Reagents which are capable of reacting with two amino groups can cross-link within the gelatine matrix to give increased solution viscosity and a higher gel melting point, even to the extent that the gelatine becomes insoluble. Exposure to formaldehyde renders the gelatine capsule increasingly insoluble, but still porous to water. Pharmacological use of formaldehyde has now been discontinued, largely because of 'after-

hardening' of the capsules which can happen naturally through impurities in the gelatine (chelation of heavy metals), or at elevated temperatures ( $> 40^{\circ} \text{C}$ ).

To overcome some of the problems associated with contamination and persistence of  $\text{C}_2\text{H}_2$  in soil, the novel incubation method (a) restricts exposure of soil cores to  $< 8 \text{ d}$  and (b) replenishes metabolised  $\text{C}_2\text{H}_2$  by a controlled infusion of the gas using gelatine capsules containing  $\text{CaC}_2$ , as described below.

### 2.7.1 Formulations for controlled-release capsules

Hard-shell gelatine capsules (size 3; Farillon, Romford, UK) were pre-treated by exposure to formaldehyde vapour in a desiccator for 72 h. This process softens the capsule, but rigidity was re-established by heating at  $100^{\circ}\text{C}$  for 30 mins. During the heating stage, the original shape of the softened capsule was preserved by placing the open 'halves' on dress-making pins, mounted vertically in a balsa wood stand. Four types of controlled release capsules were prepared, each containing 100 mg  $\text{CaC}_2$  granules ( $< 2 \text{ mm}$ ) which were coated to delay the release of  $\text{C}_2\text{H}_2$  gas, as follows:

- Type 1: Uncoated, with rapid gas release from d0 - d1
- Type 2: Coated in paraffin wax, with a delayed gas release from d2 - d3
- Type 3: Coated in a mixture of paraffin wax / beeswax (50/50), with a delayed gas release from d4 - d5
- Type 4: Coated in beeswax, with a delayed gas release from d6 - d7

The gas release times are nominal and the success of the capsules is dependent upon the persistence of  $\text{C}_2\text{H}_2$  as a nitrification inhibitor and on a degree of overlap between release


times for the different types of capsules. The pre-drilled holes in the gypsum blocks (see Fig. 2.8) each holds one of the four types of capsules, which together generate a total of 30 ml of  $C_2H_2$ .

Waxes were refrigerated before use and fine shavings were obtained using a 'cheese grater'. The capsule base was filled to about 1/4 full with wax shavings and carbide granules (approx. 100 mg) were packed above the wax to the brim of the capsule. The capsule top was filled with more wax shavings and the two halves were pressed firmly together. Coating of the carbide granules was achieved by immersing the filled capsules in molten wax, which imparted the appropriate amount of heat to melt the wax inside the capsule, without deforming the gelatine. The granules were coated evenly by rolling the capsules around in the molten wax until the wax inside was thoroughly dispersed. The capsules were then removed and any wax adhering to the outside was wiped off with absorbent paper. When the capsules had cooled, the carbide granules were observed to have become encased in a fine matrix of wax and could be stored in a desiccator until needed. On exposure to water, the capsule softens, which distorts the wax matrix, so that the carbide granules become wetted progressively; the reaction releases  $C_2H_2$  which further disrupts the wax matrix and enables the ingress of more water.

### 2.7.2 Laboratory tests on controlled release capsules

Soil cores (150 mm deep) were obtained using incubation tubes which were then placed into incubation vessels (see Fig. 2.1) embedded in moist sand to simulate the supply of soil water under tension and kept in a fume hood at 20°C. The incubation tubes were modified to improve infusion of  $C_2H_2$  into the soil core: four upward sloping holes were drilled (3.5

mm diameter, 75 mm from soil surface) at angles of 30° and equally spaced around the tube to avoid by-pass flow of leachate from the sides of the tube. A gas sampling probe was constructed from stainless steel (s.s.) tube (1 mm diameter, 125 mm long) with a rubber suba seal (No. 13) fitted to one end and a microporous (s.s.) mesh (hydrophobic, but permeable to gas) attached to the other end. Gas sampling probes were inserted into each of four soil cores (50 mm from the surface) and a gypsum block was placed beneath each soil core at the bottom of the incubation vessel and in contact with the wet sand. Each gypsum block contained a different capsule (100 mg CaC<sub>2</sub>), representing one of the four types described above. After 1, 2, 4 and 5 d, samples of the soil atmosphere from around the probes were taken by withdrawing gas through the suba seal with a syringe. The first 0.5 ml was discarded and then a further 2 ml were withdrawn slowly and injected into the sample loop (0.5 ml) of an *Autofim*\* portable gas chromatograph, fitted with a flame ionisation detector.



**Figure 2.9** Standard curve for acetylene measured on an *Autofim* GC.

The samples were compared against a set of C<sub>2</sub>H<sub>2</sub> standards prepared from a cylinder of commercial-grade C<sub>2</sub>H<sub>2</sub>. The response of the instrument was linear (see Fig. 2.9) and the results are shown in Table 2.1.

**Table 2.1** Acetylene (%) in soil atmospheres at 50 mm from surface of soil cores, incubated at 20 °C with controlled release capsules.

Capsule type		Incubation period			
		1 day	2 days	4 days	5 days
Type 1	Uncoated	0.3	0.04	0.04	-
Type 2	Paraffin wax	0.16	0.04	0.04	0.03
Type 3	Paraffin/Beeswax	0.07	0.04	0.04	-
Type 4	Beeswax	-	-	-	0.04

\*The author gratefully acknowledges the loan of the *Autofim* instrument from Research Engineers Ltd., Orsman Road, London, N1 5RD for the duration of the test.

Concentrations of C<sub>2</sub>H<sub>2</sub> in the soil atmosphere at the end of a 5 d incubation (Table 2.1) were in the range found to be completely effective in preventing nitrification by other workers, viz. 0.01% (Berg *et al.*, 1982) to 0.1% (Walter *et al.*, 1979). The concentrations determined in the soil cores are likely to be elevated to some extent when the full complement of capsules is utilized, since the effect will be to reinforce the impact of individual capsules.

### 2.7.3 Conclusions

In this chapter, the successful development of a new field incubation technique was described. The individual components of the system were thoroughly tested under laboratory and field conditions and were shown to meet the required specifications. Ion-exchange resin had sufficient capacity and a high affinity for  $\text{NO}_3^-$  to extract the N leached from soil cores. Encapsulated  $\text{CaC}_2$  (coated in wax) produced a controlled-release of  $\text{C}_2\text{H}_2$  on exposure to moist soil. The gypsum blocks provided a porous medium to support the capsules, without significantly affecting soil water conditions during incubation. Conclusive evidence of the efficiency of the controlled release capsules in preventing nitrification in soil cores could only be obtained by measuring changes in  $\text{NO}_3^-$ -N under conditions of field incubation (see Section 5.2.3). A more comprehensive examination of the incubation system was undertaken in spring and autumn (1996), during which times the prevailing conditions are often conducive to high losses of  $\text{NO}_3^-$  through denitrification and leaching, respectively (see Chapter 5).

In the next chapter, the new incubation system was further tested under field conditions. Based on these initial data, the extent of sample variability was shown; improved methods were devised and protocols for analysis of soil samples were established. The development phase of the project culminated in a model to enable several N cycling processes to be measured simultaneously using sets of paired incubations ( $\pm\text{C}_2\text{H}_2$ ).



## **CHAPTER 3 Initial field testing of the soil incubation system and protocols for analysis of soil samples**

### **3.1 Initial field tests of incubation system**

Temperature and water content are major factors influencing both mineralization and nitrification rates (Miller and Johnson, 1964). Both variables were discussed in Section 1.5 in relation to nitrification, and operating limits for the incubation system were examined for soil water content (see Section 2.6). However, in previous studies using field measurements of net mineralization, temperature was found to exert the greater influence on rates (Hatch *et al.*, 1990; Gill *et al.*, 1995). This chapter describes the performance of the incubation system under field test conditions and the extent to which the temperature of the soil core was affected. The information obtained from monitoring changes in inorganic N levels during incubation and the protocols adopted to ensure consistent extractions of soil N are also detailed. These earlier tests did not, however, include C<sub>2</sub>H<sub>2</sub> infusion of soil cores for measurements of denitrification; elaboration of relevant calculations are presented at the end of this chapter in the form of a simple model and its application is to be discussed in subsequent chapters. The incubation system, incorporating the full complement of field measurements (net mineralization, nitrification, denitrification and leaching), enabled the major N cycling processes to be investigated.

#### **3.1.1 Temperature within the incubated soil core**

The incubation system was tested on two separate occasions over 7 d periods (21/6/94-27/6/94 and 7/7/94-13/7/94), by comparing the temperature within two incubated soil cores with that of the surrounding soil, using platinum resistance (2 k $\Omega$ ) thermometers



**Figure 3.1** Temperature of incubated soil cores compared with surrounding soil: measurements began at 0900h at the start of each 7d period in a) June 1994, b) July 1994.

(thermistors). A thermistor was inserted at 100 mm depth into each soil core and two others were placed in the surrounding soil (at the same depth): temperature was then recorded hourly. Data were stored on a Delta-T data logger (Delta-T Devices, Burwell, Cambs.) and the mean temperatures of soil cores and surrounding soil were compared using a paired sample t-test, for each 7 d period. The daily temperature patterns (Fig. 3.1a and b) between the incubated soil cores and the surrounding soil were highly correlated during both periods ( $r^2 = 0.976$  and  $0.993$ , respectively). A small, but significantly ( $P < 0.001$ ) lower mean temperature was recorded within the soil cores compared with the surrounding soil during both measurement periods (differences between means of  $0.28^\circ\text{C}$  and  $0.23^\circ\text{C}$ , respectively). It was evident that these small differences were associated chiefly with the periods following peaks in diurnal temperature fluctuations (around noon) and persisted until the lowest temperatures were reached (around 0800h). Thus, the small difference between an incubated soil core and the surrounding soil was most pronounced when the air temperature was declining. The soil core, insulated within the PVC tube, lost slightly more heat than the surrounding soil, which was buffered by having the greater mass. During the warming-up phase of the diurnal cycle, the temperatures measured within and outside the incubation tube were in close agreement.

It was concluded that enclosure of soil cores in the incubation system for periods lasting 7 d had no cumulative effect on the temperature within the soil core and only a small overall difference was recorded on a day-to-day basis. If the  $Q_{10}$  of the rate of nitrification is taken to be from 1.7-3.0 (see Section 1.5.1), then the process could be slowed by up to 8% during that part of the day when the temperature of the incubated soil core falls by *c.*  $0.25^\circ\text{C}$ .

### 3.2 Experimental sites

It has been suggested (Jarvis and Barraclough, 1991) that nitrifying activity may be increased by applications of N fertilizer. Based on this observation, the present investigation seeks to extend these findings and to relate them to N inputs and losses by direct measurement of the processes involved. In order to interpret the effects of systems of field management correctly, a long-term experiment with well established treatments was needed for the investigation. These requirements were provided by a field drainage experiment (Armstrong and Garwood, 1991; Scholefield *et al.*, 1993) at the Institute of Grassland and Environmental Research (IGER) site on Rowden Moor (OS Grid ref. SX 651995) at the North Wyke Research Station in Devon. At this site, hydrologically isolated field lysimeters, each of 1 ha long-term managed grassland, allow the comparison of the effects of field drainage with undrained soil. The site has been under permanent grassland management for over 40 years on a poorly drained clay loam soil, classified as a clayey, non calcareous pelosol of the Hallsworth series derived from the Culm Measures (a typical *pelo-stagnogley* soil (Findlay *et al.*, 1984; USDA = *typic haplaquept*; FAO = *dystic gleysol*). The combination of a high annual rainfall (on average, 1050 mm) and a soil with a high clay content (*c.* 36%), means that for much of the winter period the soils are waterlogged and without artificial drainage, a large proportion of the rainfall is removed in run off through surface or sub-surface lateral flow. Surface run off and drainage waters from each plot are separately channelled to outflow weirs to enable the measurement of flow rates and the collection of water samples for analysis.

Three experimental treatments within the experimental system were identified which are likely to provide contrasting features and may, over several years, have affected the

processes concerned in this investigation. The treatments selected for the study have all been installed with a system of field drainage (field drains 40 m apart and 85 cm deep with a porous backfill with temporary mole drains 2 m apart and 55 cm deep), that has been shown to increase mineralization rates compared with undrained soil, through improved aeration (Blanter, 1991). The selected treatments were continuously grazed by cattle during the growing season and since 1983 have either received 200 kg N ha<sup>-1</sup> y<sup>-1</sup> in 9 equal applications at 3 weekly intervals (*NF* treatment), or have remained unfertilized since 1983 (*ZN* treatment), or relied on inputs of fixed N from a mixed sward, which since 1988 contained clover (*GC* treatment). The treatments are represented by two replicate plots, except in the case of *ZN*, with only a single replicate. These treatments form part of a larger experiment (Tyson *et al.*, 1993) that has been used in previous studies on nutrient cycling (Scholefield *et al.*, 1993), but due to subsequent changes, now represent the only unchanged elements of the original experimental design of 13 years ago (Fig. 3.2).

### 3.2.1 Field measurements of mineralization / nitrification

A field incubation system based on 150 mm deep soil cores (as described in Section 2.3) was used to examine mineralization / nitrification in two of the long-term treatments, described above. A comparison was made between the *ZN* plot and one of the replicate plots from the fertilised treatment (*NF*, block B). Soil cores (4) were taken at random from within the grazed areas of these plots each week, starting on 7 October 1994 and lasting for a period of four weeks. Four replicate soil cores, enclosed in incubation tubes, were placed in the ground in incubation vessels and grouped together under cages (1.0 x 1.6 m) to protect them from grazing animals. The soil from each incubation tube, together with soil from the outer cores (representing 'start' values), were crumbled, thoroughly mixed and






Figure 3.2 Map showing experimental layout of treatments used in the present study (1994)

stones and pieces of undecomposed, identifiable plant material removed. Fifty grammes moist soil were shaken with 100 ml 1 M KCl for 1 h and the suspension was filtered using Whatman No. 1 filter paper, previously washed with 1 M KCl solution. Soil extracts were analysed colorimetrically for  $\text{NH}_4^+$  by the indophenol reaction (Krom, 1980) and for  $\text{NO}_3^-$  by hydrazine reduction (Kamphake *et al.*, 1967; Kempers and Luft, 1988) using a Skalar segmented flow analyser (Skalar, U.K. Ltd., Osbaldwick, York). Net mineralization rates were calculated as the difference in the inorganic N at the start and end of each incubation period. Net nitrification was calculated from the change in  $\text{NO}_3^-$  content from the 'start' values, and included  $\text{NO}_3^-$  absorbed from the soil core by the I-E bags. Soil inorganic N and soil water content (SWC) were determined from the non-incubated soil obtained at the start of each incubation period.

### 3.2.2 Results from initial field measurements

Mean rates of net mineralization (+) or immobilization (-) for the period (Fig.3.3a) were  $-0.24$  and  $0.05 \text{ kg N ha}^{-1} \text{ d}^{-1} \pm 0.166$  (*ZN* and *NF*, respectively), but were not significantly different. Only on one occasion (*viz.* 28 Oct. 1994) was there a significant difference in rates ( $P < 0.05$ ). The whole data set was statistically analyzed using *REML* (restricted maximum likelihood) by treating 'Weeks' (the sampling frequency) as a random effect and 'Treatments' (Units: *ZN* and *NF* from individual soil cores) as a fixed effect. The variance components for net mineralization were 0.0739 (Weeks) and 0.1445 (Units). It was evident that the variability in the data set due to spatial effects (soil coring) was almost twice that associated with temporal effects (sampling frequency). Thus, it was not possible to distinguish between environmental influences that might have been expected to affect net mineralization over the sampling period and the inherent variation within individual soil



**Figure 3.3** a) Net mineralization and b) nitrification in an unfertilized (ZN) and fertilized (NF) grazed sward.


cores. Such variation might well be expected in highly heterogeneous soils, which were also grazed and demonstrates the necessity for increased replication, particularly as the system depends upon destructive sampling.

Nitrification rates (Fig.3.3b) ranged from 0.01 to 0.366 and from 0.063 to 1.132 kg NO<sub>3</sub><sup>-</sup>-N ha<sup>-1</sup> d<sup>-1</sup> (*ZN* and *NF*, respectively), but on some occasions negative rates were recorded in both treatments, indicating that recovery of NO<sub>3</sub><sup>-</sup> was incomplete. Whilst care was taken to limit or remove competition for NO<sub>3</sub><sup>-</sup> (by suppressing plant uptake and monitoring for leached N), denitrification and microbial immobilization could still deplete the inorganic N pool in the soil core and lead to misinterpretations of rates of net nitrification. It was noticeable that the negative values for nitrification were coincident with an increase in SWC, following the return of the soil to field capacity and a tendency to waterlogging in the upper horizon (SWC >60%). Anaerobic conditions, conducive to denitrification, could have developed and may have led to unquantified losses of NO<sub>3</sub><sup>-</sup>-N. Mean rates of nitrification were 0.037 and 0.060 ± 0.113 kg N ha<sup>-1</sup> d<sup>-1</sup> (*ZN* and *NF*, respectively) with spatial variances being more than three times those which were temporally attributable (0.094 and 0.028 kg N ha<sup>-1</sup> d<sup>-1</sup>, respectively).

### 3.2.3 Assessment of inorganic N in soils

Amounts of inorganic N in the soil from the two treatments were assessed from non-incubated soil obtained at the start of each incubation period (Fig. 3.4). Ammonium : nitrate ratios were generally higher in *ZN* than *NF* (means of 2.3 : 1 and 1.8 : 1, respectively), except on 3 November, when *NF* had a higher ratio ( $P < 0.05$ ). The mean levels of NO<sub>3</sub><sup>-</sup>-N in *ZN* were lower than *NF* on all sampling dates (2.62 and 3.49 ± 0.438

kg N ha<sup>-1</sup>, respectively), but were only significantly different on 28 October ( $P < 0.05$ ), which suggests greater nitrifying activity in the latter. Levels of NH<sub>4</sub><sup>+</sup>-N were similar in both treatments with means of 6.15 (ZN) and 6.12 (NF) ± 0.773 kg N ha<sup>-1</sup>.



**Figure 3.4** Inorganic N in an unfertilized (ZN) and fertilized (NF) grazed sward.

### 3.2.4 Summary

The initial field tests provided indications of increased nitrifying activity in *NF* compared with *ZN*, but the absence of firmer evidence was due to the high degree of variation in the data. Nevertheless, three key areas were identified where improvements to the assessment of the processes of mineralization and nitrification could be made:

- i) Improve the recovery and consistency of soil inorganic N extractions
- ii) Increase the replication using more incubated soil cores / treatment
- iii) Assess losses of NO<sub>3</sub><sup>-</sup>-N attributable to denitrification

The first of these requirements are discussed in the following sections and the other two aspects are the subject of future chapters.

### 3.3 Protocols for analysing soil samples

The extraction of inorganic N from soil (*viz.*  $\text{NH}_4^+$ ,  $\text{NO}_2^-$  and  $\text{NO}_3^-$ ) is most commonly achieved by displacement of the ions with a high salt concentration (e.g. KCl or  $\text{K}_2\text{SO}_4$ ) in the extractant. Various concentrations of salts have been favoured, but the objective is not necessarily to achieve a complete recovery of the inorganic N from the soil, but rather to provide an index of the availability of soil N for plant uptake. Displacement of the negatively charged ions ( $\text{NO}_2^-$  and  $\text{NO}_3^-$ ), which are not strongly attracted to soil components, could be achieved simply by dispersion of the soil particles in water (using a blender or shaker). However, ammonium ( $\text{NH}_4^+$ ), which has a positively charged ion, is attracted to the negatively charged surfaces of clay particles and organic matter and displacement depends to some degree on the protection afforded by the soil structure. Displacement of  $\text{NH}_4^+$  is achieved by overwhelming the sites with an alternative cation. To achieve effective displacement, a high ratio of extractant ions :  $\text{NH}_4^+$  is required and substitution of cations on the soil-exchange sites is facilitated by using a high extractant volume : soil ratio. Ryden *et al.* (1987) employed 1 M KCl in a 2 : 1 (extractant : soil) ratio for inorganic soil analysis, but 2 M KCl has been more generally favoured (Bremner, 1965). Ratios of 2 : 1 (Jackson *et al.*, 1988), 3 : 1 (Cabrera and Kissel, 1988), 5 : 1 (Davidson *et al.*, 1991) and 10 : 1 (Donaldson and Henderson, 1990) have also been utilized, especially where the major portion of N is likely to be in the  $\text{NH}_4^+$  form: *viz.* during anaerobic incubation or with nitrification inhibitors (e.g. N-serve, Didin and  $\text{C}_2\text{H}_2$ ). To ensure uniformity, all further work in the present study employed 2 M KCl as the standard extractant and tests were carried out to establish suitable protocols for assessing ammonification and nitrification in field incubated soil cores.

### 3.3.1 Effect of extraction shaking time

Soil samples were collected from the long-term experimental site from an undrained treatment (receiving 200 kg N ha<sup>-1</sup> y<sup>-1</sup>) which was divided into areas that were permanently grazed by cattle throughout the grazing season and also from additional areas that were reserved for conservation of herbage for winter feed (Fig. 3.2). The latter received applications of slurry that were equivalent to the over-winter output of housed animals from the grazed areas. Soil was collected with a 37 mm diameter corer to a depth of 150 mm on 27 June 1995 from the cut and grazed areas to provide four representative samples from each (4 cores to each bulked field sample). Samples for extraction were prepared within 1 h of collection by crumbling the soil cores to < 6mm (stones and large roots discarded) and a 100 g sub-sample was shaken vigorously on a rotary shaker with 2 M KCl (200 ml). The standard shaking time of 1 h was extended to include 2 and 3 h. The samples were then filtered using Whatman No. 1 filter paper and stored in a freezer, until they were analysed for NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup> on an autoanalyser (as described earlier). Results were expressed on a dry soil basis, taking into account the volume of water present in the soil sample.

**Table 3.1** The effect of shaking time on the extraction of inorganic N with 2 M KCl ( $\mu\text{g N g}^{-1}$  dry soil). Mean values are shown with SEMs in parentheses ( $n = 4$ ).

Treatment	1 h	2 h	3 h
(undrained)			
200 N Cut	34.4 ( $\pm$ 7.38)	33.8 ( $\pm$ 6.62)	37.6 ( $\pm$ 7.38)
200 N Grazed	30.2 ( $\pm$ 5.53)	31.2 ( $\pm$ 5.17)	33.6 ( $\pm$ 5.42)

Data were analysed by *ANOVA* and are shown in Table 3.1. There were no significant differences between mean values obtained from the four replicates representing cut or grazed treatments, or with extended shaking times. A standard extraction involving vigorous shaking for 1 h was, therefore, adopted both for convenience and efficiency of operation.

### 3.3.2 Effectiveness of KCl extraction on recovery of added N to soil.

Soil was collected from the Rowden experimental site using a 25 mm diameter corer to a depth of 100 mm. A composited sample, representative of the Hallsworth soil series, was obtained by thoroughly mixing sieved (4 mm) soil from 120 cores from which the stones and large pieces of root material were discarded. Sufficient moist soil was weighed into 24 conical flasks so that each contained the equivalent of 25 g dry soil. Half of the flasks received 2 ml of 22.3 mM  $\text{NH}_4\text{NO}_3$  solution to provide 25  $\mu\text{g N g}^{-1}$  dry soil (both as  $\text{NH}_4^+$  and  $\text{NO}_3^-$ -N). The other 12 flasks received 2 ml demineralised water which raised the soil water content to approximately that of field capacity (c. 60%). The additions of liquid were made using a dropping pipette and were evenly distributed over the soil samples: flasks were then loosely covered and left to stand overnight at 20 °C. Inorganic N was extracted by adding 2 M KCl in ratios of either 2 : 1, 5 : 1 or 10 : 1 (KCl (vol) : dry soil wt.) to four replicate flasks for each ratio set. The flasks were shaken vigorously on a rotary shaker for 1 h and the suspensions were filtered through Whatman No. 1 paper (previously rinsed with 2 M KCl). Soil extracts were frozen for storage and then thawed out in a refrigerator, before being analysed for  $\text{NH}_4^+$  and  $\text{NO}_3^-$  (as described in Section 3.2.1).

Recovery of added N was over 80% with all three extractant : soil ratios tested, but was


more variable in the 2 : 1 set (Table 3.2). A possible explanation is that the clay particles in the Hallsworth soil (c. 36% clay content) were less well dispersed in the more viscous soil suspensions. The unrecovered N was probably immobilized during the equilibration period into organic forms which were not assessed. Incomplete recoveries were anticipated and consistent with the findings of other workers (Bristow *et al.*, 1987): the 'disappearance' of N would have been facilitated by the warm, moist and well-aerated conditions in the flasks stimulating N-immobilization by the soil microbial biomass. However, the higher recovery found with the 10 : 1 set was confounded by greater variability in  $\text{NH}_4^+$ -N contents, which was attributed to the increased dilution of the extracts and a consequent effect on analytical precision. On balance, therefore, the most consistent (i.e. least variable) recoveries were achieved with ratios of 5 : 1, particularly for  $\text{NH}_4^+$  which is the prime concern of investigations using incubated soil cores with nitrification inhibitors.

**Table 3.2** Mean % recovery (SEM) of  $\text{NH}_4\text{NO}_3$  added to soil and extracted with 2 M KCl ( $n = 4$ ).

KCl : soil ratio	$\text{NH}_4^+$ -N	$\text{NO}_3^-$ -N	Combined recovery
2 : 1	80 (20)	88 (20)	85 (20)
5 : 1	81 (2)	81 (5)	81 (3)
10 : 1	95 (8)	84 (2)	87 (4)

### 3.4 Calculations of N cycling processes using field incubation data

The difficulties of assessing N cycling processes in a field situation were illustrated in Section 3.2 and centred mainly on the incomplete recovery of  $\text{NO}_3^-$ -N from the incubated



**Figure 3.5** Diagram showing paired incubations with and without  $C_2H_2$

soil cores. Before an effective field evaluation of these processes could be made, it was also essential to quantify losses of N through denitrification. The problem was approached by arranging two soil incubations, either with or without  $C_2H_2$ , so that denitrification was denied in the first ( $+C_2H_2$ ) and allowed to proceed in the second ( $-C_2H_2$ ) as shown in Figure 3.5. Incubations are arranged to run in parallel and the processes of net mineralization, nitrification, leaching and denitrification can then be derived from the following model:

$$M = m + D + L, \text{ therefore, } D = M - (m + L)$$

and

$$N = [\text{Increase in NH}_4\text{-N (+C}_2\text{H}_2)] - [\text{Increase in NH}_4\text{-N (-C}_2\text{H}_2)]$$

(Assumes immobilization is same for incubations  $\pm$  C<sub>2</sub>H<sub>2</sub>)

where:

- $M$  = net mineralization
- $m$  = residual net mineralization (after losses through D and L)
- $D$  = denitrification (of original plus new sources of NO<sub>3</sub><sup>-</sup>)#
- $L$  = leaching potential (obtained from ion-exchange resin bags)\*
- $N$  = nitrification
- $d$  = denitrification confined to original nitrate at start of incubation
- $l$  = leaching measured with resin bags (check on effectiveness of C<sub>2</sub>H<sub>2</sub> )

**Notes:** Change in original NO<sub>3</sub><sup>-</sup>-N content (+C<sub>2</sub>H<sub>2</sub>) is net of any removal in denitrification ( $d$ ) and/or leaching ( $l$ ), where *de novo* NO<sub>3</sub><sup>-</sup>-N production is inhibited. For controls: # $D$  is not adjusted by  $d$  (to give **actual** N denitrified in measurement period); \* $L$  is adjusted by  $l$  (to give **potential** leaching of new NO<sub>3</sub><sup>-</sup> only), defined as (see Fig. 3.5):  $d$  = denitrification (confined to original NO<sub>3</sub><sup>-</sup> at start of incubation);  $l$  = leaching of original NO<sub>3</sub><sup>-</sup> (from I-E bags to check effectiveness of C<sub>2</sub>H<sub>2</sub>).

The development of the completed incubation system, with the facility for evaluating *in situ* denitrification losses by continual C<sub>2</sub>H<sub>2</sub> infusion (described in Chapter 2), required comprehensive field tests to assess the potential role of nitrification as a controlling influence on the processes of N cycling. The next chapter examines soils from defined management systems for evidence of any differences in nitrifying activity that might be related to differences in the observed losses of N.

## CHAPTER 4 The effect of field management systems on the processes of decomposition and nitrification

### 4.1 Experimental layout

During the 1995 grazing season, sub-plots were established within the main treatments previously described in Section 3.2. There were two major reasons for these modifications:

i) To establish the extent to which long-term management systems may have altered rates of mineralization and immobilization turnover (*MIT*) and nitrification; it was important that the findings were divorced from the possible masking effect of inputs of fertilizer and excretal returns during the current season.

ii) The original experimental design, based on large scale plots of 1 ha, provided only two field replicates each of treatments *NF* (200 kg N ha<sup>-1</sup> y<sup>-1</sup>) and *GC* (grass/clover) and only one replicate of *ZN* (unfertilized) treatment (Fig. 3.2), which placed a considerable constraint on statistical analyses. The field replicates were, therefore, divided into sub-plots (7m x 7m) within the main treatment which were individually fenced to exclude animals and were cut regularly to maintain the sward at a height comparable with the grazed areas. Fertilizer N was not applied in 1995.

There were four replicate sub-plots in the *ZN* treatment (no. 1-4) and two replicate sub-plots in each of the main field treatments: *NF* Block A (no. 5-6), *NF* Block B (no. 7-8); *GC* Block A (no.9-10) and *GC* Block B (no. 11-12) as shown in Figure 4.1. Some properties of the soil from these treatments which are considered to be of particular importance in the context of *MIT* studies are shown in Table 4.1.




Figure 4.1 Map showing experimental layout and subplots of treatments used in the present study (1995-8) Scale: 1cm=25m

## 4.2 Mineralization measurement

Three approaches to measuring mineralization were adopted: (i) net mineralization by a field incubation technique; (ii) potential mineralization by anaerobic incubation; (iii) an index of potential mineralization based on chemical extraction. The analyses were all performed on soil samples obtained in June 1995 from the sub-plots described earlier, which had not been grazed or fertilized since the grazing season of 1994.

**Table 4.1** Some important properties of the soils from the main field treatments.

Treatment (drained)	Total N (mgNg <sup>-1</sup> )	Organic C (mgCg <sup>-1</sup> )	C : N	pH		Bulk density	Water content at field capacity (%) (g g <sup>-1</sup> dry soil)
				water	CaCl <sub>2</sub>		
<i>ZN</i>	5.9	57.5	9.7	5.3	4.7	0.88	67
<i>GC</i>	5.1	53.6	10.5	5.5	4.7	0.99	63
<i>NF</i>	6.8	70.5	10.4	5.6	5.0	0.95	64

### 4.2.1 Field incubation

Using a 25 mm diameter soil corer, eight soil cores to a depth of 150 mm were extracted from each of the four replicate sub-plots from the main treatments. The cores were placed in 11 glass Kilner jars (8 per jar) and 2% C<sub>2</sub>H<sub>2</sub> was added to the headspace to prevent nitrification. The jars (12) were incubated in the field in suitably-sized holes for 6 d, using the technique reported by Hatch *et al.* (1990, 1991). At the same time, further soil cores were taken which were analysed for inorganic N, as described previously, but the ratio of KCl to soil (see Section 3.3.2) was increased to 5 : 1 (vol. : wt.). Net

mineralization, in the absence of nitrification, was determined from the change in  $\text{NH}_4^+\text{-N}$  between the incubated soil samples and those taken at the start.

#### 4.2.2 Anaerobic incubation

An anaerobic incubation method (Lober and Reeder, 1993), based on a modification of an earlier technique (Waring and Bremner, 1964), was used to obtain an estimate of potentially mineralizable N in a subset of soil samples obtained at the same time as field evaluations were carried out. Fresh soil was screened to pass a 2 mm sieve and the equivalent of 5 g dry soil was placed into 60 cm<sup>3</sup> polypropylene syringes (12) fitted with Luer-Lok taps (3 treats. x 4 reps.). Demineralized water (13 cm<sup>3</sup>) was added and entrapped air bubbles were dislodged using a hand-held vibrator. The plungers were then inserted into the syringes and all the remaining air was expelled through the tap, which was then closed. The syringes were clamped to a stand using saddle clips and placed in an incubator for 7 d at 35 °C. Following incubation, 37 cm<sup>3</sup> of 2.7 M KCl were added to each syringe, resulting in a final extractant concentration of 2 M KCl and a 10 : 1 ratio of KCl to soil (vol. : wt.). The plungers were then withdrawn sufficiently to introduce headspaces of approximately 10 cm<sup>3</sup> and the stand was firmly attached to a rotary shaker with the syringes held in a horizontal plane. The syringes were shaken vigorously for 1 h and the contents filtered through Whatman No. 1 paper (pre-rinsed in 2 M KCl). Filtrates were analysed for  $\text{NH}_4^+\text{-N}$  on an autoanalyser, as previously described. Mineralizable N was calculated as the difference between post and pre-incubation  $\text{NH}_4^+\text{-N}$  concentrations.

#### 4.2.3 Chemical extraction

Samples from the subset of soil cores, taken from the three field treatments (4 replicates),

were air-dried and ground to pass a 2 mm sieve. Mineralizable N was estimated using the method described by Whitehead (1981), *i.e.* 10 g dry soil were boiled for 1 h with 50 ml 1 M KCl and the extract was then filtered, made to standard volume and analysed for  $\text{NH}_4^+$ -N and  $\text{NO}_3^-$ -N as before.

**Table 4.2** Mineralization estimates by three methods. Mean values are shown ( $\mu\text{g N g}^{-1}$  dry soil) with SEMs in parentheses ( $n = 4$ ).

Treatment (drained)	Anaerobic incubation	Field incubation	Boiling KCl extraction
<i>ZN</i>	290 (12.8)	6.70 (0.540)	31.7 (2.68)
<i>GC</i>	242 (9.2)	6.16 (1.262)	30.4 (1.60)
<i>NF</i>	264 (40.0)	13.73 (0.639)*	33.2 (1.66)
Significance	n.s.	*sig. diff.	n.s.

\* $P < 0.05$

#### 4.2.4 Results of mineralization measurements

The three methods for assessing mineralization represent contrasting approaches to the problem of predicting the amounts of N which may be released from the soil O.M. fraction during a defined period of incubation. The lowest values were obtained from the field incubation measurements (Table 4.2) which provided an estimate of the **actual** rate relating to the period of incubation (6d) and were derived from readily degradable sources of organic N. Net mineralization in the fertilized treatment (*NF*) was > two-fold ( $P < 0.05$ ) that of the unfertilized treatments (*GC* and *ZN*). The chemical extraction (KCl) indices

were up to 5-fold higher than with field incubation values, but there were no significant differences between treatments. Previous studies (Hatch *et al.*, 1990, 1991; Gill *et al.*, 1995) have found this method to be relatively insensitive to the effects of field management systems, but probably more suited to identifying differences between soil types (Whitehead, 1981). Anaerobic incubation provided an estimate of **potential** seasonal mineralization with values which were nearly an order of magnitude larger than other methods for all three treatments; the highest value that was found with the *ZN* treatment was not significantly different from the other treatments (*NF* and *GC*).

#### 4.3 Microbial biomass

The Rowden experimental site was sampled again in autumn 1995, shortly after the soil had returned to field capacity (approximately 60% SWC, by weight) as indicated by the first appearance of drainage water flowing from the field drains. Ten soil cores (25 mm diameter) to a depth of 150 mm were taken from each treatment sub-plot (*ZN*: plots 1-4; *NF*: plots 5-8 and *GC*: plots 9-12, see Figure 4.1) The soil was crumbled to pass a 4 mm sieve and stones and identifiable pieces of root were rejected. Prepared soil samples were stored at 4° C in sealed polyethylene bags until required. Soil microbial biomass (*SMB*) activities were evaluated in terms of both basal respiration (as an indication of **intrinsic** activity) by the method described by West and Sparling (1986) and substrate-induced respiration (as an indication of **potential** activity) by the method described by West (1986).

##### 4.3.1 Basal respiration

Moist soil (4 field reps. per treatment), equivalent to 10 g dry weight, was placed into 72 ml serum bottles (24) and left for 30 minutes to equilibrate at 25 °C. The bottles were then

sealed with subseals (No. 21, Fisons, Loughborough, Leics.) and incubated for a further 18 h. Headspace gas samples were withdrawn (2 ml) and analysed for CO<sub>2</sub> using a gas chromatograph fitted with a thermal conductivity detector and a packed column (*Porapak Q*). Basal respiration was expressed as  $\mu\text{l CO}_2 \text{ g}^{-1}$  (dry soil) produced  $\text{h}^{-1}$ .

#### 4.3.2 Substrate-induced respiration

Following an 18 h incubation, the bottles (24) were vented and all received sufficient glucose solution (c. 20 ml) to provide 20 mg  $\text{g}^{-1}$  (dry soil), resulting in a 2 : 1 (vol./soil wt.) ratio. Half the bottles (12) also received cyclohexamide at 60 mg  $\text{g}^{-1}$  (dry soil) incorporated with the glucose solution to inhibit eukaryotic (fungal) respiration. The mixtures were swirled to disperse the soil and left to equilibrate for 30 minutes at 25 °C. The bottles were then stoppered ( $t_0$ ) and incubation continued with samples withdrawn from the headspace at 30 minutes ( $t_1$ ) and again at 150 minutes ( $t_2$ ) for CO<sub>2</sub> analysis, as above. The hourly rate of bacterial respiration was determined from the increment in the amounts of CO<sub>2</sub> produced between  $t_1$  and  $t_2$ , divided by 2, and expressed as  $\mu\text{l CO}_2 \text{ g}^{-1}$  dry soil  $\text{h}^{-1}$ .

#### 4.3.3 SMB activity

There were no significant differences between treatments in basal respiration rates, nor were there differences when substrate (glucose) was added, although all rates were increased by more than two-fold (Table 4.3). Estimates of biomass-C were obtained from the latter using the following formula of Ocio and Brookes (1990):

$$\mu\text{g biomass-C g}^{-1} \text{ (dry soil)} = 400.4 * (\text{cm}^3 \text{ CO}_2 \text{ 100 g soil}^{-1} \text{ h}^{-1}) + 3.7$$

The amounts of biomass-C were similar in all treatments (sampled in autumn 1995), but

low when compared with the results of Lovell *et al.* (1995), who found biomass-C was highest in the spring. The present estimates may reflect the timing of sampling, or some other effect related to storage (Ross *et al.*, 1980). Although not statistically different, substrate-induced respiration indicated that just over half the total respiration in the unfertilized treatments (*ZN* and *GC*) and slightly under half in the fertilized treatment (*NF*) could be attributed to bacteria when fungal respiration was inhibited.

**Table 4.3** Soil microbial biomass respiration rates. Mean values ( $\mu\text{l CO}_2 \text{ g}^{-1} \text{ h}^{-1}$ ) are shown with SEMs in parentheses ( $n = 4$ ). All values expressed on a dry soil wt. basis.

Treatment (drained)	Basal activity	Substrate- induced	Bacterial respiration	%Bacterial respiration	Biomass ( $\mu\text{g C g}^{-1}$ )
<i>ZN</i>	3.76 (0.224)	7.98 (0.705)	4.61 (0.239)	57.8	323
<i>GC</i>	3.80 (0.221)	7.86 (0.589)	4.37 (0.125)	55.5	318
<i>NF</i>	3.47 (0.046)	8.85 (0.623)	4.37 (0.589)	49.3	358


#### 4.4 Laboratory tests of nitrifying activity

Soils from the stored subset of samples, examined previously for *SMB* activity, were also used to estimate nitrification potentials by two techniques: (i) perfusion and (ii) soil slurries.

##### 4.4.1 Nitrification potential using a perfusion system

The perfusion tubes employed in testing the performance of resin bags (Section 2.5) were modified to enable eluent to be collected from soil columns. Preliminary tests using free-

draining columns filled with a soil and sand mixture to assess nitrifying activity (after Kilham, 1987) were inconclusive, as little or no  $\text{NO}_2^-$  (the primary product) could be detected in the eluents. The reason for this was unclear, but the high ratio of sand : soil that was necessary to ensure free drainage through the clay soils, resulted in flow characteristics which may not have allowed sufficient contact between the soil particles and the substrate in the perfusate. Ultimately, the method which was adopted had a modified outlet in the form of a manometer-tube overflow (Fig. 4.2) and used the equivalent of 50 g dry weight of fresh soil (4 mm sieved) mixed with 150 g of acid-washed, horticultural-grade silver sand (screened to  $< 1$  mm) and autoclaved before use. The outlet from the perfusion tubes, via a rubber bung, was covered by a disc of fine nylon mesh (140  $\mu\text{m}$ ) and then by a filter paper (Whatman No. 40, 47 mm diam.) and the soil and sand mixture was packed loosely into the tubes above the filter paper to a depth of about 150 mm. When fully wetted and freely drained, the mixture retained about 50 ml liquid.



**Figure 4.2** Soil perfusion tube with modified outlet

Soils from each of the main treatments (*NF*, *GC* and *ZN*: four field replicates of each) were allowed to equilibrate in 12 perfusion tubes at 20 °C for three days to minimise the effects of storage. With the manometer tube outlet placed level with the base of the column (the free-draining position), pumps were set to deliver phosphate buffer (see Section 4.4.2) at a rate of 50 ml 3 h<sup>-1</sup> (once in every 24 h) to adjust the soil gradually to pH 7.5. After this period, the soil / sand mixture had consolidated to a depth of about 100 mm. On the third day, the perfusate was changed to include 2 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> and 15 mM NaHClO<sub>3</sub> in buffer solution to evaluate ammonia-oxidizing activity and was supplied at 17 ml h<sup>-1</sup>. Chlorate is an effective inhibitor of NO<sub>2</sub><sup>-</sup> oxidation (see Berg and Rosswall, 1985) because the nitrifiers convert ClO<sub>3</sub><sup>-</sup> to chlorite, which is particularly toxic to *Nitrobacter*. At the same time, the manometer tube was raised to be level with the surface of the soil / sand mixture (the overflow position), so that the column was flooded before drainage occurred. After 3 h perfusion, drainage began and eluent was collected hourly over the following 6 h. Each set of eluents was immediately analysed for NO<sub>2</sub><sup>-</sup>, using a flow-injection analyser (FIA model 5020, Perstorp Analytical, Maidenhead, Berks.) to measure the absorbance ( $\lambda$  = 540 nm) of the purple azo-dye, produced by the reaction of NO<sub>2</sub><sup>-</sup> with sulphanilamide. The response of the instrument (Beer's Law) was non-linear (over concentration range 0 - 250  $\mu$ M, cell path length 10 mm) and the standard curve was best described ( $r^2 = 0.9999$ ) by a polynomial expression of the form :

$$y = ax^2 + bx + c$$

**Where**  $y$  = the absorbance reading for the samples.

The formula was re-arranged to solve  $x$  (the concentration of NO<sub>2</sub><sup>-</sup> in the samples) as follows:

$$x = \frac{-b \pm \sqrt{b^2 - 4a(c-y)}}{2a}$$

The process was repeated with a new set of soil samples in the perfusion tubes, equilibrated to pH 7.5 for three days as before. The perfusate was then replaced with a fresh buffer solution containing 2 mM Na<sub>2</sub>NO<sub>2</sub> (without chlorate) to evaluate nitrite-oxidizing activity. An inhibitor of autotrophic ammonia-oxidation was also included in the perfusate so that no new NO<sub>2</sub><sup>-</sup> was produced during the incubation. Other workers have previously used the inhibitor nitrapyrin (N-serve), but this has now been withdrawn from commercial use. An alternative product was tested, which is marketed under the trade name of *Didin-liquid* (Omex Agriculture Ltd., Kings Lynn, Norfolk, U.K.) and contains 40% w/w dicyandiamide (DCD) as the active ingredient. The inhibitor was added to the perfusate at a rate of 100 µg *Didin* g<sup>-1</sup> dry soil, or 0.004% (w/v) of the active ingredient (DCD) in the perfusate: comparable to the concentration of nitrapyrin adopted by Dusek (1995). Tests on DCD in conjunction with NH<sub>4</sub><sup>+</sup> substrate and chlorate in the perfusate, showed this rate to be completely effective in blocking nitrite production in soils from *ZN* and *GC* and only < 0.01% of the substrate was oxidized in *NF* soils. The small amount of NO<sub>2</sub><sup>-</sup> produced in this case could have occurred because of incomplete blocking due to some property of the soil structure, or it may suggest the presence of an heterotrophic nitrifying component in the soil from the fertilized treatment. Nitrifying activity was calculated as nMoles NO<sub>2</sub><sup>-</sup>-N g<sup>-1</sup> (dry soil) h<sup>-1</sup> by taking a linear regression through data points from four replicate columns of hourly rates of NO<sub>2</sub><sup>-</sup>, either produced (ammonia-oxidation), or consumed (nitrite-oxidation).

**Table 4.4** Nitrifying activity measured with perfusion tubes. Mean values are shown with SEMs in parentheses ( $n = 4$ ).

Treatment (drained)	Ammonium oxidation ( $\text{nM NO}_2^- \text{ g}^{-1} \text{ h}^{-1}$ )	Nitrite oxidation ( $\text{nM NO}_2^- \text{ g}^{-1} \text{ h}^{-1}$ )
<i>ZN</i>	0.96 (0.069) <sup>a</sup>	0.42 (0.015) <sup>a</sup>
<i>GC</i>	1.46 (0.089) <sup>b*</sup>	0.48 (0.024) <sup>b***</sup>
<i>NF</i>	1.77 (0.160) <sup>b***</sup>	0.59 (0.023) <sup>c***</sup>

Numbers in columns followed by different letters are sig. diff. (\*  $P < 0.05$ ; \*\*\*  $P < 0.001$ ).

#### 4.4.2 Results of measurements of nitrifying activity using perfusion tubes

The whole data set was analysed using a **grouped regression analysis** to establish whether the slopes of the time courses (i.e. the rates of nitrification per unit wt. dry soil, see Fig. 4.3a and b) were statistically different. There were significant differences in nitrifying activities between all three treatments for both  $\text{NH}_4^+$  and  $\text{NO}_2^-$  oxidizing activity, as shown in Table 4.4, except for the difference between rates for the *NF* and *GC* treatments ( $\text{NH}_4^+$ -oxidizers only), which was at a lower level of significance ( $P = 0.063$ ). The  $\text{NH}_4^+$ -oxidation rates ( $\text{nMoles g}^{-1} \text{ h}^{-1}$ ) were at the bottom end of the range found with New Zealand grassland soils (1.4 - 407) by Steele *et al.* (1980) and much lower than those from arable soils (100 - 800) surveyed by Kilham (1987) using perfusion techniques. After 4 h perfusion, the linearity in  $\text{NH}_4^+$  oxidation rates ceased in *NF* and *GC* (Fig. 4.3a), possibly due to poisoning of the enzyme complex. It is possible that the chlorate (on conversion to chlorite) could have adversely affected  $\text{NH}_4^+$  oxidation (Groffman, 1987), although it has been found to be much more selective against  $\text{NO}_2^-$ -oxidizers and not to affect  $\text{NH}_4^+$

oxidizers (Belser and Mays, 1980). In this respect, it is possible that perfusion may reinforce any adverse inhibitory effects by continual supply/replacement of the inhibitor, which could have a more pronounced effect than the single dosage employed with incubated soil slurries (see Section 4.5).

#### 4.4.3 Phosphate buffers

Although many workers recommend that measurements of potential nitrification are performed on soils buffered to near neutral, or slightly alkaline pH (Belser and Mays, 1980; Kilham, 1987; Dusek, 1995), none gives detailed instructions on the formulation of suitable buffer solutions. An empirical approach was adopted to establish the combination of 2.5 mM phosphate solutions required to produce a pH 7.5 phosphate buffer. A series of titrations was made with 100 cm<sup>3</sup> of 2.5 mM Na<sub>2</sub>HPO<sub>4</sub> against a titre of 2.5 mM KH<sub>2</sub>PO<sub>4</sub> and then the reverse titration was repeated with Na<sub>2</sub>HPO<sub>4</sub> as the titre. A plot of pH against molar ratio of Na<sub>2</sub>HPO<sub>4</sub> : KH<sub>2</sub>PO<sub>4</sub> gave a linear relationship ( $r^2 = 0.9997$ ) described by the following equation:

$$y = 0.4272 \text{ Ln } (x) + 6.7691$$

where  $y = \text{pH}$

and  $x = \text{the molar ratio of Na}_2\text{HPO}_4 : \text{KH}_2\text{PO}_4$

Using this equation, a buffer solution of pH 7.5 was prepared by mixing 2.5 mM phosphate solutions in a 5.54 : 1 (vol / vol) ratio of Na<sub>2</sub>HPO<sub>4</sub> : KH<sub>2</sub>PO<sub>4</sub>. The buffer solution was checked using a pH meter and found to be pH 7.5 ± 0.1.



#### 4.5 Potential activity of nitrifiers in buffered soil slurries

The low overall potential nitrifying activities recorded with perfusion tubes, despite

buffering to pH 7.5, may either reflect the intrinsically low pH of the soils (see Table 4.1), allelopathic mechanisms of long-term systems (see Section 1.5) or poor substrate / enzyme contact. However, set against the last possibility is the fact that perfusion avoids 'end-point repression' by removing the product as it is formed (see Section 1.5.4). Thus, the natural competition for N by immobilization into plants and microbial biomass is simulated. The results from the perfusion tubes were compared with other methods using incubated soil slurries to validate the findings.

#### 4.5.1 Potential activity of ammonia-oxidisers in buffered soil slurries

Samples (10 g dry soil equivalent) of stored soil from the four field replicates of the main treatments (*NF*, *GC* and *ZN*) were weighed into twelve polyethylene bottles with screw caps and pre-incubated at 20 °C for three days at a SWC of 85% field capacity to avoid waterlogging. Evaporative losses were minimized by capping the bottles, which were vented each day to maintain aerobic conditions. In the method described by Dusek (1995), pre-incubation was found to overcome the effects of sieving and storage, resulting in reduced variances of measured factors. At the start of incubation, 50 ml phosphate buffer (pH 7.5) with chlorate (15 mM) and 2 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> were added to the bottles which were transferred to a water bath at 25 °C. The bottles were shaken for 8 h and samples of the suspensions (2 ml) were withdrawn hourly, immediately centrifuged at 4000 g (4800 rpm) at 4 °C for 30 minutes and the supernatant was then directly analysed for NO<sub>2</sub><sup>-</sup>, as previously described. Potential activity was calculated as the rate of NO<sub>2</sub><sup>-</sup> accumulation using a simple linear regression and allowance was made for the progressive reduction in the volume of the slurry due to sequential sampling.




**Figure 4.3** Nitrification potential from soil perfusion columns: a) Ammonium and b) Nitrite - oxidation.

#### 4.5.2 Potential activity of nitrite-oxidizers in buffered soil slurries

The two contrasting treatments of *NF* and *ZN* were selected for assays of the kinetics of the  $\text{NO}_2^-$ -oxidizer enzyme using the method of Belser and Mays (1980). A composite sample of the stored (4 °C) field moist soil (200 g dry soil equivalent), bulked from the four field replicate sub-plots of each treatment, was pre-incubated for three days at 20 °C at a SWC of 85% field capacity. The soil sample was then blended with 1 l phosphate buffer (pH 7.5) containing  $100 \mu\text{g g}^{-1}$  (dry soil) *Didin liquid*. Aliquots (50 ml) of the slurry (containing the equivalent of 10 g dry soil) were withdrawn with an automatic pipette, whilst vigorous stirring of the mixture was maintained and were transferred into a set of 125 ml polyethylene bottles (12 for each treatment). Duplicate bottles each received 2 ml additions from stock solutions of  $\text{Na}_2\text{NO}_3$  to produce final concentrations of 50, 100, 150, 200, 300, or  $400 \mu\text{M}$  in the slurries. The bottles were placed in a water bath at 25 °C and shaken continuously. After 1, 2, 3, 4, 6, 8, 11 and 22 h, samples were taken (2 ml), centrifuged at 4,800 r.p.m. (4 °C) for 30 minutes and analysed immediately for residual  $\text{NO}_2^-$ , as before. Potential activity was calculated as the rate of  $\text{NO}_2^-$  disappearance for each substrate concentration using a simple linear regression. In addition, the affinity of nitrite-oxidizing enzymes for substrate was evaluated using Michaelis-Menten kinetics.

#### 4.5.3 Results of ammonium-oxidation measurements using incubated soil slurries

The data were analysed using the grouped regression method as before. Overall activities of the nitrifiers were higher with slurry incubation than with perfusion tubes. Rates ( $\text{nM NO}_2^- \text{g}^{-1} \text{h}^{-1}$ ) were obtained from the slopes of fitted linear regression lines (Fig. 4.4a) and increased significantly ( $P < 0.001$ ) from  $0.82 (\pm 0.231)$  in *ZN*, to  $16.9 (\pm 1.75)$  in *GC*, to  $26.2 (\pm 2.05)$  in *NF*. An apparent lag in initial oxidizing activity of about 4 h was evident



**Figure 4.4** Nitrification potential ( $\text{NH}_4^+$ -oxidation) from incubated soil slurries: regressions of mean hourly data from a) start and b) after 4 h lag.


from the plotted data so that when re-plotted using data for 4-8 h only (Fig. 4.4b), the steady state rates were nearly two-fold higher than the slopes of the 0-8 h data set: *viz.* 1.29, 30.72 and 48.75 nM NO<sub>2</sub><sup>-</sup> g<sup>-1</sup> h<sup>-1</sup> for ZN, GC and NF, respectively. These rates are well within the ranges found by other workers who have employed incubated slurries of grassland soils, *viz.* 0-236 (Sarathchandra, 1978); 21-100 (Belser and Mays, 1982); 13-39 (Berg and Rosswall, 1987); 60-100 (Dusek, 1995) and (using arable soil) 6-36 (Hojberg *et al.*, 1996).

#### 4.5.4 Kinetics of nitrite-oxidation from incubated soil slurries



The following extract is taken from Lehninger (1981): "At low substrate concentrations [S] the initial reaction velocity ( $v_0$ ) is nearly proportional to the substrate concentration, so that the reaction is approximately first order. With increasing substrate concentrations, the increase in  $v_0$  is less and no longer proportional to [S], so that the reaction is mixed order. With further increases in substrate concentration, the reaction rate becomes independent of [S] and asymptotically approaches a constant rate (see Figs. 4.5a and 4.6a). In this range of [S], the reaction is essentially zero order and the enzyme is deemed to be saturated with its substrate. All enzyme-catalysed reactions show the saturation effect, but there is a wide variation in the concentration of the substrates required to produce it."

The Michaelis-Menten equation defines the rate equation for an enzyme-catalysed reaction (single substrate), in terms of the mathematical relationship between  $v_0$ ,  $V_{max}$ , [S] and  $K_m$ :

$$v_0 = \frac{(V_{max} \cdot [S])}{K_m + [S]}$$



**Figure 4.5** Enzyme kinetics of nitrite-oxidizers in unfertilized soil (ZN): a) Michaelis-Menten plot b) Lineweaver-Burke plot.



**Figure 4.6** Enzyme kinetics of nitrite-oxidizers in fertilized soils (*NF*): a) Michaelis-Menten plot b) Lineweaver-Burke plot.

When the initial reaction rate ( $v_0$ ) is exactly one-half the velocity ( $V_{\max}$ ), then an important numerical relationship exists where:

$$K_m = [S]$$

Thus,  $K_m$  (the Michaelis-Menten constant) can be derived and is equal to the substrate concentration at which the initial velocity is half maximal ( $1/2 V_{\max}$ ). However, because the reaction rate approaches  $V_{\max}$  asymptotically, accurate evaluation of the rate is not possible, since the point of convergence is with an infinite concentration of substrate. A useful transformation of the Michaelis-Menten equation involves plotting  $1/v$  against  $1/[S]$ , the so-called 'double reciprocal' or Lineweaver-Burke plot (see Figs. 4.5b and 4.6b). The line then has a slope of  $K_m/V_{\max}$  and an intercept of  $1/V_{\max}$  and hence both  $K_m$  and  $V_{\max}$  can be accurately determined. Additionally, the **specific affinity** of the enzyme for the substrate can be derived from  $V_{\max}/K_m$ , which describes the nutrient-sequestering ability of the nitrifying population, since it combines the qualitative ( $K_m$ ) and quantitative ( $V_{\max}$ ) components of the metabolic process.

Potential nitrite-oxidizing rates ( $V_{\max}$ ) were 62.5 (ZN) and 93.4 (NF) nMoles  $g^{-1} h^{-1}$ , with  $K_m$  values of 310 and 436  $\mu M$   $NO_2^-$ -N, respectively, and were in close agreement with data for a grassland soil from Both *et al.* (1992). Thus, confirmation of the higher nitrifying activity of the fertilized soil was obtained, but a lower affinity of the enzyme for  $NO_2^-$  substrate was indicated by the larger  $K_m$  in NF. However, when the specific affinities ( $V_{\max}/K_m$ ) of the enzymes are calculated for the soils from the two treatments, there appears to be only a small difference in the ability of the two nitrifying populations to oxidize  $NO_2^-$  (202 and 214  $10^{-6} l g^{-1} h^{-1}$ , respectively). Compensatory processes apparently exist, which enable the relatively inactive nitrifying population (ZN) to develop a higher affinity for

the  $\text{NO}_2^-$  substrate, in contrast to the higher oxidizing potential of the fertilized soil (*NF*). Hence, a decline in  $V_{\text{max}}$  appears to have been compensated for by a decrease in  $K_m$  (lower  $K_m$  = higher affinity) and vice versa, as Both *et al.* (1992) also observed. The ecological advantage of this stratagem would seem to be that highly toxic  $\text{NO}_2^-$  (to both nitrifiers and plant-life) can still be effectively converted to the harmless and mobile nitrate-ion, even in systems which are not adapted for rapid nitrification. Hence,  $\text{NO}_2^-$  is seldomly found in agricultural soils in any appreciable amounts and usually  $< 0.1 \mu\text{g g}^{-1}$  soil (Van Cleemput and Samater, 1996) unless conditions which are inhibitory to the oxidation of  $\text{NO}_2^-$  develop, e.g. high  $\text{NO}_3^-$  concentration (Boon and Laudelot, 1962); high temperature (Mahandrappa, 1966); low aeration coupled with high temperature (Laudelot *et al.*, 1976).

#### 4.6 Conclusions

The marked differences in nitrifying activities in the soils under the three management regimes ( $\text{ZN} < \text{GC} < \text{NF}$ ) represent subtle responses to the contrasting N-input systems. The measurements reported here were made during a growing season in which fertilizers were withheld and grazing animals excluded, so that the changes were clearly associated with previous conditions and did not merely reflect current inputs of N. There are implications in these findings for N cycling, in general and for the fate of mineralized N linked to nitrifying activity, in particular. Laboratory evaluations of *SMB* and potential mineralization would suggest that such tests are unlikely to predict differences in actual mineralization and the subsequent oxidation of released  $\text{NH}_4^+$  needs be assessed in situations where competing processes are allowed to prevail naturally. Further work is required to establish the extent to which nitrification may influence other processes of N cycling and these aspects are discussed in more detail in Chapter 5.

## CHAPTER 5 Concurrent measurements of net mineralization, nitrification, denitrification and leaching from field incubated soil cores

### 5.1 Experimental strategy

The **development and testing** of the improved field incubation (tube) method for measuring net nitrification concluded the first of the three major aims of this project (see Section 2.1). The second aim was to **link** the *in situ* measurements of nitrification with other N cycling processes. Experiments to date have been based on established field sites, where the combination of a high annual rainfall (on average, 1050 mm) and a soil with a high clay content (c.36%) means that for much of the winter period the soils are waterlogged. Conditions conducive to high losses of N through denitrification and leaching, therefore, often prevail in spring and autumn, when soil temperatures do not fall generally to levels which would inhibit microbial activity.

Two field experiments were therefore designed in order to exploit the particular soil and environmental conditions associated with these seasonal weather patterns. Investigations into possible linkages between the supply of  $\text{NO}_3^-$  (via nitrification and/or fertilizer) and the fate of soil N through various loss processes could then be conducted. In previous experiments, fertilizer applications were discontinued and grazing animals excluded to assess the validity of the methodology by decreasing the variability between measurements within a carefully defined set of conditions. In the following series of investigations, normal fertilizer regimes were incorporated into the schedule and cattle were only excluded from sampling areas for the duration of the measurement periods. In the autumn experiment, a field site with freely drained soil was also included to enable the important

component of N leaching to be more closely examined.

## 5.2 Investigation of microbial activity following spring applications of N

In spring 1996, microbial activity was examined over a ten week period following the first and second seasonal applications of N, by taking 'spot' measurements of *SMB*, total C and total N contents, and respiratory activity. At the same time, using the improved field incubation system, net rates of ammonification, nitrification and denitrification were obtained from an integration of measurements taken over this period, to investigate associated responses in microbial activity. Before turnout, cattle-proof enclosures were erected to provide four small sub-plots (5 x 5 m) in the 200 kg N ha<sup>-1</sup> drained treatment (NF, Block A: see Fig 4.1) of the Rowden Drainage Experiment. The field drains ceased flowing after April 1996, so that further leaching of soil N was unlikely. I-E bags were, therefore, omitted from the incubation vessels and larger gypsum blocks (depth 25 mm) were constructed to occupy the space between the incubation tube and the undisturbed soil layer. On 27 March 1996, the first fertilizer was applied at a rate of 40, 26 and 50 kg ha<sup>-1</sup> of N (as NH<sub>4</sub>NO<sub>3</sub>), P and K, respectively. Fertilizer was spread by hand in two directions (at right angles across the plots) to ensure uniformity of application. Measurements began after the first application of fertilizer, with a further application of N (60 kg ha<sup>-1</sup>) only on 26 April and continued for 10 weeks until 6 June 1996. During the experimental period, the weather was generally wet and dull, e.g. May was colder (mean temperature of 8.9 °C) and wetter (with 80.7 mm rain) than average (30 year monthly means of 10.7 °C and 65.2 mm, respectively).

### 5.2.1 Net ammonification, nitrification and denitrification

The novel ('tube') field incubation technique was employed to measure *in situ* net ammonification, nitrification and denitrification during the experimental period. Each week, 12 cores enclosed in PVC tubes (37 mm diameter, 150 mm length) were extracted (three cores per sub-plot) along with 12 paired soil samples (extracted as outer concentric layers from around the PVC tube) using a larger corer (80 mm diameter, 300 mm length). The outer soil cores were extracted within 1 h of collection in 2 M KCl (soil:extractant volume ratio of 1:5) and analysed for  $\text{NH}_4^+$  and  $\text{NO}_3^-$ -N. The sheathed cores were incubated in the field for 7 days: six cores received a continual, slow infusion of  $\text{C}_2\text{H}_2$  (from controlled release capsules, see Section 2.7.1) and the other six cores were incubated without  $\text{C}_2\text{H}_2$ . At the end of each 7 day incubation period, the sheathed cores were removed and extracted, as before and a new set of cores were obtained for incubation. Net rates for the N transformation processes were calculated from previously established relationships (see Section 3.4). Thus, an integrated measure of net mineralization (ammonification) was obtained from the increase in ammonium ( $\text{NH}_4^+$ )-N in the six cores incubated in the presence of  $\text{C}_2\text{H}_2$ , over the ten week period. An estimate of net nitrification was obtained from the difference in  $\text{NH}_4^+$ -N produced in the set of cores incubated with  $\text{C}_2\text{H}_2$  minus that produced in cores incubated without  $\text{C}_2\text{H}_2$  (i.e. the amount of  $\text{NH}_4^+$ -N that would have been nitrified, assuming immobilization was the same in both sets). Denitrification was estimated by the difference in  $\text{NO}_3^-$ -N accounted for with (denitrification denied) and without (denitrification not denied)  $\text{C}_2\text{H}_2$ .

### 5.2.2 Soil microbial biomass (SMB)

During the period of the experiment, *SMB* was investigated on five separate occasions at

intervals of 14 days. Soil was sampled using a corer (2.5 cm diameter, 15 cm deep) and the soil cores taken from each sub-plot were separately crumbled and sieved (4 mm) and stones, above-ground plant material and visible roots were discarded. A sub-sample was dried overnight at 105 °C for determination of water content. Established techniques (Brookes *et al.*, 1985; Vance *et al.*, 1987) were employed to identify and characterize *SMB* using the fumigation/extraction method (FE). On the days when soil was collected, duplicate samples of moist soil (12.5 g) were weighed into 60 ml glass bottles and the first of each pair was extracted with 50 ml 0.5 M  $K_2SO_4$  by shaking the mixture for 1 h on an end-over-end shaker. The suspension was then filtered (Whatman No. 1) and the filtrate frozen for storage. The second sample was fumigated in a desiccator, lined with wet tissue paper and containing 35 ml ethanol-free chloroform ( $CHCl_3$ ) in a beaker with glass boiling granules. Ethanol (included as a commercial stabilizer) was first removed from 50 ml  $CHCl_3$  (to avoid erroneously high measurements) by shaking with 15 ml water in a separating funnel and then drawing off the  $CHCl_3$  from below. The process was repeated twice more, each time discarding the water phase (containing the ethanol contaminant); the last traces of water were finally removed from the  $CHCl_3$  by adding fused granules of  $CaCl_2$ . The desiccator was evacuated and soil samples fumigated for 24 h, after which the beaker of  $CHCl_3$  was removed. Residual  $CHCl_3$  vapour was dispersed by alternating between evacuation and a controlled release of the vacuum back to atmospheric pressure. The soil was then extracted, as for the unfumigated sample, and frozen for storage. The extracts were used to determine biomass C and N, and results were expressed on an oven dry soil basis.

**Biomass N** The fumigated (F) and unfumigated (NF) extracts were thawed in a refrigerator

prior to Kjeldahl digestion which involved boiling 25 ml extract (including glass boiling granules) with 0.6 ml 0.19 M CuSO<sub>4</sub> and 10 ml concentrated H<sub>2</sub>SO<sub>4</sub>. The digestion process was found to proceed more gently when the acid was thoroughly incorporated in the mixture and allowed to stabilize overnight before any heat was applied. The temperature of a heating block was first raised gradually to 150 °C and when this was reached, the digest boiled vigorously. Further heat was then applied up to 200 °C and maintained until the volume had been reduced to about 10 ml (with no water remaining). Final digestion was at 400 °C, refluxed for 3 h. After cooling, the digest was made to standard volume (75 ml) with water, mixed and stored at 4 °C. The total N content (as NH<sub>4</sub><sup>+</sup>-N) was determined colorimetrically (flow injection analyser). Biomass N was calculated using the formula of Jenkinson (1988):

$$\text{Biomass N} = 2.22 (\text{Kjeldahl N in F extracts} - \text{Kjeldahl N in NF extracts})$$

Biomass C Sub-samples of the F and NF extracts (8 ml) were digested with 2 ml 0.067 M K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub>, 70 mg HgO and 15 ml of a 2:1 v/v mixture of H<sub>2</sub>SO<sub>4</sub> and H<sub>3</sub>PO<sub>4</sub> (both as concentrated acids). After boiling under a water reflux for 30 min, the excess dichromate was determined from a back titration with standardized 0.33 M (NH<sub>4</sub>)<sub>2</sub>Fe(SO<sub>4</sub>)<sub>2</sub>·6H<sub>2</sub>O in 0.4 M H<sub>2</sub>SO<sub>4</sub>. Biomass C was calculated using the formula of Vance *et al.* (1987):

$$\text{Biomass C} = 2.64 (\text{organic C in F extracts} - \text{organic C in NF extracts})$$


Biomass activity Sub-samples of the sieved soil from each sampling occasion were also placed in sealed serum bottles at 22 °C and CO<sub>2</sub> concentrations in the headspace were measured initially, and again after 4 hours of incubation, using an I.R. gas analyser.

Specific respiration was calculated as µg CO<sub>2</sub>-C produced by 1 mg biomass-C per g dry


soil per hour. An alternative approach to measuring substrate-induced respiration (SIR) to the previous method (see Section 4.3.2) was used, which employed 'dry' (as opposed to liquid) glucose additions (5 mg glucose per g equivalent dry weight of soil) which can give more sensitive responses (Anderson and Domsch, 1978). Selective respiratory inhibition (Anderson and Domsch, 1975) was again used to discriminate between fungal and bacterial components (see Section 4.3.2).

### 5.2.3 Results from the spring studies

A net release of  $\text{NH}_4^+$ -N corresponding to the pattern of *SMB* activity was recorded (Fig. 5.1a) with steadily declining rates of net ammonification following the first spring application of N, viz. 1.80 to 0.88 kg N ha<sup>-1</sup> d<sup>-1</sup>. After the second N application, rates increased significantly to 2.20 kg N ha<sup>-1</sup> d<sup>-1</sup> ( $p < 0.05$ ) when compared with samples taken immediately prior to fertilizer application (viz. 0.82) and a steady decline in rates was again repeated, reaching a minimum of 0.57 followed by a slight increase to 0.89. Therefore, N applications appeared to have directly stimulated rates of ammonification, which has been referred to as a 'priming effect' by other workers (see Haynes, 1986). Nitrification of the released  $\text{NH}_4^+$ -N was rapid and suggested a similar pattern in activity to ammonification (Fig. 5.1b), with maximum rates initiated by the two fertilizer events, 1.6 and 1.38 kg N ha<sup>-1</sup> d<sup>-1</sup>, respectively. However, the rates of the two processes were only weakly correlated ( $r^2=0.41$ ) with 33% of the variation accounted for; 63% of the net  $\text{NH}_4^+$ -N released was nitrified to  $\text{NO}_3^-$ -N. Denitrification of  $\text{NO}_3^-$ -N also appeared to have been stimulated by the N fertilizer applications (Fig. 5.2) with maxima of 0.75 and 0.54 kg N ha<sup>-1</sup> d<sup>-1</sup> recorded and an overall loss of N for the period of 20.8 kg (representing 36% of the nitrified N).




**Figure. 5.1 a** Net ammonification ( $\text{kg N ha}^{-1} \text{d}^{-1}$ ) following N applications (closed circles  $\pm$ SEM) and after withholding N (open circle  $\pm$ SEM), and specific respiration ( $\mu\text{g CO}_2\text{-C mg}^{-1}\text{Bio-C g}^{-1} \text{h}^{-1}$ ) of biomass (bars  $\pm$ SEM), in soil from a fertilized grass sward during spring (n=4). **b** Nitrification ( $\text{kg N ha}^{-1} \text{d}^{-1}$ ) following N applications (closed circles  $\pm$ SEM) and after withholding N (open circle  $\pm$ SEM), in soil from a fertilized grass sward during spring 1996 (n=4).



**Figure 5.2** Denitrification ( $\text{kg N ha}^{-1} \text{ d}^{-1}$ ) following 2nd spring (1996) application of N (closed circles  $\pm$ SEM) and weekly precipitation (mm) shown as bars ( $n=4$ )

When rates of denitrification and nitrification were compared (omitting negative or zero rates) the correlation between the two processes was poor ( $r^2=0.16$ ). However, one of the data points was shown to be associated with a high residual error and low leverage (Cooke's distance residuals/leverage test), but with no obvious/recorded analytical error. A considerably stronger correlation was obtained ( $r^2=0.98$ ) when this outlying point was omitted from the regression (Fig. 5.3). Thus, circumstantial evidence for a direct linkage between the processes was obtained, suggesting that the supply of  $\text{NO}_3^-$ -N from fertilizer and/or nitrification can strongly influence the rate of denitrification. No significant difference was found between the mean  $\text{NO}_3^-$ -N contents ( $\mu\text{g N g}^{-1}$ ) of soil sampled before ( $2.6 \pm 0.25$ ) and after incubation with  $\text{C}_2\text{H}_2$  ( $2.2 \pm 0.45$ ), which confirmed the effectiveness of the controlled-release capsules in preventing nitrification.

Despite these independent measurements showing enhanced microbial activity, the overall



**Figure 5.3** Correlation between nitrification and denitrification; one outlying value (open circle) omitted from regression analysis

size, in terms of biomass-C, was unchanged. Thus, *SMB* (measured by FE) contained nearly 1000 mg C per kg (dry) soil, but this did not alter significantly during the ten week period of the experiment (Table 5.1).

**Table 5.1** Total C and total N in *SMB* (dry soil wt. basis) from a fertilized grass sward during spring 1996 (4 reps.)

Sampling dates	Biomass C		Biomass N		Biomass C:N
	mgC kg <sup>-1</sup> soil	SEM	mgN kg <sup>-1</sup> soil	SEM	
4 April	922.2	103.53	148.2	11.63	6.22
18 April	947.7	69.61	149.6	22.29	5.73
2 May	996.4	49.85	158.8	14.46	6.35
16 May	993.4	115.78	158.6	13.34	6.21
30 May	1184.0	56.95	183.3	5.81	6.45

Biomass N content was also unchanged during the experiment with an overall mean value

of 160 mg N kg<sup>-1</sup> (dry) soil; consequently the C:N ratio of the biomass averaged 6.2 throughout (Table 5.1). Other studies have shown the *SMB* to be remarkably stable under field conditions (Bristow and Jarvis, 1991; Witter *et al.*, 1993) and additions of fertilizer N had no effect on biomass size over the course of one year (Lovell *et al.*, 1995). However, total *SMB* (as measured by FE) does not necessarily reflect its activity, since the proportions of active and inactive components may change due to environmental factors (Wardle and Parkinson, 1990), or with additions of substrate (Ocio and Brookes, 1990). **Substrate-induced respiration** (SIR) had rates between 20.3 and 27.5 µl CO<sub>2</sub> g<sup>-1</sup> h<sup>-1</sup>, but also showed no discernable trend with time. There were also no marked changes in the ratio of fungi : bacteria (assessed by selective respiratory inhibition) over the period of the experiment; overall the mean ratio was 3.0 : 1, indicating that the biomass contained a consistently higher proportion of fungi. **Specific respiration** (Fig. 5.1a), which more closely reflects the metabolic activity of the biomass, showed a decrease from a maximum of 5.1 (µg CO<sub>2</sub>-C mg<sup>-1</sup> biomass-C g<sup>-1</sup> h<sup>-1</sup>) immediately after the first application of N, to a significantly lower level of 3.0 (p<0.05) by week 5. Following the second application of N, this downward trend was halted and by week 7, specific respiration was no longer significantly different from the initial value. After a further two weeks (week 9), however, specific respiration had again fallen significantly (p<0.05) below the initial value. This response to N suggests that the activity of the microbial population may have been limited to some extent by an inadequate supply of N. In this experiment, therefore, additions of fertilizer N coincided with increased *SMB* activity, without the size of the biomass (as assessed by FE extraction) being measurably affected. Lovell and Jarvis (1996) found that in soil treated with urine, biomass-mediated processes also increased substantially, whilst biomass size remained the same. Joergensen (1996) has proposed that there is an active

component of the *SMB*, rather than its overall size, which varies during the year in response to altered conditions.

The existence of a large, but mainly dormant bacterial population, suggested by Paul and Voroney (1980), may partly explain the relatively stable component identified by FE extraction. The other changes observed in various aspects of *SMB* activity are most likely to be as an indirect effect of the increased inorganic N which increases the supply of C substrates through enhanced root growth and exudation into the rhizosphere (Fauci and Dick, 1994). Respiratory activity is relatively easy to measure and could provide an indicator of short-term responses in microbial activity to external influences. In the case of nitrifying processes the increase in substrate ( $\text{NH}_4^+\text{-N}$ ), supplied either directly from fertilizer, or indirectly as a result of stimulated mineralization (i.e. from organic N), appears to have increased the rate of nitrification, as was recently shown by Watson and Mills (1998). Thus, supporting evidence for the role of nitrification in responding to N and promoting losses through denitrification has been obtained and was examined further in the next experiment.

### 5.3 Investigation of N transformations in autumn/winter

In the autumn experiment, measurements followed a growing season with grass under conventional silage management, so that nitrifying activity *per se* could then be assessed without the complication of further inputs of fertilizer N. Two soil types, with contrasting drainage characteristics, were selected at the farm of the Institute of Grassland and Environmental Research, Okehampton, Devon, UK to test further the performance of the 'tube' incubation method. The first was a freely drained, reddish gravelly loamy soil over

Permian breccia from the Crediton Series: a typical brown earth (Findlay *et al.*, 1984). The second was a poorly drained clay loam soil of the Hallsworth series (see Section 3.2 for full classification). Both soil types had been under a 3-cut grass silage management system with similar inputs of N for the last 5 and 4 years, but had previously contrasting histories of continuous arable cropping and long-term grassland, respectively. Some important chemical and physical properties of the soils are presented in Tables 5.2, 5.3 and 5.4.

**Table 5.2** Soil particle size distribution (%) in the Crediton and Hallsworth soil series (0-150 mm depth). Particle sizes in mm.

Soil series (drainage)	Clay (<0.002)	Silt (0.002- 0.02)	Fine sand (0.02-0.2)	Coarse sand (0.2-2.0)	OM*
Crediton (free)	24.2	10.3	14.9	51.7	5.6
Hallsworth (poor)	48.5	12.7	16.3	7.7	14.1

\*OM denotes organic matter (not including living roots) assessed by loss of mass on ignition

**Table 5.3** Percentage by volume of soil, stone, water- and air-filled pore spaces at field capacity in freely and poorly drained soils at IGER, North Wyke, UK (January 1997)

Soil series (drainage)	Fine soil (%)	Stone >2 mm	Pore space (% volume)			Dry bulk density (g soil cm <sup>-3</sup> )
			Water	Air	Total	
Crediton (free)	44.9	5.0	35.7	14.4	50.1	1.19
Hallsworth (poor)	27.2	0.5	46.8	25.5	72.3	0.72

**Table 5.4** Some properties of the freely and poorly drained soils (0-150 mm depth) measured at the end of the experimental period (oven dry weight basis)

Soil series (drainage)	Inorganic N ( $\mu\text{g N g}^{-1}$ )	$\text{NH}_4^+ : \text{NO}_3^-$	Total N ( $\text{mg N g}^{-1}$ )	Organic C ( $\text{mg C g}^{-1}$ )	C : N	pH
Crediton (free)	3.4	3.8	3.3	24.0	7.3	5.3
Hallsworth (poor)	5.9	3.9	6.3	57.8	9.2	5.1

Each soil type was represented by four replicate plots (500 m<sup>2</sup>). In 1996, N applications to freely and poorly drained soils amounted to 285 and 297 (kg ha<sup>-1</sup>) fertilizer and 128 and 98 (kg ha<sup>-1</sup>) slurry total N, respectively. Three silage cuts were made in 1996, the last being on 1 October for both soil types. From 31 October 1996 to 3 January 1997, measurements were made using the 'tube' incubation system; two paired soil cores (each including associated outer cores) were taken at random from within each of the field plots every 4-5 days (11 days on one occasion) in incubation tubes, one soil core to be incubated with C<sub>2</sub>H<sub>2</sub> and the other without C<sub>2</sub>H<sub>2</sub>. Each of the 16 cores was then incubated in the field in incubation vessels with an I-E bag and gypsum block (15 mm depth) placed beneath each core, half of the blocks (8) contained controlled release capsules (incubations with C<sub>2</sub>H<sub>2</sub>). At the end of each incubation period, the soil core was removed and NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup>-N were extracted using 2 M KCl extractant (5 : 1, KCl : fresh soil) together with a new set of outer cores, representing the starting 'baseline value' for the succeeding incubation. The NO<sub>3</sub><sup>-</sup> absorbed in the I-E bags was extracted with 1.5 M H<sub>2</sub>SO<sub>4</sub>, as before

(see Section 2.4.3) and the recovery factor of 90% was used to adjust estimates of potential leaching, based on the efficiency of  $\text{NO}_3^-$  recovered in previous tests.

Denitrification was also measured independently over the experimental period by the 'jar incubation' method (Ryden *et al.*, 1987), using 8 cores (100 mm deep x 25 mm diameter) per jar (four replicate jars per soil type). At the end of the experiment, over-winter herbage production from the swards on the two soil types was assessed by taking four replicate quadrats ( $1\text{m}^2$ ) to a cutting height of 40 mm. The herbage was dried at  $85^\circ\text{C}$  for 12 h in a forced-draught oven, ground and analysed for total N content using a Carlo Erba CN Analyser. Potential nitrifying activity was estimated from soil cores obtained at the end of the experiment (four replicates of 10 bulked cores: 150 mm deep x 37 mm diameter). The soil was sieved (6 mm) and fresh soil (equivalent to 10 g dry weight) was incubated in a water bath at  $25^\circ\text{C}$  as a 'slurry' (50 ml) of 2 mM  $(\text{NH}_4)_2\text{SO}_4$ , 2 mM phosphate buffer (pH 7.5) and 15 mM  $\text{NaHClO}_3$ , as before and the rate of  $\text{NO}_2^-$  production over 8 h was measured (see Section 4.5.1).

Leaching of  $\text{NO}_3^-$ -N was independently assessed using ceramic suction cups (25 mm diameter, 55 mm length) inserted at a depth of 600 mm in the freely drained soil and at 300 mm in the poorly drained soil (12 replicates to each soil type). Estimates of net mineralization were also independently obtained by the method of Hatch *et al.* (1990) using a 7 d field incubation of 4 cores (125 mm deep x 37 mm diameter) per incubation vessel, with  $\text{C}_2\text{H}_2$  to inhibit nitrification. There were four replicate incubations to each soil type. The original method was altered slightly so that the assessment of net mineralization was based only on the change in soil  $\text{NH}_4^+$ -N during incubation, rather than the overall change

in inorganic N (i.e.  $\text{NH}_4^+ + \text{NO}_3^-$ ). This was based on the assumption that, as shown in previous studies (Hatch *et al.*, 1991; Gill *et al.*, 1995),  $\text{C}_2\text{H}_2$  is an effective inhibitor of nitrification and there would be no increase in  $\text{NO}_3^-$  levels.

### 5.3.1 Methods of soil analyses

Total C, total N and pH were also measured on oven-dried sub-samples of the bulked soils. Soil particle size analysis (Piper, 1950) was obtained from air-dried (30 °C) samples, crushed to pass a 2 mm sieve. Sub-samples, 100 g soil (Crediton series) or 50 g (Hallsworth series) were mixed with 5 ml saturated solution of sodium oxalate, 5 ml 1 M NaOH and 10 ml hydrogen peroxide (6 %  $\text{H}_2\text{O}_2$  v/v). Sufficient deionized water was added to saturate the samples, which were broken up using a pestle into a smooth lump-free paste. The Hallsworth soil also had a high O.M. content and required further additions of  $\text{H}_2\text{O}_2$  and gentle heating on a water bath to disperse the aggregates and destroy the O.M. The reaction was completed when the mixture no longer frothed (> 24 h, in the case of Hallsworth soil). The remaining  $\text{H}_2\text{O}_2$  was then destroyed by boiling for a few minutes. The mixtures were then vigorously stirred for about 15 min (30 min for the clay soil) and washed into a Bouyoucos cylinder, using a wash bottle. Further water was added up to the 1205 ml mark (100 g sample) or the 1130 ml mark (50 g sample) with the hydrometer in place. The cylinder was then inverted (without the hydrometer), ten times and then allowed to settle. Both the time and temperature of the mixture were noted and after 3 min, the hydrometer was gently inserted and read at 4 min. Prior to taking the reading, the hydrometer was depressed slightly and allowed to settle, to avoid an erroneous reading from froth or surface tension effects. A few drops of amyl alcohol (=Pentan-1-ol) helped to disperse the froth to make the recording clearer. The hydrometer was then removed and

the mixture left to settle further for 2 h, after which another hydrometer reading was obtained, along with the temperature of the mixture, at that time. Readings were corrected from the calibrated value (19.5 °C), by adding 0.3 for every 1 °C above this value and deducting 0.3 for every 1 °C below it. The second reading (representing only the clay component) was subtracted from the first reading (representing silt + clay components) to give the silt only component. Percentages were obtained by dividing these component parts by the amount of soil from which they were derived (on a dry soil weight basis), multiplied by 100. The liquid was decanted from the cylinder through a 70 mesh sieve over a bowl. After sieving in water for 1 min, the coarse sand (on the sieve) was transferred to an evaporating dish and dried in an oven (105 °C), whilst the remaining contents of the bowl were washed into a 1 l beaker to a depth of 10 cm. The fine sand was separated from the silt by vigorously stirring the sediment, followed by a period of settling-out which was determined from the temperature of the mixture, according to the following table:

**Table 5.5** Settling time required before decantation of silt to retain fine sand, as determined from the temperature of the washings, after removal of coarse sand

°C	16	17	18	19	20	21	22	23	24	25
secs	320	310	300	300	288	280	270	270	260	255

The process was continued until the liquid ran clear and the fine sand was then dried in an evaporating dish, as before. The coarse and fine sands were expressed as percentages of the original (dry) soil sample. Organic matter content (not including living root matter) was obtained from the loss of weight on ignition of soil placed in a muffle furnace and

heated overnight at 600 °C and expressed as a percentage component of the dry soil.

### 5.3.2 Results from the autumn studies

Net rates (kg N ha<sup>-1</sup> d<sup>-1</sup>) of mineralization in the soils measured by tube incubation (Fig. 5.4a and b) ranged from 0.28 - 1.64 (freely drained) and 0.55 - 1.17 (poorly drained). The mean daily rates (kg N ha<sup>-1</sup>) for the two soils over the measurement period were significantly different ( $p < 0.001$ ), 0.59 ( $\pm 0.064$ ) in freely drained soil compared with 0.88 ( $\pm 0.056$ ) in poorly drained soil. Cumulative totals for the net release of mineralized N from soil organic matter (Table 5.6) compared favourably with independent measurements using the jar incubation technique over the same measurement period.

**Table 5.6** Comparison between the tube incubation method and alternative methods of measuring several N transformation processes over 64 days (31/10/96-3/1/97)

Process	Freely drained soil (kg N ha <sup>-1</sup> )		Poorly drained soil (kg N ha <sup>-1</sup> )	
	Tube incubation /resin bags*	Jar incubation /ceramic cups#	Tube incubation /resin bags*	Jar incubation /ceramic cups#
Net mineralization	34.8 (1.29)	56.1 (0.58)	53.9 (0.39)	43.2 (1.17)
Nitrification	20.2 (2.42)	-	38.7 (0.97)	-
Denitrification	7.2 (3.39)	5.3 (0.15)	24.2 (3.51)	7.1 (1.76)
Leaching	3.7 (0.71)*	3.7 (1.73)#	6.9 (0.88)*	3.8 (2.05)#


Data are means with  $\pm$  SEMs shown in parentheses

Coefficients of variation were higher than would be anticipated under strictly controlled analytical conditions, but of the order commonly encountered with field measurements in grassland soils (mean values of 52 % and 38 %, for freely and poorly-drained soil,


respectively). In the freely drained soil, 58 % of mineralized N was nitrified, compared with 72 % in the poorly drained soil. Daily rates ( $\text{kg N ha}^{-1} \text{ d}^{-1}$ ) of nitrifying activity ranged from 0.03 - 0.82 (Fig. 5.4a) and 0.37 - 0.95 (Fig. 5.4b), with mean daily rates for the whole period of 0.24 ( $\pm 0.072$ ) and 0.60 ( $\pm 0.074$ ), respectively. No independent method for assessing nitrifying activity in the field was available, but laboratory measurements of potential nitrification (where  $\text{NH}_4^+$ -N substrate was non-limiting) gave rates of 14.7 and 8.3 nMoles  $\text{NH}_4^+$  oxidized  $\text{g}^{-1}$  (dry soil)  $\text{h}^{-1}$ , obtained from the slopes of linear plots of  $\text{NO}_2^-$  production over 8 h ( $r^2 = 0.99$  and  $0.97$ ) in the freely and poorly drained soil, respectively. Thus, higher ( $p < 0.001$ ) nitrifying rates were found in the field in the poorly drained soil, but the greater potential activity was in the freely drained soil.

Other factors which might have been expected to correspond with different rates of nitrification were similar (Table 5.4), e.g. soil pH and the ratios of  $\text{NH}_4^+ : \text{NO}_3^-$ . Nitrification was related to rates of mineralization ( $p < 0.001$ ) in both the freely drained ( $r^2 = 0.41$ ) and in the poorly drained ( $r^2 = 0.52$ ) soil, as might be expected, since the supply of substrate (*viz.*  $\text{NH}_4^+$ ) is determined by the rate of mineralization. Also,  $\text{NH}_4^+$  contents exceeded those of  $\text{NO}_3^-$  in both soil types, which confirmed this dependence and showed that there was a lag in the oxidation of  $\text{NH}_4^+$  by the nitrifying population which allowed  $\text{NH}_4^+$  to accumulate.

Rates ( $\text{kg N ha}^{-1} \text{ d}^{-1}$ ) of loss of N by denitrification (estimated by differences in inorganic N from weekly tube incubations  $\pm \text{C}_2\text{H}_2$ ) ranged from 0.04 - 0.52 (Fig. 5.5a: freely drained) and from 0.17 - 1.08 (Fig. 5.5b: poorly drained). The mean rates ( $\text{kg N ha}^{-1} \text{ d}^{-1}$ ) differed between the two soils ( $p < 0.01$ ), i.e. 0.05 ( $\pm 0.086$ ) compared with 0.40 ( $\pm 0.079$ ),





**Figure 5.4** Weekly measurements shown as rates ( $\text{kg N ha}^{-1} \text{d}^{-1}$ ) of net mineralization (closed circles  $\pm$  SEM) and nitrification (open circles  $\pm$  SEM) in a) freely and b) poorly drained soil over an experimental period (64 d) in autumn/winter 1996/7 ( $n = 4$ )



**Figure 5.5** Weekly measurements shown as rates ( $\text{kg N ha}^{-1} \text{d}^{-1}$ ) of denitrification in a) freely and b) poorly drained soil over an experimental period (64 d) in autumn/winter 1996/7. Vertical bars are  $\pm$  SEM ( $n = 4$ )

respectively. Daily rates of denitrification were closely correlated with rates of nitrification in the freely drained soil ( $p < 0.001$ ), with  $r^2 = 0.67$  (one outlying value was removed with high residual error, but with low leverage, Fig. 5.6a). However, rates of denitrification also included some negative values which were assumed to be zero (i.e. at or near a rate below the level of detection) in the calculation of the cumulative total (Table 5.6). In the poorly drained soil, there were fewer negative values (Fig. 5.6b) and the two processes were again significantly correlated ( $p < 0.001$ ), with  $r^2 = 0.68$  (two outlying values removed with high residual errors, but with low leverage). The cumulative total was  $> 3$ -fold larger than that of the freely drained soil (Table 5.6).

The cores remained tightly fitted in the incubation tubes during incubation, so that preferential leaching down the sides was unlikely and shrinkage of the soil within the tubes did not occur. Both of the techniques used for measuring leaching (by I-E bags or ceramic suction cups) and those used for denitrification (by difference in the tube incubation, or by jar incubation) were in close agreement in the freely drained soil (Table 5.6). In the poorly drained soil, however, the estimates were greater with the tube incubation method for both processes (Table 5.6). The N unaccounted for in measurements taken by tube incubation over 64 d (i.e. net mineralization minus losses from denitrification and leaching) amounted to 24.0 and 22.8 kg N in the freely and poorly drained soil, respectively. Some of this N loss could be attributed to uptake in the autumn/winter herbage growth (assessed from herbage sampled from these plots at the end of the experimental period) which yielded 26.2 ( $\pm 1.60$ ) and 55.8 ( $\pm 6.81$ ) kg N, respectively. No assessment was made of the N in herbage at the start of the experimental period (October 1996), so it is possible that more N was present in re-growth (i.e. before soil measurements began) following the last silage



**Figure 5.6** Correlations (closed circles) between nitrification and denitrification in **a)** freely drained soil and **b)** poorly drained soil from weekly measurements: outlying values (open circles) omitted

cut on the poorly drained soil (due to the wetter soil conditions) and was therefore over-estimated, whereas growth on the drier soil had ceased after cutting and regrowth only occurred after the soil had returned to field capacity. This discrepancy would have been carried forward into the over-winter assessment of uptake from N mineralized during the measurement period, but nevertheless provides an estimate of removal into herbage. Losses of N through senescence and transport into roots, which were not accounted for, would also contribute to errors in the estimates on both soil types.

### 5.3.3 Conclusions from the autumn/winter studies

The tube incubation method allows concurrent measurements of the major N transformation processes (mineralization, nitrification, denitrification and leaching) to be made which would not otherwise be possible using existing techniques. Foremost amongst the advantages is the opportunity to extend the period of measurement indefinitely, which would not be practical with other methods, particularly for gross measurements using  $^{15}\text{N}$  methodology (due to the expense and other technical limitations, see Section 6.1). An additional benefit of the improved incubation method is that integrated estimates of the various processes are obtained for defined periods (which avoids problems of long-term exposure/supply of soil to  $\text{C}_2\text{H}_2$ , see Section 2.7), rather than the alternative method of extrapolating from 'spot' measurements. The present experiment, which was designed to illustrate the use of the technique under field conditions, showed that the processes of net mineralization, nitrification and denitrification were closely associated. Such associations might be expected, given the substrate/product relationships which exist, although supporting evidence is limited. The effect that antecedent fertilizer inputs can have on increasing the rate of nitrification has been shown recently (Jarvis and Barraclough, 1991;

Hatch *et al.*, 1996; Watson and Mills, 1998) and this will have implications in terms of the supply, distribution and fate of N in soils. Further possible correlations might also have been expected between nitrifying activity and the removal of nitrate through leaching. However, the lack of drainage in December 1996 (driest December since 1959) precluded meaningful comparisons.

Errors associated with the measurements were within the range typically encountered in field experimentation. However, some negative values were found for nitrification, denitrification and also leaching which were obviously incorrect, whereas negative rates for mineralization are probably more correctly interpreted as net immobilization. The reason for these discrepancies was probably the combination of inherently low rates and high spatial variation between replicates. Greater replication (e.g. 6-8 incubation tubes to each treatment) would be required to better resolve temporal changes in these processes. In overall terms, the values obtained with tube incubation and other methods compared well, with the exception of denitrification in the poorly drained soil. Lower than expected estimates of denitrification in 'heavier' soils are recognized as particular problems of the jar incubation method (Smith and Arah, 1990). This may be due to enhanced aeration and an inability to ensure satisfactory infusion of  $C_2H_2$  to fully inhibit nitrous oxide reductase activity (i.e.  $N_2O$  to  $N_2$ ). A marked advantage of the tube incubation method is, therefore, the enclosure of the soil core in a sheath to prevent exposure of the sides of the soil cores to unnatural aeration and the improved method of providing a sustained infusion of  $C_2H_2$  via the base of the incubation tube and by gas diffusion holes placed around the sides (see Fig. 2.1). Degradation of  $C_2H_2$  by  $C_2H_2$ -metabolizing populations is also recognized as a potential problem (with the possibility of wider effects on C cycling processes), but is one

which usually only develops after a lag phase of up to 8 d (Terry and Duxbury, 1985). The greater variability in the data relating to denitrification in the freely drained soil, however, was probably caused by an inadequate supply of  $C_2H_2$  from the controlled-release capsules, because of very dry soil conditions. It may be necessary in such circumstances, periodically (e.g. halfway through the incubation period) to re-wet the gypsum blocks to ensure that the reaction with  $CaC_2$  granules proceeds satisfactorily.

Differences in assessments of nitrifying activities (actual and potential) are less easy to reconcile. The greater potential nitrifying activity in the freely drained than the poorly drained soil suggests that a more active, or abundant population of nitrifiers may have developed in the freely drained soil in response to aerobic conditions more favourable for growth of the bacteria. The higher (actual) field rate of nitrification found with the poorly drained soil is, however, compatible with having a higher rate of net mineralization (since the supply of  $NH_4^+$  substrate will be the rate-limiting step). It is also possible that this soil type, which is more highly structured, may have microsites with a higher pH which would favour nitrification, despite the apparent similarity in pH of the two soils when measured on disturbed soil samples. Such differences would not be a factor in the laboratory incubations which were based on soil 'slurries' buffered at pH 7.5.

Estimates of N leached using both I-E bags and cup-samplers were low and reflect the unusually dry winter period of 1996/7 and generally lower losses associated with cut (as opposed to grazed) swards (Ryden *et al.*, 1982). It would, therefore, be of interest to assess leaching under grazed swards in a more typical drainage period, but a considerable increase in replication would be required to overcome the greater variability in soil inorganic N

content induced by excretal returns. The 'paired' sampling approach, adopted in the present method, has an advantage over random sampling since spatial variation is largely accounted for by closely linking the 'start' cores with the incubated cores (*viz.* concentrically positioned cores). I-E bags also have the advantage over cup-samplers of operating in a 'passive' mode to the passage of soil water, whereas cup-samplers may modify the flow characteristics of soil water and are less satisfactory in 'heavier' soils (Hatch *et al.*, 1997). The tube incubation method also overcomes many of the other problems associated with enclosed incubation vessels, causing only minimal disturbance to the soil which allows natural wetting/drying processes to proceed and avoids large changes in the soil atmosphere/temperature status. Finally, the low cost of the equipment involved in the tube incubation method enables large scale monitoring for prolonged periods of experimentation to be undertaken, as well as providing the opportunity for examining the linkage between N cycling processes.

The extent, however to which nitrification may exert a controlling influence over components of the N cycle can only be determined further from direct measurements of the gross rates of the processes involved. Chapter 6 describes the final part of the project which was devoted to **examining** the practical implications of these findings under different management regimes and thereby addresses the third aim of the project as stated in Section 2.1. The natural stable isotope of nitrogen ( $^{15}\text{N}$ ) was employed, using techniques of 'mean pool dilution', to determine rates of the key processes. The tube incubation method was adapted to provide microplots for the  $^{15}\text{N}$  enrichment experiment and simultaneous estimates of net process rates. The development of the equipment, experimental techniques and the calculations involved in resolving these processes are described.

## **CHAPTER 6 Development of a multi-point soil injector to measure gross and net rates of the major N cycling processes**

### **6.1 Introduction: concepts and constraints**

A further development of the tube incubation technique (previously described in Chapter 2) was undertaken in order to combine net measurements of mineralization, immobilization and nitrification with gross rates of the same processes, obtained using standard  $^{15}\text{N}$  pool dilution methodology (Kirkham and Bartholomew, 1954; Barraclough, 1991). The basic concept was to provide an alternative calibration of the tube incubation technique, together with simultaneous assessments of the gross rates of the major processes (using a specially designed  $^{15}\text{N}$  soil injector) to underpin the findings of the previous chapter.

A popular approach employed in field evaluations using the  $^{15}\text{N}$  mean pool dilution technique involves sets of paired microplots (open-ended tubes inserted into the ground and enclosing undisturbed soil profiles). The microplots are enriched with a source of the stable isotope of nitrogen ( $^{15}\text{N}$ ):  $^{15}\text{NH}_4^+\text{-N}$  for gross rates of mineralization/immobilization and  $^{15}\text{NO}_3^-\text{-N}$  for gross rates of nitrification (Barraclough, 1991; Monaghan, 1995; Davidson *et al.*, 1991). An advantage of this approach is that only the products, rather than the substrates of these processes are supplemented, which avoids stimulation and possible disturbance of the steady state. Following injection (at  $t_0$ ), assessment of the starting point enrichment is made (after 24 h equilibration period) by removing one of the paired microplots ( $t_1$ ) and then comparing this with the resultant enrichment in the remaining microplot, after a defined period of field incubation has elapsed ( $t_1+$ ).

The underlying assumption is that both native  $^{14}\text{N}$  sources and  $^{15}\text{N}$  enrichments are jointly removed and therefore make no impact on the overall isotopic ratio, provided that three conditions are satisfied (Kirkham & Bartholomew, 1954):

1. Process rates must remain constant over the period of measurement
2. The incubation period must be short enough to avoid remineralization of labelled sources of N which have become immobilized
3. There must be no significant discrimination between isotopic sources of  $^{15}\text{N}$  and  $^{14}\text{N}$  sources by the processes under study

A fourth consideration, which is of particular importance when using intact cores (Davidson *et al.*, 1991) is that the inorganic N pools ( $\text{NH}_4^+$  and  $\text{NO}_3^-$ ) within the soil microplot should be uniformly enriched with the isotope source to avoid erroneous results.


A technical protocol was adopted which to a large extent, satisfied the requirements of the first three conditions and special equipment was designed to fulfill the fourth. Compliance with the first of these conditions was achieved by adopting a relatively short incubation period of 4 d, at a time when weather conditions were stable. The shortened incubation period also enabled problems associated with the second condition to be avoided; Barraclough (1991) has suggested that a period of no more than 3-5 d is suitable for soils at temperatures up to 23 °C. A 4 d incubation was adopted as a compromise to also produce a measurable change in inorganic N to allow accurate and simultaneous 'tube' incubation measurements (for net rates). It was not possible, however, to ensure compliance with the third condition, but this is thought unlikely to present problems when soil enrichments are employed that greatly exceed the natural abundance of  $^{15}\text{N}$  (Davidson *et al.*, 1991). Specially designed apparatus was constructed in order to achieve uniform

enrichment of undisturbed soil and is described below. Process rates were derived from the decline in the initial enrichment of the soil N pool due to the addition of unlabelled 'native'  $^{14}\text{N}$  (ammonification and nitrification). Consumptive processes (immobilization + gaseous losses) were assessed by the rate of removal of the  $^{15}\text{N}$  label, but could also have been affected by addition of the isotope (substrate). Alternative estimates of immobilization were examined, based on the appearance of  $^{15}\text{N}$  in chloroform-labile microbial biomass.

Initial analysis of soils from the experimental sites suggested that inorganic N levels ( $\text{NH}_4^+$  plus  $\text{NO}_3^-$ ) could range up to  $20 \mu\text{g N g}^{-1}$  dry soil, and the N-enriched source would add no more than an equivalent amount of N. Aspects to be considered when deciding on soil enrichment include the need to add sufficient isotope at the start of field incubation to be present in measurable amounts at the end. In practice, this means that an enrichment of  $>0.6 \text{ At.}\%$  (i.e. at least twice natural abundance) is desirable at the end of the incubation. Another constraint is the need to keep the enrichment levels at the start of the incubation within the upper limits of the capabilities of the mass spectrometer (MS) (approximately  $12 \text{ At.}\%$ ) to avoid 'spiking' samples with an unlabelled N source in order to prevent the collector of the MS from saturating. The disturbance of the 'steady state' must also be kept to a minimum to avoid excessive perturbations in the rates of processes to be measured. It is also necessary to make measurements of process rates close to the linear phase of the isotope dilution curve where processes obey zero-order kinetics to maximise accuracy (Barraclough, 1991). Because rates of mineralization in grassland soils are likely to be high, the initial enrichments were increased to  $25 \text{ At.}\%$   $^{15}\text{NH}_4^+$  and to  $20 \text{ At.}\%$   $\text{NO}_3^-$  (since rates of ammonification are most likely  $>$  nitrification; see Section 5.2.3).


## 6.2 Equipment for <sup>15</sup>N enrichment of undisturbed soil

Enclosures for the microplots were fashioned from PVC tubes (175 x 100 mm) with chamfered cutting edges (30°) at one end. Instead of paired microplots, the current technique employed individual microplots which were inserted in the ground (150 mm depth) using a drop-hammer device. A multi-point soil injector (Plate 6.1) was designed which fitted onto the rim of the microplots, based on a modified design of a gas injector described by Murphy *et al.* (1998), but adapted for liquid additions. Injection of the liquid was via a cluster of 13 spinal needles (14 gauge stainless steel, 150 mm length), mounted in brass couplings in a perspex manifold (Fig. 6.1a) and held in place with knurled screw caps. The pointed ends of the needles were sealed with epoxy resin and drilled with two side-port openings. Flow restrictors were placed over the top end of the needles to increase the back pressure within the manifold to help maintain an even flow of liquid through the needles. The flow restrictors consisted of discs (7 mm diameter) cut from an open cellular (permeable) foam material (ear-defender plugs). The manifold and needles were supported on three polished stainless (*ram*) steel rods (16 mm o.d.) which allowed the assembly to move a distance of 150 mm: equivalent to the length of the needles (Plate 6.1). In the primed position, the needles were fully extended and a 60 ml syringe (containing 40 ml liquid) was placed in a cradle and connected to the inlet of the manifold by silicon tubing using Luer-lock fittings. A handle winder, mounted on screw studding (25 mm o.d., *Acme* 5 t.p.i) was synchronised with the movement of the plunger and the travel of the needles through the soil profile. In this way, injection of liquid was always uniformly distributed, regardless of the rate at which the needles were withdrawn through the soil (see Section 6.3.1).




**Plate 6.1**  $^{15}\text{N}$  soil injector

a)



b)



**Figure 6.1 a)** Computer-aided design/manufacture (CAD/CAM) drawing of the perspex manifold from the  $^{15}\text{N}$  soil injector

**b)** Template showing location of hypodermic needles in the manifold and identical pattern used for polypropylene templates: (i) soil surface template with spiked feet (TA), (ii) thicker template used for making perpendicular holes (TB), (iii) needle-guide template (TC)

### 6.2.1 Injection of $^{15}\text{N}$

A polyethylene (100 mm diameter, 10 mm deep) template (TA), which fitted closely inside the microplot, was pressed on to the surface of the soil and kept in place by three spiked feet. It contained 13 countersunk holes in a pattern which was designed to distribute isotope evenly throughout the microplot (Fig. 6.1b). A second polyethylene (100 mm diameter, 40 mm deep) template (TB) was placed on top of TA and contained the same pattern of holes, which were aligned with TA using two metal spikes (one through the centre hole and the other through a peripheral hole) inserted into the soil. A wooden handled spike (190 mm) was used to make guide holes in the soil to a depth of 150 mm via both templates. The thicker template (TB) ensured that the guide holes were perpendicular to the soil surface in order to avoid bending the needles (n.b. this technique is most suited to investigations confined to the upper organic layer of soils where the major proportion of microbial activity occurs and in soil types which contain few stones). The guide spikes and TB were then removed and the injector positioned over the microplot. The needle cluster was aligned with TA by means of a needle-guide template (TC, Fig. 6.2) which could be positioned near to the needle points to enable them to be aligned with the countersunk holes on TA. The needles were then gently introduced into the soil, taking care to present the whole needle cluster evenly, until the base plate of the injector was seated on top of the microplot rim. By rotating the handle winder, the needles were withdrawn steadily through the soil profile whilst the plunger was advanced into the syringe barrel to force liquid (40 mls) from the manifold.

### 6.3 Experimental details

The experimental layout adopted in Section 4.1 was repeated, but for this experiment nine




Figure 6.2  $^{15}\text{N}$  Soil injector

individual plots (30 m<sup>2</sup>) were re-sited to provide new sets of three replicates for each of the three base treatments. As before, the treatments consisted of long-term (>15 years) grazed pastures which were drained by a system of field and mole drains (see Section 3.2). During the grazing season of 1997, animals were excluded from these plots and the swards were maintained in a 3-cut silage system. The three base treatments were: fertilized grass swards receiving 200 kg N ha<sup>-1</sup> y<sup>-1</sup> (*F*); unfertilized swards containing a mixture of clover (approximately 10% by weight) and grass (*C*); unfertilized grass only swards (*Z*). In April 1997, all the plots received P and K (25 and 50 kg ha<sup>-1</sup>, respectively) and the *F* plots also received 40 kg N ha<sup>-1</sup>. The nine plots were cut in June and again in July followed by fertilizer applications of 60 kg N ha<sup>-1</sup> to the *F* plots on both occasions. Thus, by the start of the measurement period, the *F* plots had received a total of 160 kg N ha<sup>-1</sup>, i.e. 80% of the intended annual application.

One week before commencement of measurements (6 August), 54 microplots were inserted in these swards (6 microplots to each of 3 replicate plots x 3 treatments). The microplots were left for one week for the swards and soils to recover from any disturbance and then on 13 August (*t*<sub>0</sub>), herbage from within the microplots was cut level with the rim (25 mm above the soil surface). Each microplot was injected uniformly with a <sup>15</sup>N solution throughout the soil core profile, using the new soil injector. In each of the base treatments, 12 replicate microplots received a solution containing 25 At. % <sup>15</sup>NH<sub>4</sub><sup>+</sup> and the other 6 microplots received 20 At. % <sup>15</sup>NO<sub>3</sub><sup>-</sup> to follow the rates of gross mineralization and nitrification, respectively. After 24 h equilibration (14 August), a soil core was removed (*t*<sub>1</sub>) from the centre of each microplot, using an incubation tube and cutting ring, as described in Section 2.3. The soil remaining in the microplot, which consisted of a

concentric ring of soil with a hollow centre, was removed from the microplot and the inorganic N extracted within 1 h using 2 M KCl. The incubation tube, containing a soil core with the same enrichment as the t1 soil, was placed in an incubation vessel in the field for a further 96 h incubation (without *phytostats* to allow normal plant uptake processes to proceed during the measurement period). Half of the microplots which had received  $^{15}\text{NH}_4^+$  were incubated over gypsum blocks without controlled-release capsules (no inhibitor, A-data set) and the other half (plus inhibitor, I-data set) were incubated with capsules to provide an infusion of  $\text{C}_2\text{H}_2$  into the soil core (see Section 2.3). The microplots receiving  $^{15}\text{NO}_3^-$  were also incubated without  $\text{C}_2\text{H}_2$  (N-data set). As there was little likelihood of leaching from the soil at that time of year (SWC had fallen below field capacity), I-E resin bags were omitted from the incubation vessels and specially constructed gypsum blocks (25 mm height) occupied fully the space beneath the soil cores. On 16 August, the incubation tubes with controlled release capsules (plus inhibitor, I-data set) were removed briefly to enable the gypsum blocks to be re-wetted (10 mls water), as soil conditions were particularly dry. This ensured that the reaction with  $\text{CaC}_2$  proceeded satisfactorily so that the infusion of  $\text{C}_2\text{H}_2$  was maintained.

At the end of the incubation period (18 August), the soil cores were removed (t5) and analysed within 1 h (as above). Over the experimental period (13-18 August), the mean maximum and minimum temperatures were 23.4 and 12.7 °C, respectively. Soil temperature at 100 mm depth averaged 17.3 °C, with a total of 49.2 hours of sunshine. Apart from precipitation on 12 August, when 3.9 mm rain was recorded, the experimental period remained dry and warm during a prolonged spell of settled weather conditions.

### 6.3.1 Testing uniformity of soil $^{15}\text{N}$ injection

An additional 8 microplots were used to test the uniformity of  $^{15}\text{N}$  injection: four received 25 At. %  $^{15}\text{NH}_4^+$  and four received 20 At. %  $^{15}\text{NO}_3^-$ , as described previously. After 24 h equilibration period (t1, 14 August), cores were removed from the centres of the microplots, using incubation tubes. The soil from the tubes, together with the soil from the associated outer cores, were extracted immediately with KCl to compare the uniformity of enrichments achieved by  $^{15}\text{N}$  injection at the starting point of an incubation period, i.e. 'start values' from the outer core and the corresponding 'start values' for soil cores undergoing a period of incubation.

### 6.4 Soil/plant processing

On each day of sampling (t1 and t5), soil was removed from microplots and incubation tubes and passed through a 6 mm mesh sieve. Stones were discarded, but all plant material (shoots plus roots) which was retained on the sieve was carefully removed, washed clean of soil over a 200  $\mu\text{m}$  mesh sieve and oven dried at 105 °C. The dried plant material was then weighed and ground to a fine consistency using a succession of grinders: hammer mill (JKA, Model A20) and ball mill (Glen Creston). Soil sub-samples (50 g fresh) were extracted with 250 ml of 2 M KCl for analysis of  $\text{NH}_4^+$  and  $\text{NO}_3^-$  (Scalar autoanalyser) and the extracts were frozen on the day of collection (t1 and t5). Other soil samples (100 g fresh) were used to assess the gravimetric water content, or extracted (12 g fresh) in 0.5 M  $\text{K}_2\text{SO}_4$  for analysis of the *SMB* (as described in Section 5.2.2).

### 6.5 Diffusion of $^{15}\text{N}$ labelled solutions

The inorganic N enriched with  $^{15}\text{N}$  was recovered from soil extracts using a diffusion

technique described by Stark and Hart (1996). A VG 602E isotope ratio mass spectrometer (see Fig. 6.3), linked to a Carlo Erba C N analyser and controlled by Europa ANCA software/interface, was used to measure the  $^{15}\text{N}$  enrichment of the soil extracts and required at least 100  $\mu\text{g N}$  to ensure a robust signal. Sufficient KCl extract to contain either 140 or 200  $\mu\text{g N}$  (initial tests suggested recoveries of 70% and 50% for  $\text{NH}_4^+$  and  $\text{NO}_3^-$  contents, respectively) were placed into 250 ml wide-neck polyethylene bottles. A pair of acidified traps was added to each bottle, consisting of two discs of filter paper (Whatman No. 1, 7 mm diameter), each treated with 5  $\mu\text{l}$  2.5 M  $\text{KHSO}_4$ . The pair of discs were sealed in a 70 mm strip (4 mm apart) of 12.5 mm wide PTFE pipe sealing tape (Teflon) using an 11 mm diameter plastic culture tube to press a concentric circle around each disc. In the case of  $^{15}\text{NH}_4^+$  labelled N, the extracts were diffused for 6 d at 25 °C with 0.3 g of MgO, whereas  $^{15}\text{NO}_3^-$  labelled N was first diffused for 7 d in open bottles (without lids) which were swirled daily to drive off  $\text{NH}_3$ . After this time, water was added to replace evaporative losses and an extra scoop (0.3g) of MgO plus 0.5 g of Devarda's alloy was added to the extract; the bottles were sealed tightly with a screw lid and left for a further 6 d to diffuse. Devarda's alloy releases  $\text{H}_2$ , in addition to the  $\text{NH}_3$  liberated by MgO and leakage from the screw cap was avoided by inverting the bottles during the diffusion process so that the gases were entrapped in the closed end of the bottles. Bottles which do leak can be identified readily by seepage of liquid and therefore discounted. Soil microbial biomass (*SMB*) was extracted from the soils by the fumigation-extraction (F-E) method (Brookes *et al.*, 1985; Vance *et al.*, 1987). The extracts were used to determine biomass C, by dichromate oxidation with back-titration against ammonium ferrous sulphate, and biomass N by Kjeldahl digestion, followed by flow injection analysis.




Figure 6.3 Schematic diagram showing a C N analyser linked to an isotope ratio mass spectrometer.

Biomass C and N were calculated using extraction correction factors of 2.64 and 2.22, respectively (see Section 5.2.2). Analysis of the Kjeldahl digests (containing 140 µg N) for  $^{15}\text{N}$  enrichment used the same procedure as for KCl extracts, but in this case the solutions in the diffusion bottles were made sufficiently alkaline (pH >11.0) to liberate  $^{15}\text{NH}_3$ , by the addition of 10 M NaOH, instead of MgO.

After diffusion, the PTFE strips were recovered, rinsed in deionised water and the top layer of PTFE was peeled back with forceps to expose the filter paper discs. Each pair of discs was impaled on a single stainless steel pin and dried in a desiccator over concentrated  $\text{H}_2\text{SO}_4$  for at least 4 h. The pairs of dried discs were placed together in one tin capsule (8 x 5 mm) and analysed for  $^{15}\text{N}$  enrichment by mass spectrometry. The two discs provided a safeguard against failure or incomplete diffusion by one of the discs, since the other would still have had sufficient capacity to absorb all of the labelled N source.

#### 6.5.1 Standards and controls

Additional extracts (75 ml) of 2 M KCl containing 100 µg N (as  $^{15}\text{NH}_4^+$ ) were placed in bottles with acid traps (diffused standards) covering a range of standards: 0, 2.5, 5.0, 7.5 and 10.0 At. %  $^{15}\text{N}$  (see Appendix 1 for calculations). Other acid traps were first 'spiked' with  $^{15}\text{N}$  in the same range and analysed in tin capsules (non-diffused standards). The bottles were incubated, as above, with 0.3 g MgO. Blank-correction of the samples was made using the standard controls to determine the extent of contamination, from the decline in enrichments between non-diffused and diffused standards. The effectiveness of the first stage of the diffusion process was also examined (i.e. liberation of all  $^{15}\text{NH}_4^+$ ), by adding 100 µg N (as 5 At. %  $^{15}\text{NH}_4^+$ ) to four replicate bottles, plus 100 µg N (as 0.3663

At. %  $^{15}\text{NO}_3^-$  viz. natural abundance). Four further replicates received only 100  $\mu\text{g N}$  as 0.3663 At. %  $^{15}\text{NO}_3^-$ . The first set was diffused with MgO (in bottles vented to remove  $\text{NH}_3$ ) and then both sets (8 bottles) were diffused again, this time with Devarda's alloy (+ extra MgO). Acid traps were only added at the second stage to capture any labelled N sources remaining in the bottles.

#### 6.6 Other measurements

C-mineralization was estimated from the amount of  $\text{CO}_2$  respired from field incubated soil cores according to a modified technique described by Gilmour *et al.* (1985). At  $t_1$ , pairs of fresh soil cores (25 x 150 mm) were removed from each of the three replicate plots and each core was placed in an individual 11 Kilner jar (6 replicate jars per treatment) containing a 70 ml serum bottle with 25 ml 0.5 M NaOH. Six other jars without soil, but containing alkali were included as controls. The jars were placed in holes in the ground for 96 h, after which the bottle of alkali was removed and stoppered ( $t_5$ ). The amount of  $\text{CO}_2$  respired was calculated after addition of 10%  $\text{BaCl}_2$  and back titration of the excess alkali against 0.1 M HCl (see Appendix 2).

Net N-mineralization was estimated using the technique described by Hatch *et al.*, 1990. Four soil cores (37 mm diameter) were analysed from each replicate plot for  $\text{NH}_4^+$ -N content at the start ( $t_0$ ) of the measurement period (see Section 4.2.1). Four additional cores were taken from the same plots and placed in 11 Kilner jars (3 replicate jars to each treatment) with 2%  $\text{C}_2\text{H}_2$  in the headspace for an incubation lasting 7 d. At  $t_7$ , the cores were removed and analysed, as for the  $t_0$  cores. Net mineralization was calculated from the difference in  $\text{NH}_4^+$ -N contents between the 'start' and incubated cores ( $t_7 - t_0$ ) and

expressed as a net rate per day.

## 6.7 Results and discussion

Each needle in the  $^{15}\text{N}$  soil injector delivered on average 2.99 ml with a precision of  $\pm 0.275$  ml, which was estimated to have increased the soil water content of the microplots by not more than 5% (dry weight soil basis). In the test microplots, no significant differences were found between the inorganic N pool sizes of either  $\text{NH}_4^+$  or  $\text{NO}_3^-$ , nor in their  $^{15}\text{N}$  enrichments (24 h after injection, t1), when the inner soil cores (incubation tubes) were compared with the associated outer cores (extracted on the same day, t1). Thus, the comparison between cores extracted at t1 and those incubated until t5 was valid in terms of the uniformity of initial  $^{15}\text{N}$  enrichments. Diffusion efficiency was subjected to the severest test under laboratory conditions (see Section 6.5.1), since any carry-over of labelled  $^{15}\text{NH}_4^+$  would be readily identified as an elevated enrichment in the  $\text{NO}_3^-$  component (added at natural abundance) which was also reduced to  $\text{NH}_4^+$  before diffusion. The mean enrichment (At. %  $^{15}\text{N}$ ) of controls without  $^{15}\text{N}$  addition was  $0.389 \pm 0.0016$ , compared with  $0.378 \pm 0.0009$  in the controls to which  $^{15}\text{NH}_4^+$  was added. The difference was significant ( $P < 0.001$ ) and probably arose through contamination ( $^{14}\text{N}$ ) in the extra MgO used in the venting stage, which would have the effect of diluting the  $^{15}\text{N}$  natural abundance ratio. Since there was no evidence, therefore of an elevated enrichment attributable to carry-over of  $^{15}\text{NH}_4^+$ , it was concluded that the diffusion stage was effective in distinguishing between the two forms of inorganic N. Similarly, where other amendments were equal (*viz.* MgO and Devarda's alloy) the apparent contamination was insignificant, so that the correlation between standards (diffused vs non diffused) was extremely close ( $r^2 = 0.9994$ ) with an intercept for the fitted line of 0.3772 At. %  $^{15}\text{N}$ ,

which is close to natural abundance. Standards (diffused and non diffused) were included in all subsequent analytical runs to estimate the extent of any contamination. However, no correction for blank contamination was found to be necessary with enrichments which greatly exceeded natural abundance, or in sample runs where close correlations were established between diffused and non diffused standards (i.e. with uniformly low levels of contamination). The calculation required for blank-correction of low enrichment samples (Stark and Hart, 1996) is, nevertheless, reproduced in Appendix 3 for reference.

In the experimental microplots, prior to injection of  $^{15}\text{N}$  (t0), the inorganic N contents of the soils were about  $5 \mu\text{g NH}_4^+\text{-N g}^{-1}$  soil in both *F* and *C* treatments (less in *Z*) and also of  $\text{NO}_3^-$  in *F*, but with approximately one-tenth of these concentrations ( $0.5 \mu\text{g N g}^{-1}$  soil) of  $\text{NO}_3^-$  in *C* and *Z* treatments (Table 6.1).

**Table 6.1** Inorganic N contents ( $\mu\text{g N g}^{-1}$  dry soil  $\pm$ SEM) of  $\text{NH}_4^+\text{-N}$  pool (A,I) or  $\text{NO}_3^-\text{-N}$  pool (N), before injection of  $20 \mu\text{g N g}^{-1}$  dry soil (t0) and at 24 h (t1) and 96 h (t5) after injection (n = 6)

Treatments (N pool)	t0	t1	t5
FA ( $\text{NH}_4^+\text{-N}$ )	5.9 (0.52)	26.2 (1.78)	34.7 (5.06)
FN ( $\text{NO}_3^-\text{-N}$ )	5.9 (2.44)	41.2 (9.70)	43.4 (9.14)
FI ( $\text{NH}_4^+\text{-N}$ )	5.9 (0.52)	28.6 (2.66)	35.3 (2.73)
CA ( $\text{NH}_4^+\text{-N}$ )	4.5 (0.31)	24.1 (1.58)	21.8 (2.11)
CN ( $\text{NO}_3^-\text{-N}$ )	0.5 (0.15)	28.2 (2.20)	26.5 (2.13)
CI ( $\text{NH}_4^+\text{-N}$ )	4.5 (0.31)	26.7 (2.21)	32.3 (2.53)
ZA ( $\text{NH}_4^+\text{-N}$ )	3.7 (0.44)	24.4 (2.21)	36.7 (8.92)
ZN ( $\text{NO}_3^-\text{-N}$ )	0.3 (0.07)	21.2 (0.58)	22.8 (1.26)
ZI ( $\text{NH}_4^+\text{-N}$ )	3.7 (0.44)	20.6 (1.04)	26.7 (5.65)

At t1, following injection of  $^{15}\text{N}$ , all treatments increased ( $P < 0.001$ ) by approximately  $20 \mu\text{g N g}^{-1}$  soil, as intended (the exception was *FN*, which probably reflects greater heterogeneity due to previous N inputs) and these concentrations persisted until t5 (Table 6.1). The  $^{15}\text{N}$  enrichments of the soils at t1 were broadly similar across all treatments (range 10.8-16.0, Table 6.2), but had declined by about one-third at t5. Consequently, of the total  $^{15}\text{N}$  recovered from all treatments at t5 (Table 6.2), the soil retained the major component (22-61%) with more variable amounts recovered in the *SMB* (0-23%); the minor component was in plant material (3-11%). Of the  $^{15}\text{N}$  added, therefore, on average 68%, 56% and 72% in total was recovered at t5 in *F*, *C* and *Z* treatments, respectively. These values are in line with total recoveries of added  $^{15}\text{N}$  in other studies (Bristow *et al.*, 1987; Schimel *et al.*, 1989), but are at variance in terms of the dominance of the *SMB* competition over plant biomass. Plant biomass was found to be the more important sink for added N by Bristow *et al.* (1987) and Ledgard *et al.* (1998). However, both the present study and other workers (Jackson *et al.*, 1989) found microbial uptake to be the dominant factor in removal of added N, and Hart *et al.* (1993) also found similar recoveries of added  $^{15}\text{N}$  in the *SMB*, which ranged from 9-15%. Recovery of  $^{15}\text{N}$  by plants (shoots + roots) was similar in all three treatments and increased by about 50% between t1 and t5 (Table 6.2). Recoveries of  $^{15}\text{N}$  in the *SMB* were far more variable and in some treatments declined with time, illustrating the transient nature of recently acquired microbial N (Bristow *et al.*, 1987). However, potential sources of error by initial, rapid abiotic reactions (Davidson *et al.*, 1991) were largely avoided by delaying the first sampling for an equilibration period of 24 h, following  $^{15}\text{N}$  injection. The competitive success of the *SMB* for available N could represent a major control over the availability of N for plant uptake and will be discussed in more detail later.

**Table 6.2** Mean (n = 6) percentage recovery of  $^{15}\text{N}$  from the  $\text{NH}_4^+$  pool (A-and I-data sets) and the  $\text{NO}_3^-$  pool (N-data sets)

Treat.	At. % $^{15}\text{N}$ soil (inorganic)		%rec. $^{15}\text{N}$ soil (inorganic)		%rec. $^{15}\text{N}$ ( <i>SMB</i> )		%rec. $^{15}\text{N}$ (plant)		Total % recovery	
	t1	t5	t1	t5	t1	t5	t1	t5	t1	t5
FA	14.49	9.02	64.0	50.3	5.2	11.7	2.7	4.8	72	67
FN	10.79	9.41	64.6	61.1	0.3	0	1.4	3.0	66	64
FI	15.07	8.81	71.5	51.9	5.2	15.2	2.9	4.8	80	72
CA	14.86	6.49	60.1	22.3	18.3	10.8	5.0	7.8	83	41
CN	13.46	11.24	63.2	49.6	12.6	0	3.9	8.8	80	58
CI	14.49	8.50	64.8	44.9	25.1	18.7	3.4	5.2	93	69
ZA	13.73	8.55	55.7	54.4	15.8	19.5	5.1	7.9	77	82
ZN	16.02	13.44	57.2	51.3	0.1	0	5.1	7.3	62	59
ZI	13.86	8.98	48.1	42.2	45.1	22.8	5.7	10.9	99	76

Process rates (expressed on a per day basis) were calculated from the rates of dilution, or removal of the  $^{15}\text{N}$  labelled  $\text{NH}_4^+$  and  $\text{NO}_3^-$  between days 1 and 5 (t5 - t1), following the equations (see Appendix 4) of Kirkham and Bartholomew (1954). Although the lowest values for gross mineralization were found with the Z treatments (Table 6.3), no significant differences were found between the treatments. Conversely, the lowest immobilization of  $\text{NH}_4^+$  was found with the F treatments and this trend was repeated in the immobilization values found in the chloroform-labile pool of the *SMB*, but the data were highly variable and no significant differences were established. It was interesting to note, however, that within each treatment the presence of the nitrification inhibitor ( $\text{C}_2\text{H}_2$ ) increased the rate of immobilization of  $^{15}\text{NH}_4^+$  into the *SMB* by 30%, 48% and 9% for F, C and Z (I-data sets), respectively. This suggests that the treatments with the highest nitrifying activities (see Table 6.4) were most affected by inhibition.

**Table 6.3** Gross rates of processes ( $\pm$ SEM) obtained from  $^{15}\text{N}$  mean pool dilution data ( $\mu\text{g N g}^{-1} \text{d}^{-1}$ )  $n = 6$

Treatment	Gross mineralization	Ammonium consumption	Gross nitrification	#Ammonium immobilization	SMB immob.
FA	4.1 (1.41)	2.8 (1.26)	1.6 (0.45)	1.2	3.0
FI	4.2 (0.73)	2.5 (0.79)	"	0.9	3.9
CA	5.4 (1.14)	6.0 (1.21)	1.2 (0.33)	4.8	3.3
CI	4.1 (0.63)	2.7 (0.48)	"	1.5	4.9
ZA	3.6 (0.55)	4.0 (0.54)	1.0 (0.06)	3.0	5.5
ZI	2.5 (0.65)	4.3 (1.06)	"	3.3	6.0

#Calculated from Davidson *et al.*, 1991:  $(\text{NH}_4^+$  consumption - gross nitrification =  $\text{NH}_4^+$  immobilization) where  $\text{NH}_4^+$  consumption includes the sum of immobilization (i.e. microbial assimilation), autotrophic nitrification, volatilization and other possible fates.

Gross rates of mineralization and  $\text{NH}_4^+$  consumption (Table 6.3) were similar between all treatments and close to the soil inorganic pool concentrations prior to injection (Table 6.1). Thus, inherently low inorganic N levels would be expected to be maintained in the 'steady state', unless amended by fertilizer inputs. A useful indication of the relative turnover rates in these treatments can be obtained from calculations of **pseudo-residence times**, defined as "the ratio of the  $\text{NH}_4^+$ -N pool size :  $\text{NH}_4^+$ -N consumption rate" (Tietema and Wessel, 1992). Ratios of 2.1, 0.8 and 0.9 ( $\text{NH}_4^+$ -N pool) and 3.9, 0.3 and 0.4 ( $\text{NO}_3^-$ -N pool) were found with *F*, *C* and *Z* treatments, respectively, representing the number of times the pool size turned over each day. Therefore, turnover of N-pools was faster in the N-limited swards (*C* and *Z*), with the  $\text{NO}_3^-$ -N pools turning over twice as fast as the  $\text{NH}_4^+$ -N pools. An explanation might be that plants experiencing only a limited mobile N supply, become effective 'scavengers' of sources of available N (Parsons *et al.*, 1991). In the fertilized swards (*F*), the turnover of the  $\text{NH}_4^+$ -N pool was on a 2-day cycle, whereas the  $\text{NO}_3^-$ -N pool took twice this time to be processed. It is clear then, that high fluxes and a rapid

turnover of the inorganic N pools are features of these grazed grassland soils, as was previously demonstrated by Ledgard *et al.* (1998).

**Table 6.4** Nitrification and consumption of nitrate ( $\mu\text{g N g}^{-1} \text{d}^{-1} \pm \text{SEM}$ )  $n = 6$

Treatment	Nitrification (gross)	Nitrification (Tube)	Nitrate consumption
F	1.6 (0.45)	0.3 (1.47)	1.5 (0.42)
C	1.2 (0.33)	2.6 (1.00)	1.6 (0.47)
Z	1.0 (0.06)	-2.5 (2.32)	0.8 (0.16)

Gross nitrification was also closely matched with  $\text{NO}_3^-$  consumption (Table 6.4) so that, again,  $\text{NO}_3^-$  would be unlikely to accumulate unless supplemented by fertilizer. The highest rate of nitrification (gross) was found with *F* compared with *Z* treatments, but this was only at a low level of significance ( $P = 0.09$ ) and with *C* intermediate (Table 6.4). Measurements of net nitrification (by tube incubation) were not consistent with other estimates (Table 6.4) and may have been adversely affected by additions of N (as  $^{15}\text{NH}_4^+$ ), since the assessment relied on differences in  $\text{NH}_4^+$  produced in the presence and absence of a nitrification inhibitor ( $\text{C}_2\text{H}_2$ ). Nitrification rates are dependent upon the supply of substrate ( $\text{NH}_4^+$ ) usually from the mineralization of soil O.M. (i.e. gross mineralization). In the present study, 39, 25 and 33% of the  $\text{NH}_4^+$  mineralized was nitrified (based on gross nitrification rates) in *F*, *C* and *Z* treatments, respectively, compared with 72% found with the same soil type in the previous autumn/winter experiment (see Section 5.3.2). Similar rates (12 - 46%) were measured during the growing season of annual grasses in California (Davidson *et al.*, 1990). Thus, the oxidizing capacity of the nitrifiers was exceeded in the summer conditions of the present study, which suggests that the  $Q_{10}$  of nitrifiers must be

somewhat lower than that of heterotrophs involved in O.M. decomposition.

Combining the tube incubation technique with the  $^{15}\text{N}$  technique enabled simultaneous assessments of gross and net rates. Net rates of mineralization (Table 6.5) obtained with tube, jar, or  $^{15}\text{N}$ -calculated rates were in reasonable agreement (particularly with *F* and *C* treatments) and approximately half the gross rates found for each of the treatments (Table 6.3). Estimates of net mineralization in the *C* treatment were intermediate between *F* and *Z*, but there were no significant differences overall between treatments. The lowest microbial activity, as measured by  $\text{CO}_2$  respiration (Table 6.5), also appeared to be in *Z*, but again was not significantly different from the other treatments (it should be noted that the simplified method, using intact soil cores, also included respiration by living roots).

**Table 6.5** Net mineralization of nitrogen ( $\mu\text{g N g}^{-1} \text{d}^{-1} \pm \text{SEM}$ ) estimated by three different techniques, and carbon ( $\mu\text{g C g}^{-1} \text{d}^{-1}$ ) measured by  $\text{CO}_2$  evolved during jar incubation

Treatment	#Net N min. (calculated)	Net N min. (Tube: n=6)	Net N min. (Jar: n=3)	Net C min. (Jar: n=6)
F	2.9	2.3 (0.66)	1.9 (0.32)	70.5 (6.73)
C	0.6	2.1 (0.84)	1.8 (0.13)	76.0 (5.62)
Z	0.6	0.6 (1.80)	1.3 (0.28)	59.9 (5.28)

#Calculated from the difference between gross min. and  $\text{NH}_4^+$  immobilization (using A-data sets)

Additionally, laboratory incubations (Table 6.6) supported the finding that microbial activity, as assessed by specific respiration, was similar in all treatments. In terms of biomass size, both biomass C and N were similar in treatments *F* and *Z* (long-term undisturbed for >50 years), but lower in *C* ( $P < 0.05$ ), which had been ploughed and reseeded in 1982 and had, therefore, not yet reached a new equilibrium.

Plant N uptake data (calculated from  $^{15}\text{N}$  recoveries) were very variable ( $cv = 60.3\%$ ), with only low rates recorded which averaged  $0.21$ ,  $0.44$  and  $0.51 \text{ kg N ha}^{-1} \text{ d}^{-1}$  for *F*, *C* and *Z*, respectively. This is in the reverse order of what might have been predicted from the known productivity of these swards, but may have been an anomaly of recently cut plants and a rapid response to applied N in those swards containing N-starved plants (viz. *C* and *Z*), compared with the lower uptake ( $P < 0.01$ ) from higher N-status plants (*F*). These results are, therefore to be considered with some reservations and may not be wholly representative of the swards in equilibrium with the management regimes. There was no effect of either the form of N ( $\text{NH}_4^+$  or  $\text{NO}_3^-$ ) or inhibitor ( $\text{C}_2\text{H}_2$ ) on the relative uptake rates within (A-, N-, or I-data sets) or between treatments (*F*, *C* or *Z*).

**Table 6.6** Soil microbial biomass N and C ( $\mu\text{g g}^{-1}$  dry soil) and specific respiration ( $\mu\text{g CO}_2\text{-C g}^{-1}$  dry soil)  $n = 6$


Treatment	Biomass N		Biomass C		Specific respiration
	t1	t5	t1	t5	
F	278.6 <sup>a</sup>	294.9 <sup>a</sup>	2025 <sup>a</sup>	2024 <sup>a</sup>	1.39 <sup>a</sup>
C	242.3 <sup>a</sup>	244.3 <sup>b</sup>	1415 <sup>b</sup>	1459 <sup>b</sup>	1.14 <sup>a</sup>
Z	282.9 <sup>a</sup>	292.3 <sup>a</sup>	1943 <sup>a</sup>	1989 <sup>a</sup>	1.13 <sup>a</sup>

Values which have different letters within columns are significantly different ( $P < 0.05$ )

Soil microbial biomass (*SMB*) data for immobilization were also very variable ( $cv = 93.1\%$ ), with rates which were lowest in *F* and highest in *Z* and with *C* intermediate, but the differences were not significant (Table 6.3). Some caution must be exercised in the interpretation of immobilization from the rate of appearance of the label in the microbial biomass. Whilst this approach may seem more direct than calculations based on the disappearance of the label from the inorganic N pool (see Appendix 4, equation 2),

questions have been raised concerning uncertainties with the measurement of *SMB* by chloroform fumigation (Voroney and Paul, 1984; Vance *et al.*, 1987; Davidson *et al.*, 1989). No adjustment was made to allow for the fraction of microbial biomass N extracted after exposure to chloroform ( $K_N$  factor), since an appropriate value for  $K_N$  has not yet been widely adopted. An alternative method (see Table 6.3) of calculating ammonium immobilization and net mineralization (by difference from gross mineralization, see Table 6.5) showed that differences in net mineralization are largely determined by rates of immobilization (Fig. 6.4).

The effect of nitrification on rates of N cycling processes was examined in swards with different nitrifying activities and compared with rates measured in the presence of a nitrification inhibitor. Where nitrification was not inhibited (Fig. 6.4a), the lowest rates in net mineralization were associated with soils from the unfertilized swards (C and Z). Ledgard *et al.* (1998) also found that immobilization was greatest in unfertilized swards, which in turn determined the ultimate differences found in net mineralization rates. Where nitrification was inhibited (Fig. 6.4b), net mineralization was not significantly different between the three soils (F, C and Z). The calculated rates for net mineralization ( $\mu\text{g N g}^{-1} \text{d}^{-1}$ ) shown in Fig. 6.4b were  $3.3 (\pm 0.92)$ ,  $2.6 (\pm 0.97)$  and  $-0.8 (\pm 0.97)$ , respectively and were in closer agreement overall (for F and C) with estimates by 'tube' and 'jar' incubation, than those based on calculations without the inhibitor (Table 6.5). When nitrification is allowed to proceed naturally, other losses may occur (see Section 3.4) which may cause an under-estimate in measured rates. In the present study, the effect of inhibitor (I-data set) was to produce higher values for immobilization of  $^{15}\text{N}$  in the *SMB* in all treatments, suggesting that  $\text{NH}_4^+$  uptake may have been enhanced when this form predominated.



**Figure 6.4** Gross mineralization,  $\text{NH}_4^+$  immobilization and net mineralization (by difference) from  $^{15}\text{N}$  data ( $\mu\text{g N g}^{-1} \text{d}^{-1}$ ) for fertilized (F), grass/clover (C) and unfertilized (Z) swards: **a)** A-data set **b)** I-data set (+ nitrification inhibitor)

### 6.7.1 Summary

The multi-point soil injector proved to be very effective at delivering  $^{15}\text{N}$  evenly throughout the soil profile under investigation, so that accurate measurements of process rates could be obtained. Future use of the apparatus will probably be confined to using separate pairs of microplots (i.e. not concentrically paired), since the combination of tube incubation with  $^{15}\text{N}$  techniques was very labour intensive, although justified in terms of verifying techniques. The importance of nitrification to N cycling in grassland soils was established in net terms in the previous chapter and examined in both net and gross terms in this chapter, which provided confirmation of rate measurements. The wider and more subtle role of nitrification in determining the uptake patterns of plant and/or microbial biomass was not resolved completely, in as much as some of the trends were not statistically established and the differences in availability of forms of N were not reflected in plant utilization. These aspects and recommendations for future work, together with an overview of the whole project are discussed in greater detail in Chapter 7.

## CHAPTER 7 Overview: project synthesis and conclusions

### 7.1 Introduction

An overview of the project is discussed in two parts:

- 1) A synthesis of the data, drawing upon salient points to illustrate the main findings, together with a re-appraisal of the major aims of the project
- 2) A synopsis of the work, identifying the limitations, implications and priorities for future investigations, culminating in a re-examination of the validity of the hypothesis, which proposed that:

**'Nitrification occupies a pivotal role in the N cycle of productive grassland soils by determining the supply of substrate (nitrate) to the two major processes involved in losses of N, i.e. denitrification and leaching'**

### 7.2 Synthesis of the data

A first step in studying nitrification was to examine soils for differences in their capacity to mineralize soil organic matter and release ammonium, which is the substrate for nitrifying organisms. However, a range of laboratory techniques for evaluating possible differences in mineralization (see Section 4.2.4) and *SMB* size (see Section 5.2.2) and activity (see Section 4.3) in the present soils did not show any differences in potential rates in these processes, even between soils with contrasting management histories. It may well be that such differences can only really be detected in truly dynamic situations under realistic conditions for field evaluation. Thus Gill *et al.* (1995) found that net mineralization was greatest in the fertilized treatments from these soils, but that

withholding N fertilizer did not have any immediate effect on rates of N-turnover. The current study, using the improved field incubation technique, gave similar results (see Section 3.2.2), but also illustrated the difficulties of making measurements in soils with such high spatial variability. In the first phase of the project, improvements in the sampling procedures were devised and introduced (see Section 3.3) to decrease some of the experimental errors; sampling replication was increased to try to overcome some of the variation within treatments. Laboratory assessments were done on bulked samples, derived from mixtures of several soil core samples, which helped to decrease the field sampling errors, but this may have stimulated mineralization. Nevertheless, the usefulness of laboratory estimates of potential mineralization as indicators of net rates in the field was limited, since much of the readily degradable organic material is derived from the litter of the current and past seasonal growth; this will not necessarily feature as a source of substrate in laboratory incubation studies. This makes the prediction of N supply particularly difficult. However, a first step towards a predictive tool has come from re-examination of previously published data in an associated study, which has resulted in the identification of some important correlations between soil thermal units (cumulative temperature at 10 cm depth) and net mineralization (Clough *et al.*, 1998).

Aspects of the microbial component of the soils under investigation were examined by standard techniques (see Section 4.3), but appeared to give only incomplete information about the decay process. Lovell *et al.* (1995) found that total biomass was inversely related to net mineralization and Bardgett and Leemans (1995) showed that activity decreased with lower fertilizer inputs, suggesting that much of the biomass may be present in a state of dormancy. Qualitative differences in microbial composition are more difficult

to identify since changes in the activities of the various population components may be only transitory. However, the dominance of fungal over bacterial activity in the fertilized treatment (see Section 4.3.3) is in accordance with the observation that an heterotrophic nitrifying component (resistant to an autotrophic blocking agent) may have been present (see Section 4.4.1). The possibility of an altered balance in the microbial composition of fertilized soils has interesting implications, the investigation of which is probably best undertaken by  $^{15}\text{N}$ -labelling techniques in combination with selective metabolic inhibitors. Such investigations, however, need to be at a microbiological scale and were beyond the scope of the present study.

More sensitive assays of nitrifying activities were used to examine the long-term field sites (see Sections 4.4 and 4.5) and clear differences were established between the soils from unfertilized and fertilized grass swards, with soils from the grass/clover swards being intermediate. Given that these differences were shown to have persisted for one year following removal of cattle and fertilizer inputs, elevated potentials for nitrification suggested increases in the potential for losses of N through denitrification and leaching (Hatch *et al.*, 1996). Interestingly, the contrasting fertilizer histories of these soils had not altered the ability of the nitrifiers to oxidize the potentially toxic nitrite ion, as demonstrated by enzyme kinetics assays (see Section 4.5.4). An important insight was provided into the ability of the nitrifying organisms to convert ammonium into the more mobile (and harmless) nitrate ion. 'Nitrification is not necessarily detrimental to ecosystems if it proceeds to the final product ( $\text{NO}_3^-$ ) and may also enhance plant uptake', as will be discussed later.

It may be that the response of mobilization and immobilization turnover (*MIT*) rates to additional inputs from SOM and the effects of environmental fluctuations are extremely transient. In fact, N-supply plays only a supporting role to the processes of *MIT* and it is the availability of readily oxidizable carbon (C) which is often the rate-limiting step, Lovell *et al.* (1996) found that the addition of N to dung (with a C : N ratio of 19.1 : 1) mixed with soil had no effect on either the size, or activity, of the *SMB*. Therefore, even though rates of supply of  $\text{NH}_4^+$  from the decomposition of SOM may not apparently differ, other reactions influencing its oxidation to  $\text{NO}_3^-$  (*viz.* nitrification) are inherently important in determining the fate of N. It was with some justification that Both *et al.* (1992) concluded that "no simple answer can be given to the question as to which of the community (nitrifying) parameters determined in (their) study is most suitable to describe or predict nitrate production in the field." They went on to suggest that "this information can probably best be obtained from intact cores incubated *in situ*."

### 7.3 Re-appraisal of the main aims

The approach of using field incubated soil cores was adopted in the present study to address the above question posed by Both *et al.* (1992), and to meet the criteria set out in the original aims of the current project which are reproduced here:

- a) To **develop** a new technique for the field measurement of nitrification, with simultaneous estimates of losses of nitrate by denitrification and leaching
- b) To **link** the *in situ* measurements of nitrification with other processes, to gain an understanding of the controlling factors determining the fate of nitrate
- c) To **examine** the practical implications of these findings under different grassland management regimes

Each of the three components above will be reviewed in turn to indicate the extent to which the project has fulfilled the three broad aims. Attention will also be given to the assessment of the implications of the findings and the requirement for future work, where aspects remain unclear or unresolved.

### 7.3.1 Development of a new field incubation technique

The information which was obtained from laboratory-based investigations represented only "snapshots" of particular conditions. The second phase of the project was, therefore, directed towards elucidating the relationships between the process of nitrification *per se* and other transformations of N, using the novel incubation system under field conditions. This new system offers significant improvements over existing methods such as 'jar incubation' (Hatch *et al.*, 1990), 'buried bag' (Eno, 1960) or 'covered tubes' (Raison *et al.*, 1987) and provided simultaneous measurements of the major N cycling processes in a combined approach which had not been attempted previously. The considerable difficulties of measuring changes in mineral N contents from the differences between incubated and non-incubated soil in previously grazed swards (affected by excretal returns) or cut swards (receiving slurry N) had hitherto been avoided assiduously by most other workers. The development of the novel incubation system provided a sampling protocol which enabled closely matching pairs of soil samples to be taken (using concentrically placed soil corers) that overcame many of the problems associated with heterogeneous areas (Hatch *et al.*, 1998). The programme of work included a field incubation study of fertilized/cut grass during spring (see Section 5.2), which was targeted at conditions where potentially high denitrification losses may be coupled to nitrifying activity (Lovell and Hatch, 1997). In autumn 1996 (see Section 5.3), these measurements were repeated on fertilized/cut swards,

following the return of the soil to field capacity, to assess interactions with potential leaching losses. Both experiments showed considerable potential for N loss, particularly through denitrification, and identified possible inaccuracies with some of the existing techniques. In particular, the much higher rates of denitrification found with tube incubation (>3-fold) compared with jar incubation, illustrated how vulnerable well-structured soils can be to disturbance (Hatch *et al.*, 1998) and the likely shortfall in the assessment associated with the latter technique.

The successful development of an effective means of supplying  $C_2H_2$  (a powerful nitrification inhibitor) using controlled-release capsules was another key step in resolving field incubation in open vessels. Whilst  $C_2H_2$  has been used successfully in other studies to assess net mineralization (Hatch *et al.*, 1990; Gill *et al.*, 1995) and denitrification (Ryden *et al.*, 1987), other methods have not utilized the opportunity to resolve several process rates simultaneously by paired incubations ( $\pm C_2H_2$ ). Further, where the scope of investigations can be focused on mineralization (known to be a key process in determining losses, see Blanton, 1991; Jarvis *et al.*, 1996), a simplified version of the tube incubation has been developed (Bhogal *et al.*, in press), which can also be adapted for use in arable soils. The method involves taking account of soil N (before and at the end of the growth period), crop uptake (herbage only), mineralization from soil cores and leaching using ion-exchange resin bags. Possible drawbacks to the use of tube incubation are the need to make contiguous measurements with short intervals between incubations (<14 d) and the difficulties encountered with soil coring in dry or stony soils. Also, the technique does not discriminate between losses of  $N_2O$  from denitrification and those attributable to nitrification, although fully recognised as a significant source under certain conditions (see

Section 1.6). The investigation of the origins of N<sub>2</sub>O (and NO) was beyond the scope of the present study. For an account of current work on the contributions from these major soil processes, see Jarvis (1997).

### 7.3.2 Linking *in situ* nitrification measurements with other processes

Close coupling between nitrification and denitrification has been demonstrated in the present study (Hatch *et al.*, 1998) and this has contributed towards our understanding of some of the causes of inefficiency of N use in high input systems. Thus, loss of N from soil O.M. may be initiated by high nitrifying activity which can persist after fertilizer inputs cease (see Section 5.3.2). Also, even though the soil O<sub>2</sub> conditions conducive to these processes are opposite (aerobic and anaerobic, respectively) the opportunity may develop for adjacent soil microsites to provide appropriate conditions due to 'hotspots' of decomposition, which can rapidly consume available O<sub>2</sub> (Tiedje, 1987).

Leaching was measured during an exceptionally dry winter so that meaningful correlations could not be established. However, the soil with the higher nitrifying activity, also lost more N through leaching (see section 5.3.2) so that further investigations of this interaction, under differing conditions of winter drainage, would be warranted. It would seem probable therefore, that the increased supply of N in a more mobile form (NO<sub>3</sub><sup>-</sup>) could well lead to increased losses and this has been shown to be the case by other workers (Scholefield *et al.*, 1993).

Whilst validation of the tube incubation technique was provided by comparisons with jar incubation, the known limitations of the latter (isolation of cores from natural drying and

wetting cycles and possible disturbance of aeration and temperature regimes), meant that a more rigorous test was required. In phase three of the project, studies using  $^{15}\text{N}$  isotope techniques were introduced to measure gross rates of nitrification and a large part of the study was devoted to the acquisition of the technology and application of the necessary analytical skills (this was facilitated by a British Council funded exchange visit to New Zealand towards the end of 1996). Only by quantifying the total supply of substrate ( $\text{NH}_4^+$ ), was it possible to make definitive statements about the processes concerned from comparisons made between soils exhibiting net differences. Initial information on these relationships was obtained at an early stage of the project in an associated study carried out in the UK with the NZ partner (Ledgard *et al.*, 1998).

The experiment, combining tube incubation with  $^{15}\text{N}$  mean pool dilution techniques, showed that the measured process rates were valid.  $^{15}\text{N}$  techniques have certain limitations (see Section 6.1), but are generally more precise in the assessment of rates since the  $^{15}\text{N}$  label is uniquely identifiable. This avoids the possibility of 'double counting' which may be an artefact of some of the measurements based purely on differences between start and end soil N concentrations. Thus the close agreement between net rates, either measured directly or calculated from  $^{15}\text{N}$  labelling, provided an alternative and independent check on techniques.


### 7.3.3 Examining the practical implications

In natural systems with low inputs, nitrification rates are also generally low (see Section 1.5) and the preservation of N (in net terms) is ecologically sound. In such situations, the small, natural inputs from N-fixation (e.g. blue-green algae, *Azotobacter* etc.) and in

precipitation, are usually balanced by minimal losses through leaching and denitrification. Where swards are fertilized and other conditions non-limiting (pH, temperature, etc.), nitrifying activity increases and may persist along with concomitant increases in losses through denitrification and leaching. The maintenance of low, stable levels of inorganic N in grassland soils, despite high rates of *MIT*, suggests a dynamic equilibrium in which the favouring of one pathway must be necessarily at the expense of another. Therefore, with restricted substrate supply and if consumption is dominated by a particular pathway, e.g. by microbial uptake, then plant uptake may be constrained. The wider role implied by microbial discrimination between  $\text{NH}_4^+$  and  $\text{NO}_3^-$  (see Section 6.7) would suggest that the balance between inorganic N forms will be of vital importance in determining uptake.

#### 7.4 Sinks within the soil for mineral N

The major sinks for mineral N within the soil are the microbial and plant biomasses, and denitrification and leaching losses constitute the most important routes to external sinks. The internal N cycle is shown in Figure 7.1, which summarises the data obtained from the  $^{15}\text{N}$  experiment (see Chapter 6) in agronomic terms (*viz.*  $\text{kg N ha}^{-1}$ ). The results from the  $^{15}\text{N}$  experiment suggest that differences in nitrification are the dominant factors in determining the equilibria between the competing processes of microbial and plant sinks for mineral N. Such differences are also reflected in the relative amounts of  $\text{NH}_4^+ : \text{NO}_3^-$  in the inorganic N pools, with ratios of approximately 1, 10 and 12 : 1 under the three sward types, i.e. fertilized grass (*F*), grass/clover (*C*) and unfertilized grass (*Z*), respectively. Furthermore, the microbial sinks in the three sward types were some 4 to 5-fold larger than the plant sinks (Fig. 7.1), with proportionately higher rates of immobilization compared (on a per day basis) with plant uptake rates.



**Figure 7.1** Diagram of N cycle processes (*italics*: kg N/ha/d) and N pools (**bold**: kg N/ha) for **a**) fertilized grass, **b**) grass/clover and **c**) unfertilized grass swards. Consumption processes (not determined separately), for nitrate include denitrification and leaching and for ammonium include volatilization

#### 7.4.1 Microbial biomass

Where nitrification was inhibited (Table 6.3), immobilization rates increased in the *SMB* in *F*, *C*, and *Z*, demonstrating the affinity of microbial uptake for  $\text{NH}_4^+$ -N. The evidence was obtained from a microcosm study under natural field conditions (see Chapter 6) which overcame criticisms of extrapolation from purely laboratory-based incubations, or from field incubations where conditions are either unrepresentative or unrealistic. More subtle effects are indicated by this work and in various independent findings in the literature. For example, the conclusion that microbial uptake of N showed a preference for  $\text{NH}_4^+$  over  $\text{NO}_3^-$  (Jackson *et al.*, 1989) would indicate strong competition between heterotrophic and autotrophic consumption of  $\text{NH}_4^+$ . The previous assumption that autotrophic nitrifiers are only poor competitors for  $\text{NH}_4^+$  (Vitousek *et al.*, 1982) was challenged by more recent studies (Schimel *et al.*, 1989; Davidson *et al.*, 1990) and also by the present work. Hence the gross rate of nitrification in the fertilized (*F*) soil ( $2.0 \text{ kg N ha}^{-1} \text{ d}^{-1}$ , Fig. 7.1) was >50% of the ammonium immobilization rate, but <30% in the mixed sward (*C*) and only 15% in the unfertilized soil (*Z*).

#### 7.4.2 Plant biomass

Whilst plants can absorb  $\text{NH}_4^+$  and  $\text{NO}_3^-$  equally (Clarkson *et al.*, 1986), the dependence of roots on mass flow (mobile ions) and diffusion gradients (concentration dependent) to acquire N (Nye and Tinker, 1977), suggests that the form and concentration of inorganic N in soil solutions will influence uptake patterns. Hence, in the present study it was found that microbial uptake outstripped plant uptake, which confirmed previous work by Jackson *et al.*, (1989) and  $\text{NO}_3^-$  uptake by unfertilized swards was more efficient than that of  $\text{NH}_4^+$ , as previously proposed by Recous *et al.*, (1988a). The rapid immobilization of fertilizer

N (Bristow *et al.*, 1987) is likely, in the short-term, to limit the efficiency of fertilizer N use. In the longer-term, it can enhance N supply by saving N which might otherwise have been lost and ultimately increase the active organic N pool (Recous *et al.*, 1988b). The addition of a nitrification inhibitor (acetylene) made no significant difference to the pattern of  $\text{NH}_4^+$  uptake in these swards (see Fig. 7.1); the reasons for which will be discussed later.

### 7.5 Limitations/drawbacks identified

Whilst the combination of measurements using the tube incubation technique and  $^{15}\text{N}$  mean pool dilution technique provided important and independent comparisons, the effectiveness of the latter was decreased by sampling the soil profile down to 150 mm depth. Since the most biologically active region will be in the upper 50 mm layer, enrichment of the deeper layers will have had the effect of diluting the impact of process rates. More sensitive measurements of process rates could be achieved by confining the enrichment to the 0-100 mm depth, although the possibility of significant mineralization below this should not be overlooked and was further investigated as part of an associated study (Patra *et al.*, in press). Less disturbance of the plant processes could also have been achieved by cutting the herbage at least one week (rather than one day) before  $^{15}\text{N}$  labelling, to avoid using plants which had been only recently perturbed. Another factor, over which less control can be exercised, is the extent of spatial variability which is encountered in these highly heterogeneous pasture soils. Greater replication could have been employed, but only at the expense of a severe restriction in the range of variables which could be investigated. It might be necessary in future work to be more selective in the choice of treatments, so that replication could be greatly increased, although this is unlikely to resolve the problem

of wide variation, since the numbers required could become unmanageable. For example, using data obtained for net mineralization by tube incubation (see Section 5.3.2) with 4 replicate soil cores to each treatment ( $CV = 52\%$ ), the number of cores which would be needed to obtain an estimate of the mean with a precision of 10%, can be found using the following formula (Goovaerts and Chiang, 1993):

$$\text{sample number} = \left[ \frac{t_{\alpha} CV}{\epsilon} \right]^2$$

**Where:**  $t$  is the Student's  $t$  statistic at the desired  $\alpha$  level,  $CV$  is the coefficient of variation and  $\epsilon$  is the degree of precision required (*viz.* 0.1). With 3 degrees of freedom,  $t = 3.17$ , where  $\alpha = 0.05$ . For the example quoted, the number of incubated cores required to estimate the mean to within  $\pm 10\%$  would be 272! Therefore, the information which can be obtained from field incubations is necessarily restricted to the comparison of trends. The resolution of temporal effects may not be possible without a large investment in extra resources of time and labour.

The mean pool dilution technique described in Section 6.1 evaluates changes in the enrichment of a labelled inorganic N pool. For example, measurements of gross mineralization are derived from the rate of addition of unlabelled native ammonium to the enriched inorganic  $^{15}\text{NH}_4^+$ -N pool, which results in the dilution of the  $^{15}\text{N}$  label by sources of  $^{14}\text{N}$ . However, further information may also be obtained from examination of the unlabelled inorganic N pools. Specifically, in the example quoted above, the nitrate pool

was not labelled, and where nitrification was measured, the ammonium N pool remained unlabelled. The inorganic N concentrations of the unlabelled N pools are shown in Table 7.1. It is clear that the presence of the nitrification inhibitor (I-data set) failed to prevent a rapid increase in  $\text{NO}_3^-$ -N, following the injection of  $^{15}\text{NH}_4^+$ . The cause of this was due to the 24 h equilibration period allowed before the incubation over controlled-release capsules took place (*viz.* t0 to t1) during which  $\text{C}_2\text{H}_2$  was not supplied. Effectiveness of the acetylene infusion was nevertheless confirmed, since no further increase in nitrate concentration occurred between t1 and t5 (I-data set only). An explanation for the apparent lack of discrimination between inorganic N forms in plant uptake is therefore offered, as there is some uncertainty concerning the form of inorganic N which may have predominated during the measurement period. A possible future solution to overcome the difficulty of combining tube incubation with standard mean pool dilution techniques, whilst preserving the advantages of both methods, is discussed in the next section.

**Table 7.1** Inorganic N contents ( $\mu\text{g N g}^{-1}$  dry soil  $\pm$ SEM) of the unlabelled N pools *viz.*  $\text{NH}_4^+$ -N in the N treatments or  $\text{NO}_3^-$ -N in the A/I treatments, before injection of 20 mg N  $\text{kg}^{-1}$  soil (t0) and at 24 h (t1) and 96 h (t5) after injection (n = 6)

Treatments (N pool)	t0		t1		t5	
FA ( $\text{NO}_3^-$ -N)	5.9	(2.44)	18.1	(7.92)	16.8	(5.44)
FN ( $\text{NH}_4^+$ -N)	5.9	(0.52)	8.5	(1.52)	12.3	(4.48)
FI ( $\text{NO}_3^-$ -N)	5.9	(2.44)	31.8	(10.31)	25.7	(8.68)
CA ( $\text{NO}_3^-$ -N)	0.5	(0.15)	6.0	(8.20)	12.1	(1.66)
CN ( $\text{NH}_4^+$ -N)	4.5	(0.31)	9.9	(0.78)	10.3	(2.56)
CI ( $\text{NO}_3^-$ -N)	0.5	(0.15)	7.1	(1.73)	6.5	(3.33)
ZA ( $\text{NO}_3^-$ -N)	0.3	(0.07)	2.4	(0.68)	7.4	(2.83)
ZN ( $\text{NH}_4^+$ -N)	3.7	(0.44)	8.8	(1.51)	6.9	(0.76)
ZI ( $\text{NO}_3^-$ -N)	0.3	(0.07)	3.2	(2.18)	1.0	(0.11)

In the labelled pools, A/I treatments received  $\text{NH}_4^+$ -N and the N treatment received  $\text{NO}_3^-$ -N

## 7.6 Future priorities for investigation

The immediate stimulatory effect on nitrifying activity by the injection of  $^{15}\text{NH}_4^+$  could be largely overcome by including acetylene in the enrichment solution. Other workers have used  $\text{C}_2\text{H}_2$  dissolved in solution as a nitrification inhibitor (Kilham, 1987, Jackson *et al.*, 1989), but a novel approach would be to equilibrate the  $^{15}\text{N}$  solution with  $\text{C}_2\text{H}_2$  prior to injection. Nitrification of the newly added  $^{15}\text{NH}_4^+$  would then be prevented and the effect could be sustained using the controlled-release capsules with tube incubation.

The role of heterotrophic nitrifiers in grassland soils has been overlooked and is deserving of attention, since low fertility soils tends to favour a higher representation of the fungal (heterotrophic) component over bacteria in the *SMB*. Labelling of the  $\text{NH}_4^+$ -N pool with  $^{15}\text{N}$  and examination of the  $\text{NO}_3^-$ -N pool for unlabelled  $\text{NO}_3^-$  would give a clearer indication of heterotrophic nitrification (i.e. organic N directly nitrified to nitrate N without ammonium substrate) as shown by Schimel *et al.* (1984) in acid forest soils. This information would be required before exercising control over autotrophic nitrification, since heterotrophs are immune to some physiological blocks which are effective against autotrophs (e.g.  $\text{C}_2\text{H}_2$ ). It is also important that future studies are carried out over extended periods (*viz.* several years) to ascertain the persistence and activity of nitrification in soils from which the stimulatory effects of high N inputs have been withdrawn.

Future work should concentrate on resolving the specific nature of the interactions between nitrifying activity and N supply. For example, feedback systems could exist in which a decrease in N-containing substrates from plant decomposition will trigger greater immobilization (and hence less mineralization to  $\text{NH}_4^+$  forms) so that nitrifiers will

decrease output. A more rapid response may be exhibited in the rhizosphere in response to cutting (often followed by fertilizer application), since the exudation of C-containing compounds from roots following removal of shoots could increase both mineralization and nitrification rates. Additionally, fertilizer N will speed up *MIT* with the result that surpluses not converted to plant/microbial biomass may be discharged into the wider environment. Nitrification will certainly be involved in determining how much of the N may succumb to this fate.

### 7.7 Options for control

Nitrification inhibitors have limited commercial benefit, but can provide invaluable comparisons and insights when used in research applications and serve as useful research tools for blocking selected pathways of N transformation reactions for comparative studies (see Section 4.1). In practical terms, however, their application would seem limited because of the expense of the formulations, their short-lived activity and concerns for the detrimental effects on the environment (nitrapyrin [*N-serve*] has now been withdrawn from commercial use). This approach to problems associated with the disadvantages of excessive rates of nitrification also addresses the symptom (i.e. excess nitrogen production), rather than the cause (i.e. excess N input). Of the options available, the more sustainable would seem to be better management of N inputs and to make fuller use of soil N supply in ways which avoid excesses and are more closely matched to plant requirements (Brown *et al.*, 1997).

### 7.8 Synopsis: acceptance/rejection of hypothesis

Superficially, the hypothesis may be deemed self-evident since the major loss processes

from fertilized grasslands (denitrification and leaching) both require nitrate as a precursor. Evidence for a direct coupling of nitrification with either of these processes has been limited to circumstantial, or inappropriate data and often obtained in conditions which were unrepresentative of field systems. The reason for this has been the difficulty in transferring established laboratory techniques to in-field measurements. The development of the new field incubation technique has now made this possible and has overcome many of the limitations of previous incubation methods. Testing of the hypothesis was based on data obtained from simultaneous measurements of the processes involved using the new incubation technique.

Acceptance of the proposed hypothesis would now seem to be justified, based on the major findings from the project, from which it has been deduced that:

- Nitrification activity increases markedly and persistently with inputs of N
- There are strong correlations between nitrification rates and denitrification losses
- Increased leaching losses may be associated with higher nitrifying activities
- Nitrification influences the outcome of competition between plants and microbes
- Net mineralization is determined by the balance between gross mineralization and immobilization (chiefly the latter, which in turn is influenced by nitrification)

Whilst verification of the correlation between leaching and nitrification requires additional investigation, the pivotal role attributed to nitrification as a controlling step in N losses, would seem justified, at least in grassland soils. In arable systems, the greater rates in turnover of plant residues and increased aeration when soils are cultivated means that

nitrification rates may not be the rate-limiting step. The wider role of nitrification in influencing the outcome of competition between plant and microbial biomass for available N sources has also been indicated.

### 7.9 Conclusions

That N inputs can stimulate nitrifying organisms and that such effects have been shown to persist into the next season, following the removal of external inputs, is of major importance in understanding the responses of biological systems in agricultural soils. This project has confirmed the original observations of Jarvis and Barraclough (1991), which were based on the distribution of mineral N between  $\text{NH}_4^+$  and  $\text{NO}_3^-$  forms as indicators of nitrifying activity, by the use of more definitive laboratory assays of nitrification (Hatch *et al.*, 1998). This is especially important when considering the supplies of mineral N derived from mineralization of soil organic matter (SOM) and the fate of N determined by subsequent transformation into more mobile forms (e.g.  $\text{NO}_3^-$ ). Nitrification will continue to feature as an important component of any investigation into N cycling in grasslands; not only is it clear that many of the N transformations within and between the lithosphere and the troposphere will be initiated by nitrifiers, but the complexity of these systems means that there is still much that is poorly understood. The dearth of field data examining the important process of nitrification ensures that the present study fills an identifiable gap in our current understanding. Whilst many other data exist in isolation, this study was able to draw together the interrelationships between major processes in the N cycle and overcame many of the limitations associated with existing methods.

---oOo---

## CHAPTER 8 References

- ALEXANDER, M. (1965) Nitrification. *In: Soil Nitrogen* p. 307-343. W.V. Bartholomew and F.E. Clark (eds.) *Agronomy* 10. *Am. Soc. Agron.* Madison, Wis.
- ALEXANDER, M. (1982) Most probable number for microbial populations. *In: Methods of Soil Analysis* p. 815-820. A.L. Page, R.H. Miller and D.R. Kenney (eds.) *Agronomy* 9, Part 2. Chemical and Microbiological Properties. *Am. Soc. Agron.* Madison, Wis.
- ALEXANDER, M., MARSHALL, K. C. and HIRSCH, P. (1960) Autotrophy and heterotrophy in nitrification. *Trans. Intern. Cong. Soil Sci.* 7th Cong. Madison 2, 586-591.
- ALEEM, M. I. H., HOCK, G. E. and VANNER, J. E. (1965) Water as the source of oxidant and reductant in bacterial chemosynthesis. *Proc. Natl. Acad. Sci. USA* 54, 869-873.
- ANDERSON, O. E. and BOSWELL, F. C. (1964) The influence of low temperature and various concentrations of ammonium nitrate on nitrification in acid soils. *Soil Sci. Soc. Am. Proc.* 28, 525-529.
- ANDERSON, J.P.E. and DOMSCH, K.H. (1975) Measurement of bacterial and fungal contributions to respiration of selected agricultural and forest soils. *Can. J. Microbiol.* 21, 314-322.
- ANDERSON, J.P.E. and DOMSCH, K.H. (1978) A physiological method for the quantitative measurement of microbial biomass in soils. *Soil Biol. Biochem.* 10, 215-221.
- ARDAKANI, M. S., SCHULZ, R. K. and McLAREN, A. D. (1974) A kinetic study of ammonium and nitrite oxidation in a soil field plot. *Soil Sci. Soc. Am. Proc.* 38, 273-277.
- ARMSTRONG, A.C. and GARWOOD, E. A. (1991) Hydrological consequences of artificial drainage of grassland. *Hydrological Processes* 5, 157-174.
- BANERJEE, N. K. and MOSIER, A. R. (1989) Coated calcium carbide as a nitrification inhibitor in upland and flooded soils. *J. Indian Soc. Soil Sci.* 37, 306-313.
- BARRACLOUGH, D. (1991) The use of  $^{15}\text{N}$  tracer experiments 1. Theory. *Plant and Soil* 131, 89-96.
- BARRACLOUGH, D. and PURI, G. (1995) The use of  $^{15}\text{N}$  pool dilution and enrichment to separate the heterotrophic and autotrophic pathways of nitrification. *Soil Biol. Biochem.* 27, 17-22.

- BARGETT, R. D. and LEEMANS, D. K. (1995) The short-term effects of cessation of fertilizer applications, liming and grazing on microbial biomass and activity in a reseeded upland grassland soil. *Biol. Fertil. Soils* **19**, 148-154.
- BARTLETT, R. J. (1981) Nonmicrobial nitrite to nitrate transformation in soils. *Soil Sci. Soc. Am. J.* **45**, 1054-1058.
- BELSER, L. W. (1979) Population ecology of nitrifying bacteria. *Ann. Rev. Microbiol.* **33**, 309-333.
- BELSER, L. W. and MAYS, W. L. (1980) Specific inhibition of nitrite oxidation by chlorate and its use in assessing nitrification in soils and sediments. *Appl. Environ. Microbiol.* **39**, 505-510.
- BELSER, L. W. and MAYS, W. L. (1982) Use of nitrifier activity measurements to estimate the efficiency of viable nitrifier counts in soils and sediments. *Appl. Environ. Microbiol.* **36**, 945-948.
- BELSER, L. W. and SCHMIDT, E. L. (1978a) Diversity in the ammonia-oxidizing nitrifier population of a soil. *Appl. Environ. Microbiol.* **36**, 584-588.
- BELSER, L. W. and SCHMIDT, E. L. (1978b). Serological diversity within a terrestrial ammonia - oxidizing population. *Appl. Environ. Microbiol.* **36**, 589-593.
- BERG, P. and ROSSWALL, T. (1985) Ammonium oxidizer numbers, potential and actual oxidation rates in two Swedish arable soils. *Biol. Fert. Soils* **1**, 131-140.
- BERG, P. and ROSSWALL, T. (1987) Seasonal variations in abundance and activity of nitrifiers in four arable cropping systems. *Microb. Ecol.* **13**, 75-87.
- BERG, P., KLEMEDTSSON, L. and ROSSWALL, T. (1982) Inhibition effect of low partial pressures of acetylene on nitrification. *Soil Biol. Biochem.* **14**, 301-303.
- BHOGAL, A., HATCH, D. J., SHEPHERD, M. A. and JARVIS, S. C. (in press) Comparison of field methodologies for field measurement of net nitrogen mineralisation in arable soils: Method development. *Plant and Soil*.
- BIRCH, H. F. (1964) Mineralization of plant nitrogen following alternate wet and dry conditions. *Plant and Soil* **20**, 43-49.
- BLANTERN, P. (1991) Factors affecting nitrogen transformations in grazed grassland soils with specific reference to the effects of artificial land drainage and N-fertilizer. Ph.D Thesis, University of Exeter.
- BOHOOL, B. B. and SCHMIDT, E. L. (1980) The immunofluorescence approach in microbial ecology. *Adv. Microb. Ecol.* **4**, 203-214.

- BOHOOL, B. B., SCHMIDT, E. L. and BEASLEY, C. (1977). Nitrification in the intertidal zone: influence of effluent type and effect of tannin on nitrifiers. *Appl. Environ. Microbiol.* **34**, 523-528.
- BOON, B. and LAUDELOUT, H. (1962) Kinetics of nitrite oxidation by *Nitrobacter winogradskyi*. *Biochem. J.* **85**, 440-447.
- BOTH, G. J. GERARDS, S. and LAANBROEK, H. J. (1992) Temporal and spatial variation in the nitrite-oxidizing bacterial community of a grassland soil. *FEMS Microbiol. Ecol.* **101**, 99-112.
- BRAR, S. S. and GIDDENS, J. (1968) Inhibition of nitrification in Bladen grassland soil. *Soil Sci. Soc. Am. Proc.* **32**, 821-823.
- BREMNER, J. M. (1965) Inorganic forms of nitrogen. In C. A. Black *et al.* (ed.) Methods of soil analysis, Part 2. Agronomy **9** : 1179-1237. *Am. Soc. of Agron., Inc.* Madison, Wis.
- BREMNER, J. M. and BLACKMER, A. M. (1978) Nitrous oxide: emission from soils during nitrification of fertilizer nitrogen. *Science* **199**, 295-296.
- BREMNER, J. M. and McCARTY G. W. (1993) Inhibition of nitrification in soils by allelochemicals derived from plants and plant residues. In: Soil Biochemistry, J. M. Bollag and G. Stotzky (eds.) Vol. 8, p.181-218. Dekker (publ.).
- BRISTOW, A. W. and JARVIS, S. C. (1991) Effects of grazing and N fertilizer on the soil microbial biomass under permanent pasture. *J. Sci. Fd. Agric.* **54**, 9-21.
- BRISTOW, A. W., RYDEN, J. C. and WHITEHEAD, D. C. (1987) The fate of several time intervals of <sup>15</sup>N-labelled ammonium nitrate applied to an established grass sward. *J. Soil Sci.* **38**, 245-254.
- BROADBENT, F. E., TYLER, K. B. and HILL, G. N. (1957) Nitrification of ammonical fertilizers in some California soils. *Hilgardia* **27**, 247-267.
- BROOKES, P.C., LANDMAN, A., PRUDEN, G. and JENKINSON, D. S. (1985) Chloroform fumigation and the release of soil N: a rapid direct extraction method to measure microbial biomass N in soil. *Soil Biol. Biochem.* **17**, 837-842.
- BROWN, L., JEWKES, E., PREEDY, N. and SCHOLEFIELD, D. (1997) An integrated soil testing/modelling approach to improve fertilizer recommendations for grassland. 18 th. International Grassland Congress, Canada.
- CAMPBELL, C. A. and BIERDERBECK, V. O. (1972) Influence of fluctuating temperatures and constant soil moistures on nitrogen changes in amended and unamended loam. *Can. J. Soil Sci.* **52**, 323-326.

- CABRERA, M. L. and KISSEL, D. E. (1988) Potentially mineralizable nitrogen in disturbed and undisturbed soil samples. *Soil Sci. Soc. Am. J.* **52**, 1010-1015.
- CHIANG, C., SINNAEVE, J. and DUBUISSON, G. (1972) Ecologie microbienne des sols du Maroc 1. Fluctuations saisonnieres. *Ann. Inst. Pasteur, Paris* **122**, 1171-1182.
- CLARK, F. E. and PAUL, E. A. (1970) The microflora of grassland. *Adv. Agron.* **22**, 375-435.
- CLARKSON, D. T., HOPPER, M. J. and JONES, L. H. P. (1986) The effect of root temperature on the uptake and the relative size of the root system in *Lolium perenne*. 1. Solutions containing both  $\text{NH}_4^+$  and  $\text{NO}_3^-$ . *Plant, Cell and Environ.* **9**, 535-545.
- CLOUGH, T. J., JARVIS, S. C. and HATCH, D. J. (1998) Relationships between soil thermal units, nitrogen mineralization and dry matter production in pastures. *Soil Use Management* **14**, 65-69.
- CRUTZEN, P. J. (1970) The influence of nitrogen oxides on the atmospheric ozone content. *Q. J. Roy. Meteorol. Soc.* **96**, 320-325.
- DALTON, H. (1977) Ammonia oxidation by the methane oxidising bacterium, *Methylococcus capsulatus* strain bath. *Arch. Microbiol.* **114**, 273-279.
- DAVIDSON, E. A., ECKERT, R. W., HART, S. C. and FIRESTONE, M. K. (1989) direct extraction of microbial biomass nitrogen from forest and grassland soils of California. *Soil Biol. Biochem.* **21**, 773-778.
- DAVIDSON, E. A., SHARK, J. M. and FIRESTONE, M. K. (1990) Microbial production and consumption of nitrate in an annual grassland. *Ecology* **71**, 1968-1975.
- DAVIDSON, E. A., HART, S. C., SHANKS, C. A. and FIRESTONE, M. K. (1991) Measuring gross nitrogen mineralization, immobilization, and nitrification by  $^{15}\text{N}$  isotopic pool dilution in intact soil cores. *J. Soil Sci.* **42**, 335-349.
- DICKINSON, C. H. and PUGH, G. J. F, eds. (1974) "Biology of plant litter decomposition". Vols. 1 and 2. Academic Press, New York.
- DOMMERMUES, Y. R. (1966) "Biologie du sol." *Presses Universitaire de France*, Paris.
- DONALDSON, J. M. and HENDERSON, G. S. (1990) Nitrification potential of secondary-succession upland oak forests: 1. Mineralization and nitrification during laboratory incubations. *Soil Sci. Soc. Am. J.* **54**, 892-897.
- DUBEY, H. D. (1968) Effects of soil moisture levels on nitrification. *Can. J. Microbiol.* **14**, 1348-1350.

- DUSEK, L. (1995) Activity of nitrifying populations in grassland soil polluted by polychlorinated biphenyls (PCBs). *Plant and Soil* **176**, 273-282.
- ENGEL, M. S. and ALEXANDER, M. (1958) Culture of *Nitrosomonas europaea* in media free of insoluble constituents. *Nature* **181**, 136.
- ENO, C. F. (1960) Nitrate production in the field by incubating the soil in polyethylene bags. *Soil Sci. Soc. Am. Proc.* **24**, 277-279.
- ETINGER-TULCZNSKA, R. (1969) A comparative study of nitrification in soils from arid areas of Israel. *J. Soil Sci.* **20**, 307-317.
- FAUCI, M.F. and DICK, R.P. (1994) Soil microbial dynamics: short- and long- term effects of inorganic and organic nitrogen. *Soil Sci. Soc. Am. J.* **58**, 801-806.
- FEDERER, C. A. (1983) Nitrogen mineralization and nitrification: Depth variation in four New England forest soils. *Soil Sci. Soc. Am. J.* **47**, 1008-1014.
- FENCHEL, J. and BLACKBURN, T. H. (1979) *Bacteria and mineral cycling*. Academic Press, London.
- FINDLAY, D. C. COLBORNE, G. J. N., COPE, D. W., HARROD, T. R., HOGAN, D. V. and STAINES, S. J. (1984) *Soils and their use in South West England. Soil Survey of England and Wales Bulletin No. 14*, Harpenden.
- FIRESTONE, M. K. (1982) Biological denitrification. *In: Stevenson, F.J. (ed.) Nitrogen in agricultural soils. Agronomy Monograph 22*, 289-326.
- FOCHT, D. D. and CHANG, A. C. (1975) Nitrification and denitrification processes related to waste water treatment. *Adv. App. Microbiol.* **19**, 153-186.
- FOCHT, D. D. and VERSTRAETE, W. (1977) Biochemical ecology of nitrification and denitrification. *Adv. Microbiol. Ecol.* **1**, 135-214.
- GILL, K., JARVIS, S. C. and HATCH, D. J. (1995) Mineralization of nitrogen in long-term pasture soils: effects of management. *Plant and Soil* **172**, 153-162.
- GILMOUR, J. T., CLARK, M. D. and SIGUA, G. C. (1985) Estimating net mineralization from carbon dioxide evolution. *Soil Sci. Soc. Am. J.* **49**, 1398-1402.
- GOLBERG, S. S. and GAINEY, P. L. (1955) Role of surface phenomenon in nitrification. *Soil Sci.* **80**, 43-53.
- GOOVAERTS, P., CHIANG, C. N. (1993) Temporal persistence of spatial patterns for mineralizable nitrogen and selected soil properties. *Soil Sci. Soc. Am. J.* **57**, 372-381.

- GROFFMAN, P. M. (1987) Nitrification and denitrification in soil: A comparison of enzyme assay, incubation and enumeration methods. *Plant and Soil* **97**, 445-450.
- HARMSSEN, G. W. and van SCHREVEN, D. A. (1955) Mineralization of organic nitrogen in soil. *Adv. Agron.* **7**, 299-398.
- HART, S. C., FIRESTONE, M. K., PAUL, E. A. and SMITH, J. L. (1993) Flow and fate of soil nitrogen in an annual grassland and a young mixed-conifer forest. *Soil Biol. Biochem.* **25**, 431-442.
- HATCH, D. J., JARVIS, S. C. and PHILIPPS, L. (1990) Field measurement of nitrogen mineralization using soil core incubation and acetylene inhibition of nitrification. *Plant and Soil* **124**, 97-107.
- HATCH, D. J., JARVIS, S. C. and REYNOLDS, S. E. (1991) An assessment of the contribution of net mineralization to N cycling in grass swards using a field incubation method. *Plant and Soil* **138**, 23-32.
- HATCH, D. J., JARVIS, S. C. and PARKINSON, R. (1996) Nitrification in grassland soils. Transactions of the 9th Nitrogen Workshop, Braunschweig, September 1996, pp 239-241. Schmidt Buchbinderei und Druckerei, Braunschweig.
- HATCH, D.J., JARVIS, S. C., ROOK, A.J. and BRISTOW, A. W. (1997) Ionic contents of leachate from grassland soils: a comparison between ceramic suction cup samples and drainage. *Soil Use and Management* **13**, 68-74.
- HATCH, D. J., JARVIS, S. C. and PARKINSON, R. J. (1998) Concurrent measurements of net mineralization, nitrification, denitrification and leaching from field incubated soil cores. *Biol. Fertil. Soils* **26**, 323-330.
- HAUCK, R. D. and BREMNER, J. M. (1976) Use of tracers for soil and fertilizer nitrogen research. *Adv. Agronom.* **28**, 219-266.
- HAYNES, R. J. (1986) Mineral nitrogen in the plant-soil system (Physiological Ecology). Academic Press, London.
- HOJBERG, O., BINNERUP, S. J. and SORENSEN, J. (1996) Potential rates of ammonium oxidation, nitrite oxidation, nitrate reduction and denitrification in the young barley rhizosphere. *Soil Biol. Biochem.* **28**, 47-54.
- HYNES, R. K. and KNOWLES, R. (1982) Effect of acetylene on autotrophic and heterotrophic nitrification. *Can. J. Microbiol.* **28**, 334-340.
- HYNES, R. K. and KNOWLES, R. (1983) Inhibition of chemoautotrophic nitrification by sodium chlorate and sodium chlorite: a re-examination. *Appl. Environ. Microbiol.* **45**, 1178-1182.

- ISHAQUE, M. and CORNFIELD, A. H. (1972) Nitrogen mineralization and nitrification during incubation of East Pakistan tea soils in relation to pH. *Plant and Soil* **37**, 91-95.
- ISHAQUE, M. and CORNFIELD, A. H. (1974) Nitrogen mineralized and nitrification in relation to incubation temperature in an acid Bangladesh soil lacking autotrophic nitrifying organisms. *Trop. Agric.* **51**, 37-41.
- JACKSON, L. E., SCHIMEL, J. P. and FIRESTONE, M. K. (1989) Short-term partitioning of ammonium and nitrate between plants and microbes in an annual grassland. *Soil Biol. Biochem.* **21**, 409-415.
- JACKSON, L. E., STRAUSS, R. B., FIRESTONE, M. K. and BARTOLOME, J. W. (1988) Plant and soil nitrogen dynamics in California annual grassland. *Plant and Soil* **110**, 9-17.
- JANSSON, S. L. (1958) Tracer studies on nitrogen transformations in soil with special attention to mineralization - immobilization relationships. *Kungl. Lantbruk. Annaler* **24**, 101-361.
- JANSSON, S. L. and PERSSON, J. (1982) Mineralization and immobilization of soil nitrogen. In: Nitrogen in agricultural soils. F.J. Stevenson (ed.). *Agronomy* No. **22**, pp. 229-252.
- JARVIS, S. C. and BARRACLOUGH, D. (1991) Variation in mineral nitrogen in soils under grazed grassland soils. *Plant and Soil* **138**, 177-188.
- JARVIS, S. C. (1997) Emission processes and their interactions in grassland soils. In: Gaseous nitrogen emissions from grasslands. S. C. Jarvis and B. F. Pain (eds.). pp. 1-17. CAB International, Wallingford, UK.
- JARVIS, S. C., STOCKDALE, E. A., SHEPHERD, M. A. and POWLSON, D. S. (1996) Nitrogen mineralization in temperate agricultural soils: processes and measurement. *Adv. Agron.* **57**, 187-235
- JENKINSON, D.S. (1988) Determination of microbial biomass C and N in soil. In: Advances in Nitrogen Cycling in Agricultural Ecosystems. J.R. Wilson (ed.), pp. 368-386. CAB International, Wallingford.
- JOERGENSEN, R.G. (1996) The function of the soil microflora for mobilization / immobilization processes in the rooting zone. Transactions of the 9th Nitrogen Workshop. (September 1996) Braunschweig, pp 35-38.
- JOHNSON, P. W. and SIEBURTH, J. McN. (1976) *In situ* morphology of nitrifying-like bacteria in aquaculture systems. *Appl. Environ. Microbiol.* **31**, 423-432.

- JUSTICE, J. K. and SMITH, R. L.. (1962) Nitrification of ammonium sulphate in a calcareous soil as influenced by combinations of moisture, temperature, and levels of added nitrogen. *Soil Sci. Soc. Am. Proc.* **26**, 246-250.
- KAMPHAKE, L. K., HANNAH, S.A. and COHEN, J.M. (1967). Automated analysis for nitrate by hydrazine reduction. *Water Res. Lab.* 205-216.
- KAPLAN, W. A. (1983) Nitrification. *In: Nitrogen in the marine environment.* Carpenter, E.J. and Capone, D.G. (eds.) Academic Press, New York. pp 139-190.
- KEENEY, D. R. and BREMNER, J. M. (1967) Determination and isotope-ratio analysis of different forms of nitrogen in soils: 6. Mineralizable Nitrogen. *Soil Sci. Soc. Am. Proc.* **31**, 34-39.
- KEMPERS, A.J. and LUFT, A.G. (1988). Re- examination of the determination of environmental nitrate as nitrite by reduction with hydrazine. *Analyst* **113**, 1117-1120.
- KILHAM, K. (1986). Heterotrophic nitrification. *In: Nitrification* (J.I. Prosser, ed.) pp. 117-126. Vol. 20, Society for General Microbiology. IRL Press, Washington, D.C.
- KILHAM, K. (1987). A new perfusion system for the measurement and characterization of potential rates of soil nitrification. *Plant and Soil* **97**, 267-272.
- KIRKHAM, D. and BARTHOLOMEW, W. V. (1954). Equations for following nutrient transformations in soil, utilizing tracer data. *Soil Sci. Soc. Am. Proc.* **18**, 33-34.
- KNOWLES, R. (1978). Common intermediates of nitrification and denitrification, and the metabolism of nitrous oxide, pp 367-371. *In: D. Schlessinger (ed.) Microbiology 1978. Am. Soc. Microbiol., Washington D.C.*
- KNOWLES, R. (1982). Denitrification. *Microbiological Review* **46**, 43-70.
- KROM, M. (1980). Spectrophometric determination of ammonia: a study of modified Berthelot reaction using salicylate and dichloroisacyanurate. *Analyst* **105**, 305-316.
- KUENEN, J. G. and ROBERTSON, L. A. (1988). Ecology of nitrification and denitrification. *In: The nitrogen and sulphur cycles.* J. A. Cole and J. S. Ferguson (eds.). General Microbiology, Symposium 42, Cambridge University Press.
- KUENEN, J. G. and ROBERTSON, L. A. (1994). Combined nitrification-denitrification processes. *FEMS Microbiology Reviews* **15**, 109-117.
- KUNIN, R. and MYERS, R. J. (1947) The anion exchange equilibria in an anion exchange resin. *J. Amer. Chem. Soc.* **69**, 2874-2878.
- LAJTHA, K. (1988) The use of ion-exchange resin bags for measuring nutrient availability in an arid ecosystem. *Plant and Soil* **105**, 105-111.

- LAUDELOUT, H.; LAMBERT, R.; FRIPIAT, J. L. and PHAM, M. L. (1974) The effect of temperature on the rate of oxidation of ammonia to nitrate by mixed cultures of nitrifying organisms. *Ann. Microbiol. (Inst. Pasteur)* **125 B**, 75-84.
- LAUDELOUT, H.; FRANKART, R.; LAMBERT, R.; MOUGENOT, F. and PHAM, M. L. (1975) *In: Modelling and simulation of water resources systems.* G. C. Vansteenkiste (ed.) pp. 361-366. Amsterdam: North Holland.
- LAUDELOUT, H.; LAMBERT, R. and PHAM, M. L. (1976) The effect of pH and partial pressure of oxygen on nitrification. *Ann. Microbiol. (Inst. Pasteur)* **127 A**, 367-382.
- LEDGARD, S. F., JARVIS, S. C. and HATCH, D. J. (1998) Short-term fluxes in grassland soils under different long-term nitrogen management regimes. *Soil Biol. Biochem.* **30**, 1233-1241.
- LEHNINGER, A. L. (1981) The molecular basis of cell structure and function. *In: Biochemistry* 2nd edition p.189. Worth publishers inc.
- LOBER, R. W. and REEDER, J. D. (1993) Modified waterlogged incubation method for assessing nitrogen mineralization in soils and soil aggregates. *Soil Sci. Soc. Am. J.* **57**, 400-403.
- LOVELL, R. D., JARVIS, S. C. and BARDGETT, R. D. (1995) Soil microbial biomass and activity in long-term grassland: effects of management changes. *Soil Biol. Biochem.* **27**, 969-975.
- LOVELL, R.D., and JARVIS, S.C. (1996) Effects of urine on soil microbial biomass, methanogenesis, nitrification and denitrification in grassland soils. *Plant and Soil* **186**, 265-273.
- LOVELL, R.D., and HATCH, D. J. (1997) Stimulation of microbial activity following spring applications of nitrogen. *Biol. Fertil. Soils* **26**, 28-30.
- MACDONALD, R. M. (1979). Population dynamics of the nitrifying bacterium *Nitrosolobus* in arable soil. *J. App Ecol.* **16**, 529-535.
- MAHENDRAPP, M. K., SMITH, R. L., and CHRISTIANSON, A. T. (1966) Nitrifying organisms affected by climatic region in Western United States. *Soil Sci. Soc. Am. Proc.* **31**, 212-215.
- MAHLI, S. S. and MCGILL, W. B. (1982) Nitrification in three Alberta soils: Effect of temperature, moisture and substrate concentration. *Soil Biol. Biochem.* **14**, 393-399.
- MAHLI, S. S. and NYBORG, M. (1979) Nitrate formation during winter from fall applied urea. *Soil Biol. Biochem.* **11**, 439-441.

- McLAREN, A. D. (1971) Kinetics of nitrification in soil: Growth of the nitrifiers. *Soil Sci. Soc. Am. Proc.* **35**, 91-95.
- McLAREN, A. D. and ARDAKANI, M. S. (1972) Competition between species during nitrification in soil. *Soil Sci. Soc. Am. Proc.* **36**, 602-605
- MILLER, R. D. and JOHNSON, D.D. (1964) The effect of soil moisture tension on CO<sub>2</sub> evolution, nitrification and mineralization. *Soil Sci. Soc. Am. Proc.* **28**, 644-647.
- MONOGHAN, R. (1995). Errors in estimates of gross rates of nitrogen mineralization due to non-uniform distributions of <sup>15</sup>N label. *Soil Biol. Biochem.* **27**, 855-859.
- MORRILL, L. G. and DAWSON, J. E. (1967) Patterns observed for the oxidation of ammonium to nitrate by soil organisms. *Soil Sci. Soc. Am. Proc.* **31**, 757-760.
- MURPHY, D. V., FILLERY, I. R. P. and SPARLING, G. P. (in press) A method to label soil cores with <sup>15</sup>NH<sub>3</sub> gas as a prerequisite for <sup>15</sup>N isotopic dilution and measurement of gross N mineralization. *Soil Biol. Biochem.*
- MYERS, R. J. K. (1975) Temperature effects on ammonification and nitrification in a tropical soil. *Soil Biol. Biochem.* **7**, 83-86.
- NELSON, D. H. (1929) The isolation of some nitrifying organisms. Iowa State Coll. *J. Sci.* **3**, 113-175.
- NYE, P. H. and TINKER, P. B. H. (1977) *Solute movement in the soil-root system*. Blackwell Scientific Publications, Oxford.
- OCIO, J. A. and BROOKES, P. C. (1990). An evaluation of methods for measuring the microbial biomass in soils following recent additions of wheat straw and the characterization of the biomass that develops. *Soil Biol. Biochem.* **22**, 685-694.
- PAINTER, H. A. (1970) A review of literature on inorganic nitrogen metabolism in microorganisms. *Water Res.* Vol. 4 pp. 393-450, Pergamon Press.
- PARSONS, A. J., ORR, R. J., PENNING, P. D. and LOCKYER, D. R. (1991) Uptake, cycling and fate of nitrogen in grass-clover swards continuously grazed by sheep. *J. Agric. Sci. Camb.* **116**, 47-61.
- PATRA, A. K., JARVIS, S. C. and HATCH, D. J. (in press) Nitrogen mineralization in soil, soil particles and macro-organic matter to depth under different grassland management systems. *Biol. Fertil. Soils*.
- PAUL, E.A. and VORONEY, R.P. (1980) Nutrient and energy flows through soil microbial biomass. In: D.C. Ellwood, J.N. Hedger, M.J. Latham, J.M. Lynch and J.H. Slater (eds.) Contemporary microbial ecology. Academic Press, London, pp 215-237.

- PATRICK, W. H. and WYATT, R. (1964) Soil nitrogen loss as a result of alternate submergence and drying. *Soil Sci. Soc. Am. Proc.* **28**, 647-657.
- PAYNE, W. S. (1978) Introduction, *In: "Microbiology - 1978"*, D. Schlessinger ed., p 287. *Am. Soc. Microbiol.* Washington, D.C.
- PIPER, C. S. (1950) *In: Soil and Plant Analysis. A monograph from the Waite Agricultural Research Insitute, University of Adelaide*, pp 47-79.
- PURCHASE, B. S. (1974a) Evaluation of the claim that grass root exudates inhibit nitrification. *Plant and Soil* **41**, 527-539.
- PURCHASE, B. S. (1974b) The influence of phosphate deficiency on nitrification. *Plant and Soil* **41**, 541-547.
- RAISON, R. J., CONNELL, M. J. and KHANNA, P. K. (1987) Methodology for studying fluxes of soil mineral-N *in situ*. *Soil Biol. Biochem.* **19**, 521-530.
- RECOUS, S., FRESNEAU, C., FAURIE, G. and MARY, B. (1988a) The fate of labelled <sup>15</sup>N urea and ammonium nitrate applied to a winter wheat crop. 1. Nitrogen transformations in the soil. *Plant and Soil* **112**, 205-214.
- RECOUS, S. and MARY, B. (1988b) The fate of labelled <sup>15</sup>N urea and ammonium nitrate applied to a winter wheat crop. 2. Plant uptake and efficiency *Plant and Soil* **112**, 215-224.
- RENNIE, R. J. and SCHMIDT, E. H. (1974) Fluorescent antibody techniques to study the autoecology of *Nitrobacter* in soils. Abstracts of the Annual Meeting of the American Society for Microbiology, Chicago.
- RICE, E. L. and PANCHOLY, S. K. (1972) Inhibition of nitrification by climax vegetation. *Am. J. Bot.* **59**, 1033-1040.
- RICE, E. L. and PANCHOLY, S. K. (1973) Inhibition of nitrification by climax ecosystems. II. Additional evidence and possible role of tannins. *Am. J. Bot.* **60**, 691-702.
- RICHARDSON, H. L. (1938) The nitrogen cycle in grassland soils: with especial reference to the Rothamsted Park Grass Experiment. *J. Agric. Sci. Camb*, **28**, 73-121.
- ROBINSON, J. B. (1963) Nitrification in a New Zealand grassland soil. *Plant and Soil* **19**, 173-183.
- ROSS, D. J., TATE, K. R., CAIRNS, A. and MEYRICK, K. F. (1980) Influence of storage on soil microbial biomass estimated by three biochemical procedures. *Soil Biol. Biochem.* **12**, 369-374.
- RUSSELL, E. W. (1973) *Soil conditions and plant growth*. 10th Ed. Longman, London.

- RYDEN, J. C. (1982) Effect of acetylene on nitrification in two soils during incubation with ammonium nitrate. *J. Soil Sci.* **33**, 263-270.
- RYDEN, J. C., BALL, R. P. and GARWOOD, E. A. (1982) Nitrate leaching from grassland. *Nature* **311**, 50-53.
- RYDEN, J. C., SKINNER, J. H. and NIXON, D. J. (1987) Soil core incubation system for the field measurement of denitrification using acetylene-inhibition. *Soil Biol. Biochem.* **19**, 753-757.
- SABEY, B. R. (1969) Influence of soil moisture tension on nitrate accumulation in soils. *Soil Sci. Soc. Am. Proc.* **33**, 263-266.
- SARATHCHANDRA, S. U. (1978) Nitrification activities of some New Zealand soils and the effect of some clay types on nitrification. *N.Z. J. Agric. Res.* **21**, 615-621.
- SAHRAWAT, K. L. (1982) Nitrification in some tropical soils. *Plant and Soil* **65**, 281-286.
- SCHIMEL, J. P., FIRESTONE, M. K. and KILLHAM, K. (1984) Identification of heterotrophic nitrification in a Sierran forest soil. *Appl. Environ. Microbiol.* **48**, 802-806.
- SCHIMEL, J. P., JACKSON, L. E. and FIRESTONE, M. K. (1989) Spatial and temporal effects on plant-microbial competition for inorganic nitrogen in a California annual grassland. *Soil Biol. Biochem.* **21**, 1059-1066.
- SCHMIDT, E. L. (1982) Nitrification in soil. In: Nitrogen in agricultural soils. Agronomy Monograph No. 22 pp 253-288. *Am. Soc. Agron.* Madison, Wis.
- SCHOLEFIELD, D., TYSON, K. C., GARWOOD, E. A., ARMSTRONG, A. C., HAWKINS, J. and STONE, A. C. (1993) Nitrate leaching from grazed grassland lysimeters: effects of fertilizer input, field drainage, age of sward and patterns of weather. *J. Soil Sci.* **44**, 601-613.
- SIERRA, J. (1992) Relationship between mineral N content and N mineralization rate in disturbed soil samples incubated under field and laboratory conditions. *Aust. J. Soil Res.* **30**, 477-492.
- SMITH, K. A. and ARAH, J. R. M. (1990) Losses of nitrogen by denitrification and emissions of nitrogen oxides from soils. *Proceedings No. 299*. The Fertilizer Society.
- SORIANO, S. and WALKER, N. (1973) The nitrifying bacteria in soils from Rothamsted classical fields and elsewhere. *J. Appl. Bacteriol.* **4**, 223-240.
- SPRENT, J. I. and SPRENT, P. (1990) Nitrogen - fixing organisms. Chapman and Hall, London, 256p.

- STANFORD, G. and EPSTEIN, E. (1974) Nitrogen mineralization-water relations in soils. *Soil Sci. Soc. Am. Proc.* **38**, 103-107.
- STARK, J. M. and HART, S. C. (1996) Diffusion technique for preparing salt solutions, kjeldahl digests, and persulphate digests for nitrogen-15 analysis. *Soil Sci. Soc. Am. J.* **60**, 1846-1855.
- STEELE, K. W., WILSON, A. T. and SAUNDERS, W. M. H. (1980) Nitrification activity in New Zealand grassland soils. *N. Z. J. Agric. Res.* **23**, 249-256.
- STEVENS, R. J., LAUGHLIN, R. J., BURNS, L. C., ARAH, J. R. M. and HOOD, R. C. (1997) Measuring the contributions of nitrification and denitrification to the flux of nitrous oxide from soil. *Soil Biol. Biochem.* **29**, 139-151.
- STOJANOVIC, B. J. and ALEXANDER, M. (1958) Effect of inorganic nitrogen on nitrification. *Soil Sci.* **86**, 208-215.
- TERRY, R. E. and DUXBURY, J. M. (1985) Acetylene decomposition in soils. *Soil Sci. Soc. Am. J.* **49**, 90-94.
- TIEDJE, T. B. (1987) Soil microsites as a source of denitrification variability. *Soil Sci. Soc. Am. J.* **51**, 1194-1199.
- TIETEMA, A. and WESSEL, W. W. (1992) Gross nitrogen transformations in the organic layer of acid forest ecosystems subjected to increased atmospheric nitrogen input. *Soil Biol. Biochem.* **24**, 943-950.
- THERON, J. J. (1951) The influence of plants on the mineralization of nitrogen and the maintenance of organic matter in soil. *J. Agric. Sci., Camb.* **41**, 289-296.
- TOPP, E. and GERMON, J. C. (1986) Acetylene metabolism and stimulation of denitrification in an agricultural soil. *Appl. Environ. Microbiol.* **52**, 802-806.
- TYLER, K. B., BROADBENT, F. E. and HILL, G. N. (1959) Low temperature effects on nitrification in four California soils. *Soil Sci.*, **87**, 123-129.
- TYSON, K. C., HAWKINS, J. M. B. and STONE, A. C. (1993) Final report on the AFRC-ADAS Drainage Experiment 1982-1993. BBSRC Institute of Grassland and Environment Research, North Wyke Research Station, Okehampton, Devon, EX20 2SB.
- VANCE, E.D., BROOKS, P. C. and JENKINSON, D.S. (1987) An extraction method for measuring soil microbial biomass. *C. Soil Biol. Biochem.* **19**, 703-707.
- VAN CLEEMUT, O. and SAMATER, A. H. (1996) Nitrite in soils: accumulation and role in the formation of gaseous compounds. *Fert. Res.* **45**, 81-89.

- VERSTRAETE, W. (1981) Nitrification. *In: Terrestrial nitrogen cycles*. F.E. Clark and T. Rosswall (eds.) *Ecol. Bull. No. 33*, pp 303-314.
- VITOUSEK, P. M., GOSZ, J. R., GRIER, C. C., MELILLO, J. M. and REINERS, W. A. (1982) A comparative analysis of potential nitrification and nitrate mobility in forest ecosystems. *Ecol. Monographs* **52**, 155-177.
- VORONEY, R. P. and PAUL, E. A. (1984) Determination of  $k_c$  and  $k_N$  *in situ* for calibration of the chloroform fumigation-incubation method. *Soil Biol. Biochem.* **16**, 9-14.
- WANIELISTA, M. P. and ECKENFELDER, Jr, W. W. (1978) Advances in water and wastewater treatment. *Biological Nutrient Removal*, Ann Arbor Science (Publishers).
- WAKSMAN, S.A. (1946) Sergei Nikolaevitch Winogradsky. The story of a great bacteriologist. *Soil Sci.* **62**, 197-226.
- WALKER, N. and WICKRAMASINGHE, K. N. (1979) Nitrification and autotrophic nitrifying bacteria in acid tea soils. *Soil Biol. Biochem.* **11**, 231-236.
- WALTER, H. M., KEENEY, D.R. and FILLERY, I. R. (1979) Inhibition of nitrification by acetylene. *Soil Sci. Soc. Am. J.* **43**, 195-196.
- WARD, B. B. (1978) Nitrification in marine environments. *In: Nitrification* (J.I. Prosser, ed.) pp 157-184. Vol. 20, Society for General Microbiology. IRL Press, Washington, D.C.
- WARDLE, D.A. and PARKINSON, D. (1990) Comparison of physiological techniques for estimating the response of the soil microbial biomass to soil moisture. *Soil Biol. Biochem.* **22**, 817-823.
- WARING, S. A. and BREMNER, J. M. (1964) Ammonium production in soil under water logged conditions as an index of nitrogen availability. *Nature* **201**, 951-952.
- WARREN, G. P. (1988) Use of acetylene to improve the reliability of the assessment of soil nitrogen mineralized on aerobic incubation. *Soil Biol. Biochem.* **20**, 119-121.
- WATSON, C. J. and MILLS, C. L. (1998) Gross nitrification transformations in grassland soils as affected by previous management intensity. *Soil Biol. Biochem.* **30**, 743-753.
- WEBER, D. F. and GAINEY, P. L. (1962) Relative sensitivity of nitrifying organisms to hydrogen ions in soils and in solutions. *Soil Science* **94**, 138-145.
- WEST, A. W. (1986). Improvement of the selective respiratory inhibition technique to measure eukaryote : prokaryote ratios in soils. *J. Microbial. Methods* **5**, 125-138.

- WEST, A. W. and SPARLING, G. P. (1986) Modifications to the substrate-induced respiration method to permit measurement of microbial biomass in soils of differing water contents. *J. Microbial. Methods* **5**, 177-189.
- WESTERMAN, D. T. and CROTHERS, S. E. (1980) Measuring soil nitrogen mineralization under field conditions. *Agron. J.* **72**, 1009-1012.
- WHITEHEAD, D. C. (1981) An improved chemical extraction method for predicting the supply of available soil nitrogen. *J. Sci. Fd. Agric.* **32**, 359-365.
- WINKLER, M. (1981) *Biological Treatment of Waste-Water*. Ellis Horwood, (Publishers).
- WINOGRADSKY, S. (1890) Sur les organismes de la nitrification. *Compte. Rend. Acad. Sci. (Paris)* **110**, 1013-1016.
- WITTER, E., MARTENSSON, A.M. and GARCIA, F.V. (1993) Size of the soil microbial biomass in a long-term field experiment as affected by different N-fertilizers and organic manures. *Soil Biol. Biochem.* **25**, 659-669.
- WYLAND, L. J. and JACKSON, L. E. (1993) Evaluating nitrate recovery by ion-exchange resin bags. *Soil Soc. Sci. Am. J.* **57**, 1208-1211.
- YOSHIDA, J. and ALEXANDER, M. (1970) Nitrous oxide formation by *Nitrosomonas europaea* and heterotrophic organisms. *Soil Sci. Soc. Am. Proc.* **34**, 880-882.

## APPENDICES

### Appendix 1

#### Preparation of $^{15}\text{N}$ enriched solutions

Stable isotopes of nitrogen, although expensive, are readily available in various formulations which are suitable for experiments which require labelling of plant, soil, animal excreta, etc. However, particular enrichments may not be obtainable, so that individual requirements are best catered for by dilution from a concentrated stock.

The following example requiring a nominal 20 At.%  $^{15}\text{N}$  enrichment will serve to illustrate the calculations needed (taken from Hauck and Bremner, 1976):

Stock isotope source: Z g of 99.9 At.%  $(^{15}\text{NH}_4)_2\text{SO}_4$

Required stock: X g of 20 At.%  $^{15}\text{N}$

The procedure is to 'dilute' the  $^{15}\text{N}$  stock with a  $^{14}\text{N}$  source from standard  $(\text{NH}_4)_2\text{SO}_4$  (at natural abundance: 0.3663 At.%  $^{15}\text{N}$ )

*Then:*  $(X * 20) = (Z * 99.9) + ((X - Z) * 0.3663)$

$$20X = 99.9Z + 0.3663X - 0.3663Z$$

*Rearranging:*

$$19.6337X = 99.534Z$$

$$Z / X = 19.6337 / 99.534$$

Thus, Z needs to be 0.1973X

From these proportions an enrichment solution requiring 1.25 g N l<sup>-1</sup> @ 20 At.%\* (i.e. X = 1.25) would be prepared from:

$$Z = (1.25 * 0.1973)$$

$$Z = 0.246 \text{ g N @ } 99.9 \text{ At.}\% \text{ } ^{15}\text{N}$$

Plus: (X - Z)

$$\text{or } (1.25 - 0.246)$$

$$= 1.004 \text{ g N @ } 0.3663 \text{ At.}\% \text{ } ^{15}\text{N}$$

\*Nb This is nominally 20 At.% based for convenience on proportions by mass (i.e. not atomic weights).

The calculation for the actual enrichment is shown below:

Using <sup>12</sup>C as the standard, atomic weights are 15.0001088 (<sup>15</sup>N) and 14.0030732 (<sup>14</sup>N)

The effective atomic mass (of N) when enriched to A% is:

$$[(A * 15.0001) + \{(100-A) * 14.0031\}] / 100$$

$$= 0.0099704A + 14.003073$$

Therefore, g atoms N in 1.25 g enriched N = 1.25 / (0.0099704A + 14.003073)

Effective atomic mass of N in 99.9% enriched source is

$$[(14.0031 * 0.01) + (15.0001 * 99.9)] / 100 = 14.999111$$

Similarly, effective atomic mass of N in natural abundance source is

$$[(14.0031 * 96.63) + (15.0001 * 0.3663)] / 100 = 14.0067$$

$$\text{g atoms in } 0.246 \text{ g enriched source} = 0.246 / 14.999111 = 0.0164009$$

$$\text{g atoms in } 1.004 \text{ g natural abundance source} = 1.004 / 14.0067 = 0.0716799$$

$$\text{Total g atoms in } 1.25 \text{ g mixture} = (0.0164009 + 0.0716799) = 0.0880808$$

Therefore,

$$1.25 / (0.0099704A + 14.003073) = 0.0880808$$

Actual enrichment (A) = 18.902 At.%  $^{15}\text{N}$ .

Using the above equations the following range of standards were prepared, for the diffused and non-diffused standard (i.e. blank-correction) tests, giving nominal and actual enrichments achieved. Since the components are inter-dependent and pure forms of the sources of  $^{15}\text{N}$  and  $^{14}\text{N}$  are not readily available, this represents the most direct method of calculation:

**Table 1** Weights required for 10 mls of each standard (containing 100  $\mu\text{g N}/10\mu\text{l}$ ) as  $(\text{NH}_4)_2\text{SO}_4$  or  $\text{KNO}_3$  from stocks of unlabelled ( $^{14}\text{N}$ ) @ 0.3663 At.% or labelled ( $^{15}\text{N}$ ) @ 99.9 At.%

Standards	<i>a</i>	<i>b</i>	<i>c</i>	<i>d</i>
At. % $^{15}\text{N}$	$^{15}\text{NH}_4^+$ (mg)	$^{14}\text{NH}_4^+$ (g)	$^{15}\text{NO}_3^-$ (mg)	$^{14}\text{NO}_3^-$ (g)
nominal (actual)				
2.5 (2.38)	10.142	0.4613	14.247	0.6508
5.0 (4.84)	21.951	0.4498	30.856	0.6337
7.5 (8.33)	33.775	0.4376	47.653	0.6169
10.0 (9.36)	45.624	0.4254	64.319	0.6001

Mix: (*a* + *b*) and (*c* + *d*) to produce range of  $^{15}\text{NH}_4^+$  and  $^{15}\text{NO}_3^-$  standards as specified.

## Appendix 2

### Calculation of Microbial Respiration (CO<sub>2</sub> - C)

The amount of CO<sub>2</sub> - C respired is calculated from the titration values using 10 ml aliquots of the exposed 0.5 M NaOH, using the following formula:

$$\mu\text{g C g}^{-1} \text{ soil d}^{-1} = [12000 (A_1 - A_2) M \text{ HCl } V] / (2 Wvd)$$

where:

- A<sub>1</sub> = volume of acid using to titrate controls (ml)
- A<sub>2</sub> = volume of acid used to titrate soil samples (ml)
- M<sub>HCl</sub> = molarity of acid
- W = weight of soil (dry)
- V = volume of alkali in trap (ml)
- v = volume of alkali titrated (ml)
- 2 = molar ratio factor
- 12000 = conversion from mMol to μg C
- d = number of days incubated

### Appendix 3

#### Blank-Correction of Samples for <sup>15</sup>N diffusion

Since the mass spectrometer (VG Micromass 602E) used in this work requires at least 100 µg N in each sample to generate a robust and reliable signal, it is clear that the evaluation of enrichments in control (blanks) samples (with inherently low amounts of contaminant <sup>14</sup>N and << 100 µg N) will be unreliable.

The method of blank-correction is taken from Kelly *et al.*, 1991, and the necessary information is obtained from comparisons between diffused and non-diffused standards in which the level of contamination of <sup>14</sup>N can be found from the differential effect on the dilution of known <sup>15</sup>N standards. The mass of N in the blank (M<sub>b</sub>) can be found from the following isotope dilution equation:

$$M_b = \frac{M_{std} (E_m - E_{std})}{(E_b - E_m)} \quad (1)$$

where M<sub>std</sub> is the mass of N in the standard, E<sub>std</sub> is the <sup>15</sup>N enrichment of the non-diffused standards, E<sub>m</sub> is the enrichment measured in the diffused standards and E<sub>b</sub> is the enrichment of the blank (assumed to be 0.366%). The mass of N in the blank calculated using Equation (1), can then be used in the following equation (2) to calculate the blank-corrected <sup>15</sup>N enrichment of the sample:

$$E_s = E_m + \frac{M_b (E_m - E_b)}{M_s} \quad (2)$$

where: E<sub>s</sub> is the corrected <sup>15</sup>N enrichment of the sample and M<sub>s</sub> is the mass of N in the sample, prior to diffusion.

## Appendix 4

### Calculation of MIT Rates

Gross mineralization rates were calculated from the resultant dilutions of  $^{15}\text{N}$ -labelled  $\text{NH}_4^+$  between days 1 and 4, following the equation of Kirkham and Bartholomew (1954):

$$m = \frac{M_0 - M_1}{t} \cdot \frac{\log (H_0 M_1 / H_1 M_0)}{\log (M_0 / M_1)} \quad (1)$$

Consumption of  $^{15}\text{NH}_4^+$  includes the sum of immobilization (microbial assimilation, see later), autotrophic nitrification, volatilization and other possible fates:

$$c = \frac{M_0 - M_1}{t} \cdot \frac{\log (H_0 / H_1)}{\log (M_0 / M_1)} \quad (2)$$

where:

- $M_0$  = initial  $^{14+15}\text{N}$  pool ( $\mu\text{g N g}^{-1}$  dry soil)
- $M_1$  = post-incubation  $^{14+15}\text{N}$  pool ( $\mu\text{g N g}^{-1}$  dry soil)
- $H_0$  = initial  $^{15}\text{N}$  pool ( $\mu\text{g N g}^{-1}$  dry soil)
- $H_1$  = post-incubation  $^{15}\text{N}$  pool ( $\mu\text{g N g}^{-1}$  dry soil)
- $m$  = mineralization rate ( $\mu\text{g N g}^{-1}$  soil  $\text{d}^{-1}$ )
- $c$  = consumption rate ( $\mu\text{g N g}^{-1}$  soil  $\text{d}^{-1}$ )
- $t$  = time (4d for present study:  $t_5 - t_1$ )

The symbols M and H refer to the  $\text{NH}_4^+$ -N pool, where  $^{15}\text{NH}_4^+$  was applied, but refer to the  $\text{NO}_3^-$ -N pool where  $^{15}\text{NO}_3^-$  was applied. For nitrification rates, the symbol m is replaced by n ( $\mu\text{g N g}^{-1}$  soil  $\text{d}^{-1}$ ).

The rate at which  $^{15}\text{N}$  enters the chloroform-labile pool (v) from F-E extraction of the soil microbial biomass can be found from the following equations of Davidson *et al.* (1991):

$$i = \frac{v_t}{\left(\frac{y_0}{x_0}\right) \cdot \left(\frac{1 - e^{-kt}}{k}\right)} \quad (3)$$

where:

- i = immobilization rate ( $\mu\text{g N g}^{-1}$  dry soil)
- $v_t$  =  $^{15}\text{N}$  content of the chloroform-labile pool ( $\mu\text{g N g}^{-1}$  dry soil) in excess of background  $^{15}\text{N}$  at the end of incubation (4d for the present study,  $t_5-t_1$ )
- $(y_0/x_0)$  = initial enrichment of the  $\text{NH}_4^+$  pool
- k = rate constant, assuming the  $\text{NH}_4^+$  pool declines exponentially as organic-  $^{14}\text{N}$  is mineralized to  $^{14}\text{NH}_4^+$

where:

$$k = \left[ - \ln \left( \frac{H_1 M_0}{H_0 M_1} \right) \right] / t \quad (4)$$

In this case, the chloroform-labile pool (microbial biomass-N) represents an infinite sink for immobilized  $^{15}\text{N}$ , assuming that none of the  $^{15}\text{N}$  is remineralized during the incubation period. A simpler method of obtaining K, assumes a linear decline in the inorganic N pool

enrichment during incubation (Barraclough *et al.*, 1985), which probably holds true for short incubations. Immobilization is then calculated as the  $^{15}\text{N}$  flush from chloroform-fumigation divided by the average enrichment at the beginning and end of incubation.

## List of abbreviations

CV	Coefficient of variation
DCD	Dicyandiamide
EC	Ethyl cellulose
FA	Fluorescent antibody (stain)
FIA	Fow-injection analyser
FE	Fumigation-extraction
HMPC	Hydroxypropylmethyl cellulose
I-E	Ion-exchange (resin)
MIT	Mobilization and immobilization turnover
MS	Mass spectrometer
MNP	Most probable number
OM	Organic matter
PTFE	Polytetrafluoroethylene
PVC	Polyvinyl chloride
SEM	Standard error of the mean
SIR	Substrate induced respiration
SMB	Soil microbial biomass
SOM	Soil organic matter
SWC	Soil water content