

2014-03-12

Improving Dental Experiences by Using Virtual Reality Distraction: A Simulation Study

Tanja-Dijkstra, K

<http://hdl.handle.net/10026.1/2924>

10.1371/journal.pone.0091276

PLoS One

Public Library of Science

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

1 Improving dental experiences by using Virtual Reality distraction: a simulation study

2
3
4 Karin Tanja-Dijkstra^a, Sabine Pahl^a, Mathew P. White^b, Jackie Andrade^a, Cheng Qian^c,
5 Malcolm Bruce^d, Jon May^a, David R. Moles^d

6
7 ^a School of Psychology, Plymouth University, Plymouth, UK

8 ^b University of Exeter Medical School, Truro, UK

9 ^c School of Electronic, Electrical and Computer Engineering, University of Birmingham, UK

10 ^d Plymouth University Peninsula Schools of Medicine and Dentistry, Plymouth, UK

11
12
13 Corresponding author:

14 Karin Tanja-Dijkstra, PhD.

15 Plymouth University

16 School of Psychology

17 Drake Circus

18 Plymouth PL4 8AA

19 United Kingdom

20 Telephone: +44 1752 587560

21 Fax: +44 1752 584808

22 Email: karin.tanja-dijkstra@plymouth.ac.uk

This is the Author Version of the paper
published as:
Tanja-Dijkstra K, Pahl S, White MP,
Andrade J, Qian C, et al. (2014) Improving
Dental Experiences by Using Virtual
Reality Distraction: A Simulation Study.
PLoS ONE 9(3): e91276.
doi:10.1371/journal.pone.0091276

23

24

25

26 **Abstract**

27 Dental anxiety creates significant problems for both patients and the dental profession. Some
28 distraction interventions are already used by healthcare professionals to help patients cope
29 with unpleasant procedures. The present study is novel because it a) builds on evidence that
30 natural scenery is beneficial for patients, and b) uses a Virtual Reality (VR) representation of
31 nature to distract participants. Extending previous work that has investigated pain and anxiety
32 during treatment, c) we also consider the longer term effects in terms of more positive
33 memories of the treatment, building on a cognitive theory of memory (Elaborated Intrusions).
34 Participants (n = 69) took part in a simulated dental experience and were randomly assigned
35 to one of three VR conditions (active vs. passive vs. control). In addition participants were
36 distinguished into high and low dentally anxious according to a median split resulting in a 3 x
37 2 between-subjects design. VR distraction in a simulated dental context affected memories a
38 week later. The VR distraction had effects not only on concurrent experiences, such as
39 perceived control, but longitudinally upon the vividness of memories after the dental
40 experience had ended. Participants with higher dental anxiety (for whom the dental
41 procedures were presumably more aversive) showed a greater reduction in memory vividness
42 than lower dental-anxiety participants. This study thus suggests that VR distractions can be
43 considered as a relevant intervention for cycles of care in which people's previous
44 experiences affect their behaviour for future events.

45

46

47

48 **Introduction**

49 Patient pain and anxiety are undesirable side-effects of many medical procedures and can
50 affect the patient's willingness to undergo treatment [1], [2]. Medical (e.g. analgesic)
51 interventions to reduce pain during treatments are frequently used but can be expensive and
52 may have their own side-effects. Simple, non-invasive alternatives, such as "distraction
53 therapy" are therefore desirable. The use of virtual reality (VR) as a distraction tool is
54 receiving growing attention in medical contexts.

55 Distraction is thought to help patients cope with pain and other aversive experiences
56 and is often combined with relaxation or pleasant imagery [3], although the psychological
57 mechanisms underlying its effects are not well understood [4]. VR distraction during aversive
58 experiences can improve coping with pain [5], lower experienced level of itching for chronic
59 puritis patients [6], and reduce the perceived duration of procedures [7]. A recent systematic
60 review of eleven studies looked at the effectiveness of virtual reality distraction on pain
61 reduction [8]. They concluded that more sophisticated VR techniques, capable of completely
62 immersing the individual were associated with greater pain relief. According to Gold and
63 colleagues [9] VR provides a powerful means of modifying affect, because of its immersive
64 nature.

65 Most previous work has considered the effects of VR distraction on pain and anxiety
66 during treatment. Distraction may also have lasting effects in terms of more positive
67 memories of the treatment, leading to a greater willingness to return for treatment. The aim of
68 the current study was to study both immediate and more long-term effects of VR distraction
69 in a simulated dental context. We chose a simulated rather than real treatment for ethical
70 reasons, as we wanted to include participants high in dental anxiety, for whom a simulated
71 treatment would be stressful already.

72 Dentistry has received relatively little attention from VR researchers, yet it is one of the most
73 common healthcare encounters. Dental anxiety is very common [10] and anxious patients are
74 less likely to keep their appointments [11], take longer to treat and feel less satisfied with
75 their treatment [12], and make their dentists feel anxious too [13]. Armfield and colleagues
76 [14] described a vicious cycle of dental anxiety. This suggests that people with high dental
77 fear delay dental treatment, which can lead to more extensive dental problems and
78 symptomatic visiting patterns which in turn maintain or exacerbate existing dental fear.
79 Memories and expectations thus play a crucial role in sustaining dental anxiety. Although we
80 focus on dental treatment, experiences and expectancies are very important in determining
81 future uptake of treatment in a range of medical contexts, e.g., unpleasant bowel
82 examinations [15].

83 VR distraction during dental treatment may improve the treatment experience and, by
84 doing so, help break the cycle of negative experiences leading to negative memories and
85 expectations about future treatment. The Elaborated Intrusion theory [16] argues that
86 unconscious cognitive activity triggered by cues in the world, mind or body can lead to
87 apparently spontaneous intrusive thoughts, and that salience of the intrusion can lead to the
88 thought being elaborated, through the construction of mental imagery. Heightened emotion
89 and arousal during a dental examination will increase the likelihood of recollections of the
90 event being triggered uncontrollably by situational cues [17], as a whiff of antiseptic might
91 trigger thoughts about dental treatment. Attempts at suppressing these intrusive thoughts tend
92 to be counterproductive [18], and once triggered, intrusive thoughts tend to be elaborated [5].
93 For example, an intrusive thought about going to the dentist might lead to the patient
94 imagining how uncomfortable the next visit is going to be and experiencing some of the
95 negative sensations and emotions they associate with dental treatment. Interfering with the
96 processing of negative stimuli during treatment, through VR distraction, would counteract the

97 effects of heightened emotion and arousal and so reduce the likelihood of intrusive thoughts
98 and negative elaborations following treatment. Additionally, it would be desirable to identify
99 if VR distraction is a suitable technique for patients with all levels of dental anxiety or
100 whether specific patients would be most likely to benefit. We therefore included level of
101 dental anxiety as a moderating variable.

102 There have been a few studies of VR in a dental context. A case study showed that
103 VR distraction is more effective in offering pain control than watching a video or a standard
104 care situation without distraction [19]. One study investigated the effects of using an A/V
105 eyeglass system displaying an instructional video [20]. Adult patients scheduled for dental
106 prophylaxis were distracted during half of their treatment. Patients reported less anxiety and
107 discomfort when using the equipment. In another study, patients undergoing periodontal
108 scaling and root planning procedures were presented with either a control situation (only
109 wearing the headgear), a video (i.e. the animation movie Cars) and a VR environment (of a
110 botanical garden in Second Life) [21]. Both distracters, relative to the control condition,
111 resulted in less pain and discomfort and lower blood pressure and pulse rate, but the VR
112 environment was better on all indicators compared to the movie. This difference can possibly
113 be explained by looking at the level of interactivity VR distraction offers compared to
114 passively watching a video.

115 Dahlquist and colleagues [5] tested the role of interactivity more directly, by assessing
116 pain tolerance and pain threshold in children using the cold pressor task. In a within-subjects
117 design, the children played a computer game, Finding Nemo, and watched a video of
118 someone else playing the exact same computer game. Both types of distraction reduced pain
119 threshold, but pain tolerance was almost twice as long during interactive distraction relative
120 to passive distraction. The authors suggested that the interactive distraction involved two

121 additional sensory attentional pathways and that the game required problem-solving,
122 providing an active cognitive processing component.

123 The current study used a VR environment of a coastal nature area to distract
124 participants during simulated dental treatment. This environment was chosen as previous
125 research demonstrated the beneficial effects of nature [22], [23], in particular coastal
126 environments [24], [25]. We investigated whether offering such distraction improved the
127 dental experience both immediately and a week later. We also investigated active versus
128 passive use of the same VR environment and the role of pre-existing dental anxiety. One of
129 the concerns with VR distraction is that it might affect patient-clinician communication.
130 Therefore we tested whether the VR interfered with this by recording compliance with the
131 dentist's requests.

132 In terms of overall experience, first we hypothesized that providing VR distraction
133 during simulated dental treatment would result in lower time perception compared to no VR
134 distraction, based on previous research suggesting that the use of VR can affect time
135 perception [7]. Second, in accordance with EI theory, we proposed that offering VR
136 distraction results in less vivid memories and less intrusive thoughts a week later.

137 The second set of hypotheses focussed on the comparison of active and passive VR.
138 We predicted that the active VR group would experience a higher level of control
139 (manipulation check) and a higher level of presence. Third, we predicted that the effects for
140 the overall experience both immediately and after a week would be stronger for the active VR
141 group compared to the passive VR group.

142 The third set of hypotheses proposed that pre-existing dental anxiety would moderate
143 these effects. We hypothesized that the effects for the dental experience, the VR experience,
144 and the follow-up effects, would be more pronounced for participants higher in dental anxiety.

145

146 **Method**

147 *Ethics statement*

148 The study was approved by the Faculty of Science and Technology ethics review board,
149 Plymouth University. Participants signed a consent form prior to participating, which was
150 approved by the ethics review board.

151

152 *Participants*

153 Seventy-five participants were recruited through a participant pool containing general
154 public as well as university staff and students. They received £4 for their participation. Data
155 from six participants were excluded because of technical failures (crashed VR environment;
156 remote control stopped working) that required intervention from the experimenter, leaving
157 data from 69 people (28 male, mean age = 33.1 years, SD = 12.7). A one-week follow-up
158 telephone interview (mean = 7.13 days, SD = .42) collected data from 62 participants. We
159 called participants up to 3 times within the set-up appointment time frame and sent an email
160 to reschedule if they did not respond to the phone calls. Seven participants did not pick up
161 their phone on any of the occasions or responded to the email so their follow-up data is
162 missing. Of the seven participants who did not complete the follow-up part of the study, five
163 (71%) were part of the control condition. Please refer to Figure 1 for the participant flow-
164 chart.

165 Data on oral health characteristics showed that 29% of the participants had no fillings,
166 52% between 1 and 5 fillings, 16% between 6 and 10 fillings and 3% had more than ten
167 fillings. One third of the study population had had at least one wisdom tooth removed. The
168 last visit to the dentist was in the last month for 13% of the participants. Another 20% went
169 2-3 months ago, 13% 4-6 months ago, 28% 6-12 months ago, 16% 1-2 years ago and another
170 10% longer than 2 years ago.

171

172 ***Design***

173 Participants were exposed to one of three conditions in a fully randomised between
174 participant design: Control no VR; Active VR; Passive VR). In addition participants were
175 split into high vs. low Dental Anxiety based on their Dental Anxiety scores collected at the
176 start of the study. This effectively produced a 3 (Condition: Control; Active VR; Passive VR)
177 by 2 (Baseline Dental anxiety: High/Low) between participant design.

178 The difference between the active and the passive VR groups was that the first group
179 was able to actively navigate the VR environment by using a controller. The passive group
180 was a yoked control group; participants in this group watched a recording of the VR walk that
181 the previous participant in the active condition generated. A total number of 22 walks were
182 generated by the active participants and each of these walks was shown to a participant in the
183 passive group. Taken together, both VR groups were thus shown the exact same content.
184 Participants in the control group wore the head-mounted device (HMD) but only saw a black
185 screen. In most research on VR distraction, a VR group is compared to a standard care
186 situation (either between or within-subjects) [4], [8]. Although such a set-up allows for
187 conclusions to be drawn regarding the effectiveness of VR distraction, it does not provide an
188 answer to the question if it is the presence of the VR environment or the exclusion of the
189 medical environment that accounts for the effect. In the current study we chose to include a
190 black-screen control group to add this perspective and to be able to attribute the effects to the
191 presence of a VR environment.

192

193 ***Procedure***

194 Participants completed an online dental anxiety questionnaire when they enrolled in the study,
195 at least 24 hours prior to the experimental session.

196

197 *Setting*

198 A simulated dental waiting and treatment area was created, using cues usually present in
199 those areas. One part of the lab represented a waiting area with a row of chairs, and posters
200 on the wall depicting dental information (see Figure 2). Here we took informed consent,
201 collected baseline data and explained the procedure. A simulated treatment area was created
202 in the other part of the lab (see Figure 3), with a dental chair, overhead light, dental
203 instruments and a dentistry-related smell (drops of oil of cloves on cotton wool). The
204 experimenter was wearing a white lab coat.

205

206 *The simulated dental experience*

207 Participants sat in the dental chair and listened to an audio tape of a dental treatment
208 (performed by a practicing dentist), involving the administration of local anaesthetics, cavity
209 preparation and filling, and an uncomplicated removal of a small upper wisdom tooth. They
210 were asked to open their mouth during this simulated dental treatment and follow the
211 instructions on the recording, for example ‘to open their mouth really wide’. They were
212 reassured that their mouth would not be touched at any point. At baseline we measured heart
213 rate and blood pressure. During the simulated treatment we measured heart rate, and
214 immediately following treatment we measured blood pressure¹. Afterwards we collected
215 measures on their experience of the event, the VR experience, demographic (age, gender and
216 education) and background information (number of fillings, removal of a wisdom tooth, last
217 and next dental visit, familiarity with the VR environment) with computer-based
218 questionnaires. An appointment was made for a telephone call one week later and participants
219 received their honorarium. Following research using the Elaborated Intrusions paradigm [26],

¹ Preliminary analysis found no significant differences in the temporal patterns of heart rate and blood pressure as a function of condition so physiological results are not considered further.

220 [27], one week later intrusive thoughts and vividness of memories were measured and
221 participants were debriefed.

222

223 *Virtual environment and VR equipment*

224 The virtual environment (VE) depicted an existing environment, which consists of a coastal
225 path, complete with sea, beach and field areas (see Figure 4), originally developed for
226 restorative and rehabilitative environment studies [28]. The VE was constructed using
227 commercially-sourced topographical geometry and aerial photographic images, and the
228 resulting 3D model was used as a template to enable the VE to be populated with additional
229 3D assets and photographic textures, including the accurate representations of the few
230 buildings at the site, trees, plants and other features.

231 A Vuzix iWear VR920 headset was connected to an Alienware M11X laptop (dual-
232 core, 1.3GHz Intel processor with Nvidia GT 540M graphics card) and used to display the
233 VE. The headset consists of two LCD displays with a 640x480 resolution, provides a 32-
234 degree field of view and weighs 3.2 ounces. Head tracking of the HMD was switched off due
235 to the context, since it would be inadvisable for the participant to move their head during
236 dental treatment. Participants in the active condition were able to explore the VE in a first-
237 person perspective, by using a Zeemote JS1 Thumbstick Controller. This controller was also
238 used to look around.

239

240 *Measures*

241

242 *Moderator*

243 *Dental Anxiety* was measured using the modified dental anxiety scale, which is often used in
244 clinical practice to assess patients' level of dental anxiety [29]. This 5-point scale, ranging

245 from not anxious [1] to extremely anxious [5] contains 5 items and a sum score was
246 calculated as in indicator of dental anxiety. Participants were divided into high-and low-
247 dental anxiety groups based on a median split (median = 13, range 6-22), resulting in a low
248 dental anxiety group of 37 participants scoring 6 to 13 ($M = 9.76$, $SD = 2.23$) and a high
249 dental anxiety group of 32 participants scoring 14 to 22 ($M = 17.06$, $SD = 2.26$).

250

251 *Immediate dental experience*

252 *Compliance* with the four requests made by the dentist on the tape was recorded; participants
253 were for example instructed to open their mouth really wide. This resulted in a score between
254 two and four since there was no non-compliance amongst the participants. The sum score of
255 the four items was used as a measure of compliance and totally compliant participants
256 (scoring 4) were compared with not totally compliant participants (scoring 2 or 3).

257 To measure *time perception* participants were asked to estimate how long they thought the
258 simulated treatment lasted for (actual time: 5 minutes and 43 seconds). The ratio of subjective
259 duration to objective duration was calculated. A perfect estimation is indicated by a ratio of
260 1.0, whereas ratios higher than 1.0 indicate overestimation and ratios lower than 1.0 indicate
261 underestimation.

262

263 *VR experience*

264 *Perceived control* ($\alpha = .66$) was included as a manipulation check for the active versus
265 passive VR manipulation using a scale based on the dominance dimension of the PAD-model
266 [30]. This bipolar scale ranged from [1] to [9]. Sample items include “in control/controlled”
267 and “guided/autonomous”.

268 Level of *presence* ($\alpha = .86$) was assessed in both VR groups using six items selected from the
269 IGroup Presence Questionnaire [31] and the Reality Presence Questionnaire [32] and the

270 average score was calculated as an indicator of level of presence. An 11-point verbal rating
271 scale, ranging from [1] to [11] was used and sample items include “I was completely
272 captivated by the virtual world” and “How real did the virtual world seem to you?”.
273 Participants were also asked to indicate their *awareness of the surrounding environment*
274 when wearing the HMD and to indicate to what extent they would choose to wear goggles or
275 use VR during a real dental visit as a measure of behavioural intention. Both items were
276 measured on an 11-point verbal rating scale, ranging from [1] to [11].

277

278 *Follow-up dental experience*

279 For the purpose of the current study we developed a questionnaire that assessed intrusive
280 thoughts of the experience and vividness of memories of the experience. This questionnaire is
281 based on the Alcohol Craving Experience Questionnaire [33] which was developed to
282 measure vividness of memories and intrusive thoughts in a different context. We assessed
283 whether participants suffered from *intrusive thoughts* about the experience ($\alpha = .81$) and the
284 *vividness of memories* ($\alpha = .69$). Intrusive thoughts were assessed with two items on an 11-
285 point verbal rating scale ranging from *not at all* [0] to *constantly/extremely* [10] and an
286 average score was calculated. The items were “How often have you thought about the visit in
287 the past week?” and “To what extent did your thoughts about the visit pop into your mind
288 spontaneously?”. The vividness of memories was measured with 5 items on an 11-point
289 verbal rating scale ranging from *not at all* [0] to *extremely vividly* [10] and the average score
290 was calculated. Sample items include “How vividly do you do you feel the emotions you
291 experienced?”, “How vividly do you remember the discomfort of holding your mouth open?”,
292 and “How vividly do you imagine the sounds?”.

293

294 *Statistical procedure*

295 A series of Analyses of Variance (ANOVA) with a 3 (condition: VR active, VR passive,
296 control) x 2 (dental anxiety: high, low) between-participant design with planned contrasts
297 were carried out. The first contrast tested the difference between VR (both active and passive
298 together) and the no VR control group. The second contrast tested the difference between the
299 active and passive VR groups. Additionally, the interaction effects between VR condition and
300 dental anxiety were examined to understand the role of dental anxiety. Degrees of freedom
301 may vary across analyses due to the loss of participants at follow-up and not all measures
302 being relevant for all groups in the study. A chi-square test was used for the not normally
303 distributed data of the compliance measure.

304

305 **Results**

306 Table 2 includes the means and standard deviations for the three groups on the different
307 outcome measures. All met assumptions of normality with acceptable skewness and kurtosis
308 apart from compliance, which was high with 75% of all participants complying with all four
309 requests, and no-one missing more than two requests.

310

311 *Baseline characteristics*

312 No baseline differences between the experimental groups were found regarding participants'
313 demographic variables, oral health characteristics, and familiarity with the VR environment,
314 all $ps > .05$.

315

316 *Immediate dental experience*

317 Comparing totally compliant and not totally compliant participants, the passive group were
318 most compliant with only one person not being totally compliant; five participants in the
319 active group and eight in the control group missed one or two requests ($\chi^2(2) = 6.27$,

320 $p = .043$). No moderating effect of dental anxiety was present, all $ps > .05$. No effects of VR
321 condition were found on time perception ($F < 1$), but the main effect of dental anxiety
322 approached significance ($F(1,63) = 3.76, p = .057, \eta_p^2 = .06$). Participants with higher dental
323 anxiety made a larger overestimation ($M = 1.42, SD = .52$) than those with lower dental
324 anxiety ($M = 1.18, SD = .48$). While the actual time of the treatment was 5.7 minutes,
325 participants with high dental anxiety estimated it lasted for 8.1 ($SD = 3.0$) minutes and
326 participants with lower dental anxiety estimated 6.8 ($SD = 2.7$) minutes.

327

328 *VR experience*

329 The manipulation check of perceived control showed that participants in the active VR group
330 experienced a higher level of control than those in the passive VR group ($F(1,66) = 4.38, p$
331 $= .040, \eta_p^2 = .06$).

332 The active VR group experienced a higher level of presence than the passive VR
333 group ($F(1,41) = 4.77, p = .035, \eta_p^2 = .10$). An interaction between VR condition and dental
334 anxiety was found ($F(1,41) = 4.23, p = .046, \eta_p^2 = .09$). Participants with a higher level of
335 dental anxiety felt more presence in the VR if they could actively control it ($M = 6.86, SD =$
336 1.57) than if they were passively watching it ($M = 4.92, SD = 1.89; F(1,41) = 9.22, p = .004,$
337 $\eta_p^2 = .18$; see Figure 5).

338 Although the results for participants' awareness of the surrounding environment were
339 in the expected direction, with the active VR group being the least aware, the passive VR
340 group slightly more aware and the control group most aware, these differences did not reach
341 statistical significance and we found no interaction effect for dental anxiety ($F < 1$).

342 Participants were asked to indicate to what extent they would choose to wear goggles
343 or use VR during a real dental visit. Participants in the VR groups were more interested to use
344 VR during a dental visit than participant in the control group ($F(1,63) = 4.19, p = .045, \eta_p^2$

345 = .06). And more importantly, we also found a main effect of dental anxiety. Participants
346 with more dental anxiety ($M = 8.63, SD = 1.88$) were more interested to use VR during real
347 dental treatment than those with lower levels of dental anxiety ($M = 7.16, SD = 2.75; F(1,63)$
348 $= 4.93, p = .030, \eta_p^2 = .07$).

349

350 *Follow-up dental experience*

351 No effects were found on the intrusive thoughts participants experienced as a consequence of
352 VR distraction ($F < 1$), or on vividness of memories ($F(1,56) = 2.55, p = .12$). A
353 main effect was found for dental anxiety ($F(1,56) = 4.89, p = .031, \eta_p^2 = .08$) on intrusive
354 thoughts. Participants with more dental anxiety ($M = 3.10, SD = 1.44$) experienced more
355 intrusive thoughts than those with lower levels of dental anxiety ($M = 2.11, SD = 1.84$). No
356 interaction effect for dental anxiety was found ($F < 1$). A main effect for dental anxiety was
357 also found ($F(1,56) = 4.92, p = .031, \eta_p^2 = .08$) for vividness of memories.

358 Most importantly, a significant interaction between VR condition and dental anxiety
359 was found for vividness of memories ($F(2,56) = 4.06, p = .023, \eta_p^2 = .13$). Simple main
360 effect analysis showed that for participants with higher dental anxiety, both active ($M = 4.35,$
361 $SD = .79$) and passive VR ($M = 4.34, SD = 1.69$) distraction resulted in less vivid memories
362 compared to the black-screen control group ($M = 6.23, SD = 1.46; F(2,56) = 3.89, p = .026,$
363 $\eta_p^2 = .12$; see Figure 6). This shows that VR was successful at interrupting the memory
364 process in particular for highly anxious participants.

365

366 **Discussion**

367 Our research extends previous VR studies by showing that VR distraction in a simulated
368 dental context affected memories a week later. The VR distraction had effects not only on
369 concurrent experiences, but also longitudinally upon the vividness of memories after the

370 dental experience had ended. Participants higher in dental anxiety (for whom the procedures
371 were presumably more aversive) showed a greater reduction in memory vividness than lower
372 dental-anxiety participants. This is an important extension because it helps us understand the
373 cognitive processes by which VR distraction can work.

374 Dental anxiety is associated with the tendency to experience negative or threatening
375 thoughts concerning treatment [34] and this may prevent patients arranging and attending
376 dental appointments. Our findings suggest that VR distraction has the potential to influence
377 people's memories of a potentially anxiety-inducing medical event. Our results are promising
378 for real dental procedures in suggesting that VR distraction during dental treatment has the
379 potential to interrupt the cycle of dental anxiety [14], by blocking the development of vivid
380 memories.

381 It is important to note that the current study took place in a simulated environment.
382 We chose a simulated rather than real treatment for ethical reasons, as we wanted to include
383 participants high in dental anxiety, for whom a simulated treatment would be stressful already.
384 And whilst we do find differential effects for participants high and low in dental anxiety,
385 there were no differences on the physiological measures between these two groups. This does
386 mean that we are currently unable to draw any conclusions regarding the effectiveness of VR
387 distraction during real dental treatment. As with any experimental study, there always is the
388 worry about demand characteristics. We do however find a moderating effect on dental
389 anxiety, and do not think it likely that people with higher or lower dental anxiety would differ
390 in their desire to comply with an experimenter. Next, we collected the measure of vividness
391 of memories one week later. If demand characteristics really were at play in this study, we
392 would expect a lot of participants to still remember all the details and what we would
393 possibly want them to answer. Also, we would presume that demand characteristics would

394 play a greater role in within-subject designs where participants are exposed to all conditions,
395 while the current study employed a between-subjects design.

396 Frere and colleagues [20] suggest that the use of VR equipment will be particularly
397 useful for long procedures or treatment of patients who have to have repeated procedures. In
398 order to realize cost-effective VR distraction interventions, it would be desirable to identify
399 those patients that will most benefit from this. Our findings suggest that anxious patients,
400 rather than being resistant to distraction interventions, would be most likely to benefit from
401 VR. Interestingly, participants with more dental anxiety were also more interested to use VR
402 during real dental treatment than those with lower levels of dental anxiety, and especially
403 participants with a higher level of dental anxiety felt more presence in the VR if they could
404 actively control it than if they were passively watching a recording. These results are in line
405 with the ideas about how anxiety influences attention [35] and suggest that VR distraction, or
406 possibly any distraction intervention, could be particularly suitable for this high anxiety
407 group. We recognise that no real-time recordings of anxiety were gathered during the
408 simulated treatment, primarily to avoid the participant having to disengage with the
409 immersive scenario, and thus we are unable to comment on the temporal patterns in anxiety
410 during VR distraction. Future research could monitor how anxiety might be affected at
411 different stages during treatment.

412 Previous research found that interactive VR was better than passive VR in children
413 experiencing experimentally induced pain [5], [36]. Our participants in the active group
414 experienced a higher level of control and presence, and participants in the passive VR group
415 were more compliant than active and control participants, yet active versus passive VR had
416 no effects on immediate outcomes or a week later. More research is needed to decide whether
417 this is because we used a calming natural environment that people simply walked around in
418 (rather than an interactive game), or whether this was due to our simulated context or adult

419 sample. Most research in the domain of VR distraction made use of existing video games as
420 the distractor, e.g. [5], [37], or games developed for the purpose of using it as a VR distractor
421 [38], [39]. Both types of games have proven to be effective distractors, but it is unclear
422 whether gaming elements, such as providing a goal, are required for a VR distraction
423 intervention to be effective.

424 A variety of other imagery and stimuli has been used to distract patients in previous
425 research including natural contexts such as forests [40] and a botanical garden [21]. Research
426 on restorative environments suggests that certain environments are capable of relaxing people,
427 especially natural environments [41]. Hence we would call for more research that addresses
428 the content of VR interventions to help us understand which specific elements are successful.

429 The cognitive effects were measured at one-week follow-up, following the Elaborated
430 Intrusions paradigm. One might argue that a one-week follow up assessment of memories
431 does not reflect the amount of time that is usually present between dental appointments.
432 While it is not the most common situation, a variety of treatments do require patients to
433 return a week later for the next part of their treatment, for example when crowns or dentures
434 are needed. Also, the current study only offered a first test of this elaborated intrusions
435 account, so it did seem prudent to test the effect at one week follow-up first before investing
436 in studies with a more longitudinal character. Arguably, the week immediately following
437 such an experience is crucial for consolidating and processing any relevant memories.

438 One of the claims that is often made for the usage of VR as a distraction technique is
439 that wearing a HMD effectively excludes the surrounding medical environment. For example,
440 the appearance of the nurse who cleans patient's wounds may be a strong enough cue to
441 create anxiety [38]. The overhead light and the dental instruments may induce anxiety in a
442 similar way even in a simulated context. In the current study we chose to include a black-
443 screen control group to add this perspective and to be able to attribute the effects to the

444 presence of a VR environment. However, further research is needed to decide if it is the
445 presence of the VR environment or the exclusion of the medical environment that accounts
446 for the effect.

447 Taken together, the current study provides evidence that a VR distraction intervention
448 can not only impact the experience of a simulated aversive event, it can also reduce the
449 vividness of memories of such an event a week later. This study thus suggests that VR
450 distractions can be considered as a relevant intervention for cycles of care in which people's
451 previous experiences affect their behaviour for future events. If a dental patient for example
452 has a more positive experience of a treatment due to the VR distraction intervention, that
453 patient might have less vivid memories and as a consequence might be less likely to postpone
454 a future dental visit.

455

456

457 **Acknowledgements**

458 The authors thank Kimberley Schenke for her assistance in data collection and Prof. R.J.
459 Stone for his involvement in developing the 3D coastal environment.

460

461 **References**

462 1. Maggiriias J, Locker D (2002) Psychological factors and perceptions of pain associated
463 with dental treatment. *Community Dent Oral Epidemiol* 30: 151-159.

464 2. Bare LC, Dundes L (2004) Strategies for combating dental anxiety. *J Dent Educ* 68: 1172-
465 1177.

466 3. McCaul KD, Monson N, Maki RH (1992) Does distraction reduce pain-produced distress
467 among college students? *Health Psychol* 11: 210-217.

- 468 4. Mahrer N, Gold J (2009) The use of virtual reality for pain control: A review. *Cur Pain*
469 *Headache Rep* 13: 100-109.
- 470 5. Dahlquist LM, McKenna K D, Jones KK, Dillinger L, Weiss KE, Ackerman CS (2007)
471 Active and passive distraction using a head-mounted display helmet: effects on cold
472 pressor pain in children. *Health Psychol* 26: 794-801.
- 473 6. Leibovici V, Magora F, Cohen S, Ingber A (2009) Effects of virtual reality immersion and
474 audiovisual distraction techniques for patients with pruritus. *Pain Res Manag* 14: 283-286.
- 475 7. Schneider SM, Hood LE (2007) Virtual reality: A distraction intervention for
476 chemotherapy. *Oncol Nurs Forum* 34: 39-46.
- 477 8. Malloy KM, Milling LS (2010) The effectiveness of virtual reality distraction for pain
478 reduction: A systematic review. *Clin Psychol Rev* 30: 1011-1018.
- 479 9. Gold JI, Belmont KA, Thomas DA (2007) The neurobiology of virtual reality pain
480 attenuation. *CyberPsychol Behav* 10: 536-544.
- 481 10. Oosterink FMD, de Jongh A, Hoogstraten J (2009) Prevalence of dental fear and phobia
482 relative to other fear and phobia subtypes. *Eur J Oral Sci* 117: 135-143.
- 483 11. Kleinknecht RA, Bernstein DA (1978) The assessment of dental fear. *Behav Ther* 9: 626-
484 634.
- 485 12. Locker D, Liddell AM (1991) Correlates of dental anxiety among older adults. *J Dent Res*
486 70: 198-203.
- 487 13. Hill KB, Hainsworth JM, Burke FJT, Fairbrother KJ (2008) Evaluation of dentists'
488 perceived needs regarding treatment of the anxious patient. *Br Dent J* 204: E13.
- 489 14. Armfield J, Stewart J, & Spencer, AJ (2007) The vicious cycle of dental fear: exploring
490 the interplay between oral health, service utilization and dental fear. *BMC Oral Health* 7: 1.
- 491 15. Redelmeier DA, Katz J, Kahneman, D (2003) Memories of colonoscopy: a randomized
492 trial. *Pain* 104: 187-194.

- 493 16. Kavanagh DJ, Andrade J, May J (1995) Imaginary relish and exquisite torture: The
494 Elaborated Intrusion theory of desire. *Psychol Rev* 112: 446-467.
- 495 17. Brewin CR, Dalgleish
496 T, Joseph S (1996) A dual representation theory of posttraumatic stress disorder. *Psychol*
497 *Rev* 103: 670-686
- 498 18. Wegner DM, Scheider DJ, Carter SR, White TL (1987) Paradoxical effects of thought
499 suppression. *J Pers Soc Psychol* 53: 5-13.
- 500 19. Hoffman HG, Garcia-Palacios A, Patterson DR, Jensen M, Furness T, Ammons WF
501 (2001) The effectiveness of virtual reality for dental pain control: A case study.
502 *CyberPsychol Behav* 4: 527-535.
- 503 20. Frere CL, Crout R, Yorty J, McNeil DW (2001) Effects of audiovisual distraction during
504 dental prophylaxis. *J Am Dent Assoc* 132: 1031-1038.
- 505 21. Furman E, Jasinevicius TR, Bissada NF, Victoroff KZ, Skillicorn R, Buchner M (2009)
506 Virtual reality distraction for pain control during periodontal scaling and root planing
507 procedures. *J Am Dent Assoc* 140: 1508-1516.
- 508 22. Beukeboom CJ, Langeveld D, Tanja-Dijkstra K (2012) Stress-reducing effects of real and
509 artificial nature in a hospital waiting room. *J Altern Complement Med* 18: 329-333.
- 510 23. Dijkstra K, Pieterse ME, Pruyn A (2008) Stress-reducing effects of indoor plants in the
511 built healthcare environment: The mediating role of perceived attractiveness. *Prev Med* 47:
512 279-283.
- 513 24. Wheeler B, White MP, Stahl-Timmins W, Depledge MH (2012) Does living by the coast
514 improve health and wellbeing? *Health Place* 18: 1198-1201.
- 515 25. Volker S, Kistemann T (2011) The impact of blue space on human health and well-being
516 – Salutogenic health effects of inland surface waters: A review. *Int J Hyg Environ Health*
214: 449-460.

- 517 26. Krans J, Näring G, Holmes EA, Becker ES (2010) "I see what you're saying": Intrusive
518 images from listening to a traumatic verbal report. *J Anxiety Disord* 24: 134-140.
- 519 27. Holmes EA, Brewin CR, Hennessey RG (2004) Trauma films, information processing, and
520 intrusive memory development. *J Exp Psychol Gen* 133: 3-22.
- 521 28. Knight JF, Stone RJ, Qian C Virtual restorative environments: preliminary studies in
522 scene, sound and smell. *International Journal of Gaming and Computer-Mediated*
523 *Simulations* 2012, 4:In Press.
- 524 29. Humphris GM, Morrison T, & Linsay SJE (1995) The Modified Dental Anxiety Scale:
525 validation and United Kingdom norms. *Community Dent Health*12:143-150.
- 526 30. Mehrabian A, Russell JA (1974) *An approach to environmental psychology*. M.I.T. Press,
527 Cambridge, Mass.
- 528 31. Schubert T, Friedmann F, Regenbrecht H (2001) The experience of presence: Factor
529 analytic insights. *Presence* 10:266-281.
- 530 32. Baños R M, Botella C, Garcia-Palacios A, Villa H, Perpina C, Alcaniz M (2000) Presence
531 and reality judgment in virtual environments: A unitary construct? *CyberPsychol Behav* 3:
532 327-335.
- 533 33. Statham DJ, Connor JP, Kavanagh DJ, Feeny GFX, Young RMD, May J, Andrade J
534 (2011) Measuring alcohol craving: development of the Alcohol Craving Experience
535 Questionnaire. *Addict*106: 1230-1238.
- 536 34. De Jongh A, ter Horst G (1995) Dutch students' dental anxiety and occurrence of thoughts
537 related to treatment. *Community Dent Oral Epidemiol* 23: 170-172.
- 538 35. Derakshan N, Eysenck MW (2009) Anxiety, processing efficiency, and cognitive
539 performance. *New developments from attentional control theory*. *Eur Psychol* 14: 168-176.

- 540 36. Law EF, Dahlquist LM, Sil S, Weiss KE, Herbert LJ, Wohlheiter K, Horn SB (2011)
541 Videogame distraction using virtual reality technology for children experiencing cold
542 pressor pain: The role of cognitive processing. *J Pediatr Psychol* 36: 84-94.
- 543 37. Schneider SM, Ellis M, Coombs WT, Shonkwiler EL, Folsom LC (2003) Virtual reality
544 intervention for older women with breast cancer. *CyberPsychol Behav* 6: 103-107.
- 545 38. Hoffman HG, Doctor JN, Patterson DR, Carrougher GJ, Furness III, TA (2000) Virtual
546 reality as an adjunctive pain control during burn wound care in adolescent patients. *Pain*
547 85: 305-309.
- 548 39. Schmitt YS, Hoffman HG, Blough DK, Patterson DR, Jensen MP, Soltani M, Carrougher
549 GJ, Nakamura D, Sharar R (2011) A randomized, controlled trial of immersive virtual
550 reality analgesia, during physical therapy for pediatric burns. *Burns* 37: 61-68.
- 551 40. Mühlberger A, Wieser MJ, Kenntner-Mabiala R, Pauli P, Wiederhold BK. (2007) Pain
552 modulation during drives through cold and hot virtual environments. *CyberPsychol Behav*
553 10: 516-522.
- 554 41. Hartig T, Böök A, Garvill J, Olsson T, Gärling T (1996) Environmental influences on
555 psychological restoration. *Scand Journal Psychol* 37: 378-393.
- 556

557 **Figure legends**

558

559 Figure 1. Flowchart of participants

560 Figures 2 and 3. Set-up of the study. The person depicted in the images has given written
561 informed consent, as outlined in the PLOS consent form, to publication of their photograph.

562

563 Figure 4. Screenshot of the VR environment

564 Figure 5. The interaction effect of VR and dental anxiety on feelings of presence

565 Figure 6. The interaction effect of VR and dental anxiety on vividness of memories

Table 1. Overview of the means and standard deviations (between brackets) for the dependent variables

DV	Active VR (n=22)	Passive VR (n=23)	No VR control (n=24)
Perceived control	3.94 (1.57)	3.13 (1.20)	N/A
Compliance	3.67 (.66)	3.95 (.21)	3.57 (.66)
Low dental anxiety	3.60 (.70)	4.00 (.00)	3.71 (.47)
High dental anxiety	3.73 (.65)	3.90 (.32)	3.33 (.87)
Time perception (ratio)	1.33 (.50)	1.24 (.48)	1.31 (.57)
Low dental anxiety	1.12 (.40)	1.22 (.55)	1.26 (.54)
High dental anxiety	1.51 (.51)	1.35 (.46)	1.39 (.63)
Presence	6.21 (1.51)	5.16 (1.65)	N/A
Low dental anxiety	5.43 (1.04)	5.27 (1.44)	
High dental anxiety	6.86 (1.57)	4.92 (1.89)	
Awareness of the surrounding environment	4.05 (2.36)	4.61 (2.21)	5.17 (2.48)
Low dental anxiety	4.10 (2.23)	4.85 (2.15)	4.73 (2.71)
High dental anxiety	4.00 (2.56)	4.45 (2.30)	5.89 (1.97)
Interest in using VR during real dental visit	8.59 (1.94)	8.09 (2.41)	6.92 (2.78)
Low dental anxiety	7.90 (2.28)	7.31 (2.96)	6.47 (2.80)
High dental anxiety	9.17 (1.47)	8.82 (1.17)	7.67 (2.74)
Sample sizes for follow-up measures	n=21	n=22	n=19
Intrusive thoughts	1.68 (.98)	1.83 (1.26)	1.61 (1.25)
Low dental anxiety			
High dental anxiety			
Vividness of memories	4.26 (.88)	4.40 (1.40)	4.55 (2.23)
Low dental anxiety			
High dental anxiety			

