

2013-02

# Loss of SOX10 function contributes to the phenotype of human Merlin-null schwannoma cells

Doddrell, RDS

<http://hdl.handle.net/10026.1/3571>

---

10.1093/brain/aws353

Brain

Oxford University Press (OUP)

---

*All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.*

# Loss of SOX10 function contributes to the phenotype of human Merlin-null schwannoma cells

Robin D. S. Doddrell,<sup>1</sup> Xin-Peng Dun,<sup>1</sup> Aditya Shivane,<sup>1</sup> M. Laura Feltri,<sup>2</sup> Lawrence Wrabetz,<sup>2</sup> Michael Wegner,<sup>3</sup> Elisabeth Sock,<sup>3</sup> C. Oliver Hanemann<sup>1</sup> and David B. Parkinson<sup>1</sup>

1 Plymouth University Peninsula Schools of Medicine and Dentistry, Plymouth, Devon, PL6 8BU, UK

2 Hunter James Kelly Research Institute, School of Medicine and Biomedical Sciences, State University of New York at Buffalo, Buffalo NY 14203, USA

3 Institut für Biochemie, Emil-Fischer-Zentrum, Universität Erlangen-Nürnberg, Erlangen, Germany

Correspondence to: Dr David Parkinson,  
Peninsula School of Medicine and Dentistry,  
University of Plymouth,  
Devon, PL6 8BU, UK  
E-mail: david.parkinson@pms.ac.uk

Loss of the Merlin tumour suppressor causes abnormal de-differentiation and proliferation of Schwann cells and formation of schwannoma tumours in patients with neurofibromatosis type 2. Within the mature peripheral nerve the normal development, differentiation and maintenance of myelinating and non-myelinating Schwann cells is regulated by a network of transcription factors that include SOX10, OCT6 (now known as POU3F1), NFATC4 and KROX20 (also known as Egr2). We have examined for the first time how their regulation of Schwann cell development is disrupted in primary human schwannoma cells. We find that induction of both KROX20 and OCT6 is impaired, whereas enforced expression of KROX20 drives both myelin gene expression and cell cycle arrest in Merlin-null cells. Importantly, we show that human schwannoma cells have reduced expression of SOX10 protein and messenger RNA. Analysis of mouse SOX10-null Schwann cells shows they display many of the characteristics of human schwannoma cells, including increased expression of platelet derived growth factor receptor beta (PDGFRB) messenger RNA and protein, enhanced proliferation, increased focal adhesions and schwannoma-like morphology. Correspondingly, re-introduction of SOX10 into human Merlin-null cells restores the ability of these cells to induce KROX20 and myelin protein zero (MPZ), localizes NFATC4 to the nucleus, reduces cell proliferation and suppresses PDGFRB expression. Thus, we propose that loss of the SOX10 protein, which is vital for normal Schwann cell development, is also key to the pathology of Merlin-null schwannoma tumours.

**Keywords:** Schwann; merlin; SOX10; KROX20; schwannoma

**Abbreviations:** AMP = adenosine monophosphate; GFP = green fluorescent protein; P<sub>0</sub> = myelin protein zero; PDGF = platelet derived growth factor; MAPK = mitogen activated protein kinase; MBP = myelin basic protein

## Introduction

The genetic condition neurofibromatosis type 2 is characterized by formation of multiple tumours of the peripheral nervous system. The predominant tumours formed are vestibular

schwannomas of the eighth cranial nerve (vestibulocochlear), but also include cutaneous schwannomas, meningiomas and ependymomas (Hanemann, 2008). Neurofibromatosis type 2 is caused by biallelic loss of the *NF2* gene that encodes the tumour suppressor protein merlin (NF2), which is also lost in all sporadic schwannoma

Received August 9, 2012. Revised November 7, 2012. Accepted November 12, 2012

© The Author (2012). Published by Oxford University Press on behalf of the Guarantors of Brain.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

tumours (Rouleau *et al.*, 1993; Trofatter *et al.*, 1993). Research has shown that merlin acts to ensure cell–cell contact inhibition of growth and regulates protein ubiquitination in the nucleus through interaction with DCAF1 (McClatchey and Giovannini, 2005; Curto *et al.*, 2007; Lallemand *et al.*, 2009; Li *et al.*, 2010). Loss of merlin in cultured Schwann cells leads to altered morphology, enhanced proliferation and over activation of multiple signalling pathways including Rac GTPase, extracellular regulated kinase (ERK), c-Jun N-terminal kinase (JNK) and phosphatidylinositol-3 kinase (PI-3 kinase) MAP kinase pathways (Shaw *et al.*, 2001; Kaempchen *et al.*, 2003; Ammoun *et al.*, 2008). During Schwann cell development, a number of signalling pathways and transcription factors regulate axon–Schwann cell interaction, cell cycle withdrawal and differentiation of Schwann cells within peripheral nerves (Jessen and Mirsky, 2005; Pereira *et al.*, 2012), and the impact of merlin loss on these regulators of normal Schwann cell differentiation has not yet been investigated. The zinc finger transcription factor KROX20 is essential for the differentiation of myelinating Schwann cells and regulates both cell cycle withdrawal and myelin gene expression (Topilko *et al.*, 1994; Zorick *et al.*, 1999; Parkinson *et al.*, 2004). Expression of KROX20 in turn is regulated by a number of other transcription factors such as SOX10, OCT6, NFATC4 and YY1, which bind and co-operatively activate the myelinating Schwann cell enhancer of the *KROX20* promoter (Ghislain and Charnay, 2006; Kao *et al.*, 2009; He *et al.*, 2010; Kipanyula *et al.*, 2012). In addition to its role in myelin gene regulation, in synergy with KROX20, SOX10 functions to regulate Schwann cell proliferation and controls the dynamics of Schwann cell–axon interactions during development (Finzsch *et al.*, 2010). SOX10 is both essential for the generation of Schwann cell precursors from the neural crest (Britsch *et al.*, 2001) and the sequential induction of OCT6 and KROX20 during myelination (Finzsch *et al.*, 2010). Given that the key characteristics of human schwannoma tumours are loss of cell differentiation, loss of axonal contact and increased cell proliferation, we have investigated this transcriptional network in these cells.

As a first step, we tested both the regulation and function of KROX20 in human Merlin-null schwannoma cells. We found that the induction of both KROX20 and OCT6 is impaired in these cells, whereas enforced expression of KROX20 induces cell cycle arrest and differentiation identically in human control Schwann cells and Merlin-null schwannoma cells. As SOX10 is key in the regulation of both OCT6 and KROX20, we measured the levels of SOX10 in schwannoma cells and found reduced levels of *SOX10* messenger RNA and protein in all the tumours we analysed ( $n = 17$ ). In keeping with a role for SOX10 in these tumours, SOX10 re-expression restores the ability of these cells to induce Krox-20, relocalizes NFATC4 to the nucleus and drives myelin gene expression. Human schwannoma cells express raised levels of PDGFRB and consequently show increased signalling and proliferation in response to platelet derived growth factor (PDGF) (Ammoun *et al.*, 2008). We find that re-expression of SOX10 in human schwannoma cells reduces PDGFRB expression and consequently PDGF-induced signalling and proliferation. To separately study the roles of SOX10 and merlin, we have used conditional mouse knockouts for both genes together with small interfering RNA knockdown of merlin in Schwann cells. We demonstrate that loss of merlin alone, both

*in vitro* and *in vivo*, does not reduce SOX10 levels in Schwann cells. Similarly, restoration of merlin expression in human schwannoma cells does not rescue SOX10 expression. Analysis of SOX10-null mouse Schwann cells shows that loss of SOX10 alone appears to give many of the phenotypes of human schwannoma cells, namely increased levels of PDGFRB, increased cell proliferation, schwannoma-like morphology with increased focal adhesions and the failure to myelinate *in vitro* (Flaix *et al.*, 2009). These findings link, for the first time, the processes of normal cell differentiation and tumourigenesis in Schwann cells and demonstrate that loss of both merlin and SOX10 contribute to the aberrant phenotype of human schwannoma cells.

## Materials and methods

### Reagents

Adenoviruses expressing green fluorescent protein (GFP), KROX20/GFP, Cre recombinase, JNK-binding domain of JNK-interacting protein 1 (JIP-1) and  $\Delta 169$  c-Jun have been described elsewhere (Parkinson *et al.*, 2001, 2004, 2008). Adenoviruses expressing NF2/GFP (Xiao *et al.*, 2005) and dominant negative Rac1 (Wojciak-Stothard *et al.*, 2001) were kind gifts from J. Testa (Fox Chase Cancer Centre, Philadelphia, USA) and A. Ridley (King's College London, UK), respectively. The SOX10 adenoviral construct was generated using a C-Terminal haemagglutinin (HA)-tagged human SOX10, using the AdEasy™ system (He *et al.*, 1998) to generate an adenovirus co-expressing GFP and HA-tagged SOX10. Antibodies against OCT6 (sc-11661), NFATC4 (sc-13036), Rho GDI (sc-359), Beta2a tubulin (sc-134229) and merlin (sc-55575) were from Santa Cruz Biotechnology (Wembley, UK). Guinea pig and rabbit antibodies against SOX10 were as described (Stolt *et al.*, 2003; Maka *et al.*, 2005). Phospho-NF $\kappa$ B (Ser<sup>276</sup>) p65 antibody (#3037), phospho-CREB (Ser<sup>133</sup>, #9198) and PDGFRB (#3169) were obtained from New England Biolabs. S100 $\beta$  antibodies were from Dako, Ki67 antibodies from Abcam, KROX20 antibodies were from Cambridge Bioscience, phospho-ERK1/2 (V8031) antibody was from Promega and paxillin (5H11) antibody from Millipore. Antibody to periaxin was a gift from P. Brophy (University of Edinburgh, UK), and antibody to myelin protein zero (MPZ or P<sub>0</sub>) a gift from J. Archelos (Medical University of Graz, Austria). Biotinylated anti-guinea pig, anti-goat, anti-mouse and anti-rabbit were obtained from Vector laboratories. Alexa Fluor® fluorescently conjugated anti-rabbit, mouse or goat IgGs and fluorescently conjugated streptavidin were from Life Technologies (Paisley, UK). Superfrost glass slides were from VWR Merck. All reagents for small interfering RNA knockdown of merlin were obtained from Qiagen (Crawley, UK).

### Preparation of Schwann cell cultures

Schwann cells were prepared from the sciatic nerve and brachial plexus of post-natal day 3 rodents as previously described (Brookes *et al.*, 1979). Human Schwann cells were isolated from nerve donors and schwannoma cells from tumour samples of patients with neurofibromatosis type 2 (following informed consent). Neurofibromatosis type 2 diagnosis was established using criteria set down in the NIH Consensus Conference (National Institutes of Health Consensus Development Conference, 1988; Mulvihill *et al.*, 1990). Cells were harvested and cultured as previously described (Hanemann *et al.*, 1998; Rosenbaum *et al.*, 1998). All experiments were performed in

serum-free supplemented medium known as defined medium (Jessen *et al.*, 1994; Parkinson *et al.*, 2008).

## Immunocytochemistry, western blotting, polymerase chain reaction and adenoviral infections

Immunostaining and counting were performed as previously described (Parkinson *et al.*, 2001, 2003, 2004, 2008). Anti-SOX10 (both antibodies at 1:1000), anti-OCT6 (1:150), anti phospho-NF $\kappa$ B (Ser<sup>276</sup>) p65 (1:100), phospho-CREB (1:100), NFATC4 (1:1000) and PDGFRB (1:100) primary antibodies were incubated overnight at 4°C. Biotinylated secondary antibodies (1:200), anti-guinea pig, anti-goat and anti-rabbit, were incubated at room temperature for 45 min and the streptavidin-568 third layer (1:500) incubated for 30 min at room temperature. Confocal imaging was performed using the Zeiss meta 510 confocal microscope and the Zeiss LSM software (Zeiss). Fluorescence microscopy was performed on a Nikon Eclipse 80i. Unless otherwise stated, western blotting, PCR and adenoviral infections were performed as previously described (Parkinson *et al.*, 2001, 2003, 2004, 2008; Kaempchen *et al.*, 2003; Flaiz *et al.*, 2007). For semi-quantitative PCR, the following primers were used: 18S rRNA For, CAGCCACCCGAGATTGAGCA, Rev, TAGTAGCGACGGCGGTG TG; SOX10 For: CAGGCGAGCTGGGCAAGGTCAAG, Rev: CTGCCT GAGCCCACACCATGAAG; PDGFRB For: CGAGTTGGACCTGAACA TGAC, Rev: CGCACAATCTCGATCTTTCTC; CD44 For: TGAATATAA CCTGCCGCTTTG, Rev: GTCATACTGGGAGGTGTTGGA.

## Schwann cell differentiation and proliferation assays

For myelination assays, defined medium was supplemented with 1 mM dibutyryl cAMP for 48 h unless otherwise stated. Human Schwann and schwannoma cells were infected with adenoviruses, in defined medium without forskolin, for 24 h before cAMP stimulation in fresh defined medium. For proliferation assays in human schwannoma cells, 20 ng/ml NRG1, 100 ng/ml IGF-1 or 100 ng/ml PDGF-DD was added to fresh defined medium for 24 h or 72 h after the initial adenoviral infection. For proliferation assays in mouse Schwann cells defined medium was supplemented with 10 ng/ml PDGF-DD with or without 2  $\mu$ M forskolin for 48 h.

## Transgenic mice

Transgenic mouse breeding and experiments were performed following guidelines from the UK Home Office. Mice with the conditional SOX10<sup>fl/fl</sup> allele have been described (Finzsch *et al.*, 2010), and mice with the conditional NF2<sup>fl/fl</sup> allele (Giovannini *et al.*, 2000) were obtained from the Riken BioResource Centre (Experimental Animal Division, Japan). For *in vitro* analysis of null cells, mouse Schwann cells were prepared from the sciatic nerves of either Sox10<sup>fl/fl</sup> or Nf2<sup>fl/fl</sup> mouse pups and SOX10 or merlin gene expression removed in culture by infection with Cre-recombinase expressing adenovirus. Schwann cells from the same culture were infected with GFP virus and used as a control. Mice expressing Cre recombinase under the control of the P<sub>0</sub> promoter (mTOTA P<sub>0</sub>-Cre) have been previously described (Feltri *et al.*, 1999). For analysis of Merlin-null sciatic nerve *in vivo*, NF2<sup>fl/fl</sup> mice were crossed with P<sub>0</sub>-Cre mice to generate NF2<sup>fl/wt</sup> Cre+ animals that were then back-crossed to NF2<sup>fl/fl</sup> mice to generate control (NF2<sup>fl/fl</sup> Cre-) and Merlin-null (NF2<sup>fl/fl</sup> Cre+) sciatic nerve for analysis.

## Small interfering RNA knockdown

Merlin was knocked down in Schwann cells using 3 nM of validated small interfering RNA oligos Rn\_Nf2\_2 and Rn\_Nf2\_3 in combination. Alternatively, 3 nM Allstars negative control was used as a scrambled control. Small interfering RNA/lipid complexes were prepared using 6  $\mu$ l of HiPerfect® reagent and incubated with cell cultures for 20 h. Transfection was performed twice at 0 and 48 h, and protein expression assayed a further 24 or 48 h later.

## Fluorescent immunohistochemistry

Transverse 7- $\mu$ m thick cryostat sections of schwannoma and control peroneal nerve were fixed with 4% paraformaldehyde on Superfrost plus glass slides. Sections were then stained sequentially with antibodies against SOX10 (1:1000) and S100 $\beta$  (1:100) following the fluorescent immunocytochemistry protocol.

## Scanning electron microscopy

For scanning electron microscopy, mouse Schwann cells were cultured on glass coverslips and infected with GFP or Cre expressing adenovirus as above. Forty-eight hours after addition of adenovirus, coverslips were fixed in 2.5% w/v glutaraldehyde, diluted in 0.2 M sodium cacodylate buffer, at room temperature for 2 h and then washed twice in 0.2 M sodium cacodylate buffer. Coverslips were washed in 30%, 50%, 70%, 90% and 100% absolute ethanol for 15 min each at room temperature. Samples then underwent critical point drying through CO<sub>2</sub> in a K850 Critical Point Dryer (Emitech), and coverslips were mounted on support stubs and gold coated (nominal thickness 10 nm) in a K550 Sputter Coater (Emitech). Samples were imaged on a JSM-5600LV scanning electron microscope (JEOL Ltd.) operated at 15 kV, with an 11 mm working distance and at  $\times$ 1000 or  $\times$ 2000 magnification.

## Statistics

For all experiments,  $n = 3$  was used unless otherwise stated. All graphs display the arithmetic mean with error bars representing one standard error of the mean.  $P$ -values were calculated using a Student  $t$ -test and are denoted on figures as: \* $P < 0.05$ , \*\* $P < 0.01$  and \*\*\* $P < 0.005$ . For all cell differentiation and proliferation assays,  $\sim$ 200 cells were counted in duplicate. In adenoviral experiments, the number of positive cells was divided by the number of GFP positive cells. For all other experiments, the number of positive cells was divided by the number of Hoechst positive cells. A minimum of 500 cells were counted for SOX10 positivity in each cryostat section.

## Results

### KROX20 drives myelin gene expression in Merlin-null schwannoma cells

It has been well characterized that KROX20 is the key regulator of Schwann cell myelination. Enforced expression of KROX20 *in vitro* is sufficient to drive increased expression of compact myelin proteins (P<sub>0</sub> and MBP), myelin associated proteins (myelin associated glycoprotein and periaxin) and essential enzymes in myelin lipid synthesis (Nagarajan *et al.*, 2001; Parkinson *et al.*, 2003).



KROX20 expression also downregulates expression of the myelination inhibitory transcription factors *c-Jun* and *Sox2* (Parkinson *et al.*, 2004, 2008; Le *et al.*, 2005). As a first step in this analysis of Merlin-null schwannoma cells, we tested whether these events downstream of KROX20 could be normally regulated in these tumour cells. No significant basal differences in the expression of myelin proteins or negative myelin-regulators were observed in schwannoma cells. To investigate whether KROX20 could drive myelination in schwannoma cells, we infected primary human schwannoma (NF2<sup>-/-</sup>) and control Schwann (NF2<sup>+/+</sup>) cells with adenoviral constructs co-expressing KROX20 and GFP or GFP alone. After KROX20 expression, we assessed, by immunolabelling and western blotting, the induction of myelin targets and downregulation of *c-Jun* expression.

KROX20 was able to drive strong upregulation of the myelin protein P<sub>0</sub> and the myelin marker periaxin in both wild-type Schwann cells and Merlin-null schwannoma cells, shown by both fluorescent immunocytochemistry and western blotting (all  $P < 0.02$ ). Similarly, KROX20 was also able to downregulate the inhibitory transcription factor *c-Jun* in Merlin-null schwannoma cells ( $P < 0.001$ ) (Fig. 1). The regulation of P<sub>0</sub>, periaxin and *c-Jun* by KROX-20 in human Schwann and schwannoma cells was indistinguishable from that seen in primary rat Schwann cells (data not shown). These data suggest that once expressed, KROX-20 is apparently fully able to drive the downstream myelination programme in Merlin-null schwannoma cells.

## KROX20 expression inhibits the proliferation of Merlin-null schwannoma cells

In addition to controlling myelin gene expression, KROX20 has been shown to regulate the proliferation of Schwann cells, inhibiting the proliferation of cells in response to mitogens such as beta-neuregulin (NRG1) (Zorick *et al.*, 1999; Parkinson *et al.*, 2004). Loss of merlin causes schwannoma cells to proliferate at an increased basal rate, to lose cell–cell contact dependent inhibition of cell proliferation and continue to grow in overlapping layers past confluency *in vitro* (Lallemand *et al.*, 2003; Curto *et al.*, 2007; Flaiz *et al.*, 2007, 2008). This proliferation could be driven by either PDGF or IGF-1, which have both been shown to be mitogenic for schwannoma cells *in vitro* (Ammoun *et al.*, 2008, 2012).


To test whether KROX20 effectively controls proliferation in schwannoma cells, primary human Merlin-null cells were infected with either GFP- or KROX20/GFP-expressing adenoviruses in defined medium and then treated with the Schwann cell mitogens NRG1 (10 ng/ml), PDGF-DD (100 ng/ml) or IGF-1 (100 ng/ml). Proliferation was measured by immunofluorescence using an antibody against Ki-67, which has a punctate nuclear stain in proliferating cells (Fig. 2). Basal proliferation of schwannoma cells in defined medium alone was  $2.3 \pm 0.5\%$ , which was increased by NRG1, PDGF or IGF-1 addition to  $8.1 \pm 1.1$ ,  $8.7 \pm 1.1$  and  $11.2 \pm 2.1\%$ , respectively, in control GFP infected schwannoma cells (Fig. 2). This proliferation was dramatically and significantly reduced to  $<1\%$  with all mitogens in KROX20-expressing

schwannoma cells (Fig. 2; NRG1,  $P < 0.001$ ; PDGF,  $P < 0.001$ ; IGF-1,  $P = 0.002$ ;  $n = 4$ ). KROX20 therefore effectively controls proliferation of schwannoma cells in mitogen-stimulated conditions. In conclusion, we saw that enforced KROX20 expression in Merlin-null schwannoma cells is sufficient to drive myelin gene expression, downregulate inhibitors of myelination and to inhibit the proliferation of schwannoma cells in response to a range of mitogens. These findings implied that the differentiation network downstream of KROX20 is intact in Merlin-null schwannoma cells; therefore, we next investigated the regulatory events upstream of KROX20 in these cells.


## Impaired induction of KROX20 and OCT6 in schwannoma cells

During Schwann cell myelination *in vivo*, expression of KROX20 is controlled by a complex signalling cascade involving the transcription factors OCT6, SOX10, YY1 and NFATC4, as well as through activation of the G-protein coupled receptor GPR126 (Topilko *et al.*, 1994; Jaegle *et al.*, 1996; Kao *et al.*, 2009; Monk *et al.*, 2009, 2011; Finzsch *et al.*, 2010; He *et al.*, 2010). This process of myelination can be mimicked *in vitro* by addition of cyclic AMP, which causes Schwann cell flattening and upregulation of myelin proteins (e.g. P<sub>0</sub>, myelin basic protein and periaxin), myelin lipids (e.g. O4) and myelinating transcription factors (e.g. OCT6 and KROX20) (Morgan *et al.*, 1991; Parkinson *et al.*, 2008). The ability of Merlin-null schwannoma cells to upregulate KROX20 in response to cyclic AMP was tested by incubating primary human Schwann and schwannoma cells in defined medium alone or in defined medium supplemented with 1 mM of the non-hydrolysable cyclic AMP analogue dibutyryl-cAMP (hereafter referred to as cAMP) for 24, 48 and 72 h and examining KROX20 induction by immunofluorescence and western blotting. Human control Schwann cells displayed no detectable KROX20 expression in basal conditions, but dramatically induced KROX20 on cAMP treatment, peaking at  $34.8 \pm 4.2\%$  of nuclei positive for KROX20 at 72 h (Fig. 3 at 48 h and Supplementary Fig. 1). In contrast, Merlin-null schwannoma cells from seven independent tumours were significantly reduced in their ability to induce KROX20 protein at all time points examined (Fig. 3 and data not shown) peaking at only  $9.1 \pm 2.3\%$  at 72 h, with a significantly decreased induction at 24 h ( $P = 0.037$ ), 48 h ( $P < 0.001$ ) and 72 h ( $P < 0.001$ ). Schwannoma cells from three of these tumours displayed an absolute block in KROX20 induction, with  $<1\%$  of cells KROX20 positive after any duration of cAMP treatment. This result was confirmed by western blotting at the 48 h time point in control human Schwann and schwannoma cells, again showing no apparent induction of KROX20 in Merlin-null schwannoma cells from a further two schwannoma tumours (Fig. 3). The myelinating Schwann cell marker periaxin is also induced by cAMP in Schwann cells (Parkinson *et al.*, 2003) and we found that the induction of periaxin was also significantly blocked in Merlin-null schwannoma cells at all timepoints examined (Fig. 3, 48 h and Supplementary Fig. 1,  $P < 0.001$ , 72 h).

KROX20 induction may be inhibited by increased activity of the ERK1/2 and JNK1/2 MAP kinase pathways (Harrisingh *et al.*,



**Figure 1** Krox-20 induces periaxin and P<sub>0</sub> and downregulates c-Jun expression in both control and Merlin-null human Schwann cells. (A–H) Immunofluorescence of control Schwann +/+ (A, B, E and F) and Merlin-null schwannoma -/- (C, D, G and H) cells infected with control GFP (A–D) or GFP/KROX20 (E–H, K20) expressing adenoviruses showing equal induction of periaxin (Prx) protein in both control and Merlin-null cells (F and H). (I–P) Immunofluorescence of control +/+ (I, J, M and N) and Merlin-null -/- (K, L, O and P) cells infected with GFP and GFP/KROX20 (K20) expressing adenoviruses, showing down regulation of c-Jun in both control and Merlin-null cells (N and P). Scale bars = 20  $\mu$ m. (Q and R) Graphs showing percentage periaxin/GFP (Q) and c-Jun/GFP (R) positive control Schwann (+/+) and schwannoma (-/-) cells following infection with GFP control and GFP/KROX20 expressing adenoviruses. (S) Western blot showing similar upregulation of periaxin and P<sub>0</sub> protein and downregulation of c-Jun expression by KROX20 in both control Schwann (+/+) and Merlin-null schwannoma cells (-/-).



**Figure 2** KROX20 halts the proliferation of Merlin-null schwannoma cells. (A–D) Human Merlin-null schwannoma cells were infected with GFP (A and B) or GFP/KROX20 (K20) adenoviruses (C and D) for 24 h and then treated with 20 ng/ml of NRG-1 (+ NRG-1) for a further 24 h. Cells were fixed and immunolabelled with antibodies against Ki-67. Arrows in A and B indicate Ki-67 positive nuclei. Scale bar = 20  $\mu$ m. (E) Graph showing that KROX20 expression causes a significant decrease in schwannoma cell proliferation in response to NRG-1 ( $P < 0.001$ ), PDGF ( $P < 0.001$ ) or IGF-1 ( $P = 0.002$ ).

2004; Ogata *et al.*, 2004), and both pathways have been shown to be elevated in Merlin-null schwannoma cells (Ammoun *et al.*, 2011). We therefore tested whether inhibition of either or both of these pathways would restore KROX20 induction. Experiments using a JNK1/2 inhibitor (SP600125), expression of the JNK-binding domain of JIP-1, both of which have previously been shown to inhibit JNK activity in Schwann cells (Parkinson *et al.*, 2004), or even expression of dominant negative c-Jun ( $\Delta 169$  c-Jun; Parkinson *et al.*, 2001) did not restore KROX20 induction in schwannoma cells. Similarly, inhibition of the MEK/ERK pathway by U0126 or AZD6244, the PI3-kinase pathway using BEZ235, or even use of the ErbB2 inhibitor Lapatinib or the PDGFR inhibitors Nilotinib or Sorafenib, all of which inhibit schwannoma cell proliferation at the concentrations used, were unable to increase KROX20 induction in response to cAMP in schwannoma cells (Supplementary Fig. 2 and data not shown). Similarly, both mouse and human Merlin-null cells show increased GTP loading of Rac, and such enhanced Rac activity has been shown to block elongation of schwannoma cells along dorsal root ganglion axons in co-culture assays, an early step in myelination (Nakai *et al.*, 2006). Inhibition of Rac activity by adenovirally mediated expression of dominant negative N17-Rac also failed to rescue KROX20 expression in schwannoma cells in response to cAMP (data not shown).

Next, we tested whether there was normal induction of the POU domain transcription factor OCT6 in Merlin-null schwannoma cells. Forty-eight hours after addition of cAMP to control human Schwann cells, we observed induction of nuclear OCT6 by


immunocytochemistry and western blotting, which was significantly ( $P = 0.003$ ) blocked in Merlin-null schwannoma cells (Fig. 3).

Phosphorylation and activation of NF $\kappa$ B and CREB following cAMP treatment are thought to be important for the induction of OCT6 and KROX20, respectively (Nickols *et al.*, 2003; Arthur-Farraj *et al.*, 2011). Using phospho-specific antibodies for the p65 subunit of NF $\kappa$ B (Ser276) and CREB (Ser133), we measured the response to cAMP in control and Merlin-null schwannoma cells. Analysis by immunocytochemistry with these antibodies after cAMP addition showed an identical amplitude and duration of activation (i.e. phosphorylation) of NF $\kappa$ B and CREB in control Schwann and Merlin-null schwannoma cells (Supplementary Fig. 3). Additionally, the zinc finger transcription factor YY1 has also been shown to regulate KROX20 expression during myelination. Analysis by immunolabelling showed clear nuclear YY1 expression in both control Schwann and Merlin-null schwannoma cells (data not shown). These experiments identify the regulation of both OCT6 and KROX20 by cyclic AMP as abnormal in Merlin-null schwannoma cells.

## SOX10 levels are reduced in Merlin-null schwannoma cells *in vitro* and *in vivo*

The high mobility group transcription factor SOX10 is first expressed in the neural crest and is required for Schwann cell specification and subsequent myelination in the peripheral nervous system (Britsch *et al.*, 2001; Finzsch *et al.*, 2010). The control of both KROX20 and OCT6 have been shown to be dependent on






**Figure 3** Merlin-null schwannoma cells do not induce OCT6 or KROX20 in response to cyclic AMP. (A–F) Impaired induction of OCT6 in schwannoma cells. Control (NF2 +/+) and Merlin-null (NF2 -/-) cells were treated for 48 h with 1 mM cAMP and OCT6 levels measured by immunolabelling (A–D) and western blotting (F). Note clear induction of OCT6 in nuclei of control (+/+) cells (C), whereas no OCT6 is observed in Merlin-null (-/-) cells (D). Scale bar = 20  $\mu$ m (E). Graph of percentage OCT6-positive Schwann and schwannoma cells in response to cAMP. (F and G) Western blot showing lack of OCT6 (F) and KROX20 (G) induction in response to cAMP at 48 h. (H and I) Graphs showing percentage KROX20 (H) and periaxin (I) positive control (NF2 +/+) and Merlin-null (NF2 -/-) cells after cAMP treatment (48 h). DM = defined medium control.

SOX10 expression (Ghislain and Charnay 2006; Reiprich *et al.*, 2010; Jagalur *et al.*, 2011); therefore, we next measured SOX10 levels *in vitro* in control and Merlin-null schwannoma cells and *in vivo* by analysis of cryostat sections of control human nerve and schwannoma tumour samples. Figure 4 shows the results of immunocytochemistry and western blotting of human Schwann and schwannoma cells. *In vitro*, control Schwann cells displayed a uniform intense nuclear expression in all cells. In contrast, both schwannoma cells *in vitro* and sections of schwannoma tumours showed a consistent decreased expression in all cells or a heterogeneous staining pattern with some cells expressing SOX10 and some containing no detectable SOX10 ( $n = 8$ ; Fig. 4). These findings were confirmed with two separate SOX10 antibodies. In total, we analysed cultured cells from eight schwannoma tumours and

cryostat sections from a further six independent tumours and observed a significant reduction of SOX10 in all samples analysed. Counts of SOX10-positive nuclei in control nerve and schwannoma sections showed overall a significant (60.3%,  $P < 0.001$ ) decrease in the percentage of SOX10-positive nuclei (Supplementary Fig. 4). Western blotting of cell extracts from three further independent schwannoma cell cultures compared with three control human primary Schwann cell extracts also showed a variable, but consistent, reduction in SOX10 expression in Merlin-null schwannoma cells. Thus, in total, we have analysed 17 human schwannoma samples, all of which showed a reduction in SOX10 protein levels. Similarly, we observed a reduction of SOX10 messenger RNA levels in schwannoma cells compared with control Schwann cells (Fig. 4).





**Figure 4** Expression of SOX10 protein and messenger RNA is reduced in Merlin-null human schwannoma cells. (A–F) Cryostat sections of control human peroneal nerve (+/+), (A and B) and human schwannoma tumours (–/–, C–F) stained with SOX10 antibody and the Schwann cell marker S100 $\beta$  (S100). Nuclei are counterstained with Hoechst dye (Ho). Almost all nuclei in control nerve (A and B) are SOX10 positive. For the tumour shown in panels C and D, a mosaic pattern is observed with many cells being SOX10 negative. For the tumour shown in panels E and F, almost no SOX10 stain is observed. Scale bar = 20  $\mu$ m. (G–J) Immunolabelling of dissociated control Schwann (+/+), (G and H) and Merlin-null schwannoma (–/–, I and J) cells in culture with SOX10 antibody. Note mosaic pattern in SOX10 stain in panel J; arrows show SOX10-negative cells in I and J. Scale bar = 20  $\mu$ m. (K) Western blot analysis of three independent human control (+/+) and Merlin-null (–/–) schwannoma tumour samples. Note reduction of SOX10 expression in all tumour samples. (L) Semi-quantitative PCR showing reduction of SOX10 messenger RNA in two independent Merlin-null tumours (–/–) compared with normal Schwann cell controls (+/+).

## Re-expression of SOX10 in human Merlin-null schwannoma cells rescues KROX20 and P<sub>0</sub> induction

Having identified a consistent reduction in SOX10 levels in schwannoma cells as a possible cause of decreased KROX20 and OCT6, we measured the effects of haemagglutinin-tagged SOX10 re-expression in human schwannoma cells. Immunolabelling and western blotting of schwannoma cells with either haemagglutinin or SOX10 antibodies showed strong nuclear expression of SOX10 in cells following adenoviral infection (Fig. 5 and data not shown). Next, we tested whether reintroduction of SOX10 into schwannoma cells would allow induction of Krox-20 in response to cAMP. SOX10 expression was sufficient, in the absence of cAMP, to induce KROX20 in a small percentage of schwannoma cells ( $P=0.001$  versus controls), which was greatly enhanced by the addition of cAMP ( $P < 0.001$  versus controls). In addition, we also observed the induction of P<sub>0</sub> protein in these cells (Fig. 5).


## SOX10 re-expression causes nuclear localization of NFATC4 in schwannoma cells

Signalling through the calcineurin/NFAT pathway has also recently been shown to be required for KROX20 induction and myelination

(Kao *et al.*, 2009; Kipanyula *et al.*, 2012). Although Schwann cells express NFATC1–C4 (Kipanyula *et al.*, 2012), only NFATC4 has been shown to bind to the myelinating Schwann cell enhancer of the KROX20 gene and drive KROX20 expression in Schwann cells (Kao *et al.*, 2009). We therefore examined the levels and localization of NFATC4 in schwannoma cells infected with control GFP and SOX10/GFP expressing adenoviruses with or without cAMP treatment. This analysis showed that, in contrast to control GFP virus-infected cells, SOX10 reintroduction causes nuclear localization of NFATC4 that is further boosted upon cAMP treatment ( $P < 0.001$ ). Western blotting demonstrated that this nuclear localisation occurred without any SOX10 driven increase in NFATC4 protein expression (Fig. 5). In addition to this, expression of SOX10 also seems to cause a shift in the molecular weight of NFATC4, which is consistent with its potential dephosphorylation, and nuclear localization, in response to SOX10 expression (Kao *et al.*, 2009).

## Loss of SOX10 expression in schwannoma cells is independent of Merlin loss

Having observed the clear loss of SOX10 in schwannoma cells and the effects of SOX10 re-expression in these cells, we next tested whether SOX10 expression is directly controlled by merlin in




**Figure 5** Re-introduction of SOX10 into Merlin-null human schwannoma cells rescues induction of KROX20, P<sub>0</sub> and NFATC4. (A–H) Merlin-null cells were infected with GFP (A–D) and GFP/SOX10 (E–H) expressing adenoviruses and either maintained in defined medium alone (A, B, E and F) or defined medium plus 1 mM cAMP (C, D, G and H). Note that SOX10 expression alone is sufficient to drive KROX20 expression in a small number of cells (F, I and K), which is further enhanced by cAMP treatment (H, I and K). Scale bar = 20 μm. (I) Percentage KROX20-positive cells in GFP- and SOX10-infected schwannoma cells. (J) Graph showing that SOX10-null mouse Schwann cells do not induce KROX20 in response to cAMP + NRG-1, but re-expression of SOX10 restores KROX20 induction. (K) Western blot showing regulation of KROX20 and P<sub>0</sub> by SOX10 in Merlin-null cells. Probing with HA antibody detects adenovirally mediated SOX10 expression. (L–S) Re-expression of SOX10 in Merlin-null cells restores nuclear localization of NFATC4. Merlin-null cells were infected with GFP- (L–O) and GFP/SOX10- (P–S) expressing adenoviruses and either maintained in defined medium alone (L, M, P and Q) or defined medium plus 1 mM cAMP (N, O, R and S). Scale bar = 20 μm. (T) Percentage NFATC4-positive cells in GFP- and GFP/SOX10-infected schwannoma cells. (U) Graph showing that SOX10-null mouse Schwann cells do not induce nuclear localization of NFATC4 in response to cAMP + NRG-1, but re-expression of SOX10 restores NFATC4 relocalization. (V) Western blot showing expression of NFATC4 in GFP and GFP/SOX10 infected schwannoma cells. DM = defined medium.

Schwann cells. To do this, we used a number of *in vitro* and *in vivo* approaches. Firstly, using small interfering RNA, we silenced merlin expression in rat Schwann cells and measured SOX10 expression by western blot and immunocytochemistry. No effect on SOX10 levels was seen at either 72 or 96 h time points after transfection, despite almost complete silencing of merlin (Fig. 6). Next, we analysed Merlin-null Schwann cells, either *in vitro* by infection of *NF2<sup>fl/fl</sup>* Schwann cells (Giovannini *et al.*, 2000) with adenovirus expressing Cre recombinase, or *in vivo* by analysis of Merlin-null sciatic nerve prepared by breeding *NF2<sup>fl/fl</sup>* mice with P<sub>0</sub>-Cre expressing mice to specifically ablate merlin expression in Schwann cells of the nerve (Feltri *et al.*, 1999). Using both paradigms, we did not observe a reduction of SOX10 protein expression in Merlin-null Schwann cells, either by

## Loss of SOX10 alone results in phenotypes typical of human schwannoma cells

Loss of SOX10 *in vivo* leads to abnormalities in Schwann cell–axon interactions, a complete block in myelination and increased post-natal proliferation of Schwann cells in peripheral nerves (Finzsch *et al.*, 2010; Bremer *et al.*, 2011). To characterize changes in Schwann cells that are SOX10-dependent, we next prepared, *in vitro*, SOX10-null Schwann cells by treatment of SOX10<sup>fl/fl</sup> cells with Cre-recombinase expressing adenovirus. *In vitro* SOX10-null cells show a flattened non-bipolar morphology and increased focal adhesions as measured by paxillin immunolabelling (Fig. 7), both



**Figure 6** Regulation of SOX10 expression in Schwann cells is independent of merlin. (A) Western blot of rat Schwann cells transfected with control scrambled (asterisk) or *NF2* small interfering RNA (NF2) at 72 h and 96 h after transfection; untransfected cells (–). Note that knockdown of merlin at both 72 and 96 h time points, but no decrease in SOX10 protein expression. (B) Western blot of *NF2<sup>fl/fl</sup>* mouse Schwann cells infected with control GFP- or Cre-expressing adenoviruses, 72 h after infection. (C) Western blot of control (*NF2<sup>fl/fl</sup>* Cre–) and Merlin-null (*NF2<sup>fl/fl</sup>* Cre+) post-natal Day 6 (P6) mouse sciatic nerve, showing no reduction of SOX10 protein levels. (D–G) Cryostat sections from control (+/+; D and E) and Merlin-null (–/–; F and G) post-natal Day 6 mouse sciatic nerve shows no difference in nuclear SOX10 stain (E and G); nuclei are counterstained with Hoechst dye (Ho; D and F). Scale bar = 100  $\mu$ m. Insets in panels D–G show higher magnification views of cells, confirming that SOX10 is localized to the nuclei of cells. (H–L) Re-introduction of merlin into Merlin-null human schwannoma cells does not restore expression of SOX10. Human schwannoma cells were infected with either GFP control- (H, I) or GFP/Merlin- (J and K) expressing adenoviruses and immunolabelled with SOX10 antibodies 48 h after infection (I and K). Scale bar = 20  $\mu$ m. (L) Western blot showing human schwannoma cells 48 h after infection with control GFP or merlin (NF2) expressing adenoviruses. Merlin re-introduction into Merlin-null human schwannoma cells does not increase SOX10 expression.

western blot or staining of nerve cryosections (Fig. 6), demonstrating that loss of merlin does not directly lead to lower SOX10 levels. Finally, we performed experiments to reintroduce merlin into human schwannoma cells and measure levels of SOX10. Adenovirally mediated re-expression of merlin did not increase SOX10 levels in cells after 48 h as measured by either immunocytochemistry or western blot in human schwannoma cells (Fig. 6).

phenotypes of human schwannoma cells (Rosenbaum *et al.*, 1998; Flaiz *et al.*, 2009). Similar to human schwannoma cells, SOX10-null cells show significantly increased ( $P = 0.03$ ) adhesion in a cell adhesion assay (Utermark *et al.*, 2003) compared with control GFP virus infected cells (Supplementary Fig. 5). To confirm our idea that loss of SOX10 in Merlin-null human schwannoma cells causes reduced KROX20 induction, we performed *in vitro* myelination assays as



described (Arthur-Farraj *et al.*, 2011) with control and *Sox10*-null mouse Schwann cells. *SOX10*-null Schwann cells displayed minimal NFATC4 nuclear localization and KROX20 induction that was significantly increased by *SOX10* reintroduction ( $P = 0.003$ , NFATC4;  $P < 0.001$ , KROX20) (Fig. 5). Similarly, in line with the *in vivo* evidence (Finzsch *et al.*, 2010), *SOX10*-null mouse Schwann cells fail to induce OCT6 in response to cAMP plus neurotrophin (data not shown).


In addition, human Merlin-null schwannoma cells show a range of phenotypes, but two of the key changes in these cells are increased expression of PDGFRB, causing increased proliferation of cells, and the CD44 cell surface glycoprotein, which is involved in cell adhesion and migration (Morrison *et al.*, 2001; Hanemann *et al.*, 2006; Bai *et al.*, 2007; Ammoun *et al.*, 2008). To study whether the regulation of PDGFRB and CD44 in Schwann cells is *SOX10*-dependent, we analysed their expression in control and *SOX10*-null mouse Schwann cells and the effects of *SOX10* reintroduction in Merlin-null human schwannoma cells. *SOX10*-null cells show higher expression levels of PDGFRB compared with control cells and also show significantly increased basal ( $P = 0.002$ ) and PDGF-induced ( $P < 0.001$ ) rates of proliferation (Fig. 7).

In agreement with this, reintroduction of *SOX10* into human schwannoma cells greatly reduces expression of PDGFRB and CD44 messenger RNA and protein. *SOX10* re-expression also significantly reduces basal proliferation ( $P = 0.03$ ) and both PDGF stimulated proliferation ( $P = 0.01$ ) and MAP kinase signalling in these cells (Fig. 8).

Schwannoma cells re-expressing *SOX10*, when viewed by fluorescence microscopy, showed a much more regular cell shape with fewer filopodia and appearing to grow over each other less than GFP control cells. Scanning electron microscopy of GFP- and GFP/*SOX10*-infected schwannoma showed that *SOX10* infected cells do not overlap as frequently as control cells, displaying greater respect for the cell membrane of neighbouring cells (Fig. 8). The small GTPase Rac is a well-established regulator of the actin cytoskeleton and cell morphology, including the formation of lamellopodia. Enhanced Rac activity is a key feature of Merlin-null murine and human schwannoma cells (Shaw *et al.*, 2001; Kaempchen *et al.*, 2003). Rac activation was measured in control GFP- and *SOX10*/GFP-infected human schwannoma cells using a Rac G-LISA kit that found *SOX10* re-expression caused a small but significant decrease (0.86-fold of control levels,  $P = 0.031$ ) in GTP bound Rac (Supplementary Fig. 6).

## Discussion


Roles for the merlin tumour suppressor have been characterized in many cell types and interaction of merlin with many other proteins regulates a diverse number of cell functions ranging from signalling at the cell membrane to controlling protein degradation in the nucleus (McClatchey and Giovannini 2005; Ammoun *et al.*, 2008; McClatchey and Fehon 2009; Li *et al.*, 2010; Cooper *et al.*, 2011). The hallmark feature of neurofibromatosis type 2 is the occurrence of Merlin-null vestibular schwannomas, often bilaterally, and currently there is no effective clinical treatment for these tumours (Hanemann, 2008). Whereas a lot is now known about the



**Figure 7** Loss of *SOX10* alone in mouse Schwann cells results in phenotypes typical of human schwannoma cells. (A) Western blot of *SOX10*<sup>fl/fl</sup> mouse Schwann cells infected with GFP- or Cre-expressing adenoviruses, showing increased expression of PDGFRB in *SOX10*-null mouse cells. (B and C) Scanning electron Microscopy showing flattening of *SOX10*-null cells (C) in comparison with control cells (B). Scale bar = 10  $\mu$ m. (D and E) Immunolabelling of control (D) and *SOX10* null (E) cells with paxillin antibody (Pax) to reveal increased numbers of focal adhesions in *SOX10* null cells (E). Scale bar = 20  $\mu$ m. (F) Graph showing significantly ( $P < 0.001$ ) increased proliferation of *SOX10*-null mouse Schwann cells in response to 10 ng/ml of PDGF or PDGF plus 2  $\mu$ M of forskolin (PDGF/For). Reintroduction of *SOX10* into *SOX10*-null cells once more decreases PDGF-induced proliferation.

transcription factors and signalling that control normal Schwann cell-axon interaction, cell cycle exit and differentiation, little is known about how merlin loss impacts on these processes and leads to tumour development. We have performed a number of experiments to examine the process of differentiation and cell





**Figure 8** Re-introduction of SOX10 reduces PDGFRB and CD44 levels, reduces PDGF-induced signalling and prevents cell overgrowth in human Merlin-null schwannoma cells. (A–D) GFP-control- (A and B) and GFP/SOX10- (C and D) infected human schwannoma cells were immunolabelled with PDGFRB antibodies. SOX10 causes clear downregulation of membrane staining for PDGFRB. Scale bar = 20  $\mu$ m. (E) Western blot showing that adenovirally mediated re-expression of SOX10 lowers PDGFRB protein levels in schwannoma cells. (F) Semi-quantitative PCR- of duplicate complementary DNA samples prepared from two independent human schwannoma cell cultures infected with control GFP or GFP/SOX10 adenoviruses. Note that downregulation of *CD44* and *PDGFRB* messenger RNAs following SOX10 re-expression. (G) SOX10 reexpression in human schwannoma cells reduces PDGF-induced ERK1/2 MAP kinase activation. Human schwannoma cells infected with either control GFP or GFP/SOX10 adenoviruses were treated with PDGF-DD for 15 min. Blotting with HA antibody shows exogenously re-expressed SOX10 protein. (H) Re-introduction of SOX10 reduces both basal and PDGF-induced proliferation of human Merlin-null schwannoma cells at 72 h. (I–L) Immunolabelling of control GFP-infected (I and J) human schwannoma cells shows clear membrane-associated staining of CD44 (arrows in J), which is lost on SOX10 re-expression (K and L). Scale bar = 20  $\mu$ m. (M and N) Scanning electron microscopy of GFP- (M) and GFP/SOX10- (N) infected human schwannoma cells. Scale bar = 10  $\mu$ m. Note that control cells frequently overgrow one another, upon SOX10 re-expression the cells appear to respect cell–cell boundaries.

cycle exit in human Merlin-null schwannoma cells. The zinc finger transcription factor KROX20 is a key regulator of Schwann cell differentiation, regulating both cell cycle exit and the gene expression that drives myelination (Topilko *et al.*, 1994; Zorick *et al.*, 1999; Parkinson *et al.*, 2004, 2008). The normal control by KROX20 of cell cycle exit and myelin gene induction, using enforced KROX20 expression, is unchanged in Merlin-null schwannoma cells, but the ability of these cells to induce endogenous markers of differentiation, such as KROX20 and OCT6, in response to a myelination cue is markedly reduced, implying a Merlin-dependent loss of correct signalling upstream of KROX20. The mouse SC4 immortalized Merlin-null schwannoma cell line has been widely used in understanding the biology of merlin (Morrison *et al.*, 2007; Hennigan *et al.*, 2012). Interestingly, in SC4 cells, we did not observe a significant increase in either periaxin expression or downregulation of c-Jun when KROX20 was expressed (data not shown). We presume that the additional genetic events in the immortalization of these cells preclude their normal differentiation in response to KROX20 in contrast to primary human Schwann or schwannoma cells.

The transcriptional control of OCT6 and KROX20 have been well characterized in Schwann cells, and both rely on the high mobility group transcription factor SOX10 for their expression in developing Schwann cells (Ghislain and Charnay 2006; Finzsch *et al.*, 2010; Reiprich *et al.*, 2010; Bremer *et al.*, 2011). Correspondingly, we have found reduced levels of *SOX10* messenger RNA and protein in all the human schwannoma tumours we have examined, and in further confirmation of the role of

SOX10, re-expression of SOX10 in human schwannoma cells reverts many of the changes seen in these cells back to a normal phenotype, critically including restoring the capacity to express myelin proteins *in vitro* and suppressing basal and PDGF-induced proliferation.

To separate the effects of merlin and SOX10 loss in Schwann cells, we have used small interfering RNA and mice with conditional alleles of these genes, and, both *in vitro* and *in vivo*, we see that loss of merlin has no apparent direct effect on SOX10 expression in Schwann cells. Analysis of the role of SOX10 in Schwann cells *in vivo* has shown that it is required for normal Schwann cell–axon interaction, cell cycle withdrawal, expression of both OCT6 and KROX20 and the normal induction and maintenance of myelination in peripheral nerves. In addition, loss of SOX10 leads to an increase in other cell types within the nerve, such as endothelial cells and pericytes and an apparent decrease in S100-expressing Schwann cells (Finzsch *et al.*, 2010; Bremer *et al.*, 2011), although we, in agreement with other studies (Rosenbaum *et al.*, 1998; Kanaan *et al.*, 2008), find no decrease in S100 staining in the schwannoma tumours we have analysed; in accordance with this, we find no increase in S100 $\beta$  messenger RNA levels when SOX10 is re-expressed in schwannoma cells (data not shown).

Our examination of SOX10-null mouse Schwann cells *in vitro* shows that loss of SOX10 alone replicates phenotypic changes associated with human schwannoma cells, such as increased numbers of focal adhesions, increased proliferation and increased expression of and signalling through the PDGF receptor beta

(PDGFRB). From these data, we would propose a model by which loss of Merlin and SOX10 contribute to the phenotype of human schwannoma cells. For the example of the PDGFRB, loss of Merlin in cells only causes a modest (1.3-fold) increase in PDGFRB messenger RNA levels but is involved in recycling of the receptor protein at the cell surface (Lallemand *et al.*, 2009), whereas our data show that SOX10 appears to transcriptionally repress the PDGFRB gene; thus, the combined effects of both merlin and SOX10 loss will lead to the large increases seen in PDGFRB protein levels in human schwannoma cells. Similarly, for *CD44* messenger RNA, which encodes the hyaluronate receptor that transmits proliferative signalling in Merlin-null cells (Morrison *et al.*, 2001; Bai *et al.*, 2007), SOX10 transcriptionally represses *CD44* messenger RNA levels when reintroduced into human schwannoma cells.

The mechanism by which SOX10 expression is lost in human schwannomas is unclear; the simplest explanation would be that loss of merlin leads to changes in cell signalling that repress *SOX10* transcription, but our *in vitro* and *in vivo* analysis show this to not be the case. Analysis of both Merlin-null knockout cells and Merlin re-expression experiments in human schwannoma cells shows that the acute loss of Merlin has no effect on levels of SOX10. What regulates SOX10 in Schwann cells is presently unclear, but an upstream enhancer, known as U3, contributes to SOX10 expression and can be activated by a number of transcription factors such as SOX10, SOX9, AP2 $\alpha$  (TFAP2A) and FOXD3 *in vitro* (Wahlbuhl *et al.*, 2012). A role for activated Notch has also been shown in repressing SOX10 expression in Schwann cells (Li *et al.*, 2004), and Merlin loss in *Drosophila* leads to increased Notch signalling (Maitra *et al.*, 2006), but inhibition of Notch signalling by the  $\gamma$ -secretase inhibitor *N*-[*N*-(3, 5-difluorophenacetyl)-l-alanyl]-*S*-phenylglycine *t*-butyl ester; 25  $\mu$ M of the  $\gamma$ -secretase SOX10 expression in schwannoma cells (data not shown).

The key to the loss of SOX10 in human schwannomas, we propose, may be the chromosomal location of SOX10 and Merlin on chromosome 22. Remarkably, the two genes are close to one another on the long arm of chromosome 22 (merlin 22q12.2, SOX10 22q13.1). The second 'hit' and loss of merlin function in neurofibromatosis type 2 commonly involves loss of heterozygosity or a large deletion of chromosome 22, and our meta-analysis of cytogenetic studies on schwannoma tumours indicates that the region encoding SOX10 would be lost in >50% of these tumours (Bruder *et al.*, 1999; Antinheimo *et al.*, 2000; Mantripragada *et al.*, 2003; Koutsimpelas *et al.*, 2011), although this is probably an underestimate as the chromosomal marker resolution of several of these studies is not particularly good. SOX10 has been shown to bind and activate its own promoter (Wahlbuhl *et al.*, 2012); therefore, it is possible that the loss of one allele could lead to the transcriptional silencing of the remaining intact allele in Schwann cells and the corresponding fall in both SOX10 messenger RNA and protein observed in schwannoma tumours. Another alternative, which has been proposed, given the differences observed between patients with neurofibromatosis type 2 and the phenotype of mice with Merlin-null Schwann cells, is that there is a third 'hit' at another locus in addition to the loss of merlin (Woods *et al.*, 2003). Such an additional genetic change could directly or indirectly affect SOX10 expression and

contribute to the phenotype we observe in human schwannoma cells. The production and characterization of a conditional SOX10 allele, and the separation of merlin and SOX10 on different mouse chromosomes (chromosomes 11 and 15, respectively) would now allow the testing of the effects of SOX10 hemizyosity or complete loss on the biology of Merlin-null cells and may lead, perhaps, to the development of a mouse model that more closely matches the human disease.

In conclusion, we have identified consistent loss of SOX10 in human schwannoma cells, and our results identify that this loss strongly contributes to the phenotype of this tumour cell type. These results, for the first time, provide a direct link between the processes involved in normal Schwann cell differentiation and homeostasis in the peripheral nerve and the development and pathology of schwannoma tumours. Further characterization of the relative contributions of merlin and SOX10 to the biology of schwannoma cells may give important insights into how to treat these clinically important tumours.

## Acknowledgements

The authors wish to thank Dr Roy Moate, Peter Bond and Glenn Harper from the University of Plymouth electron microscopy centre for excellent technical assistance with electron microscopy experiments.

## Funding

This work was funded by a Wellcome Trust. Wellcome Trust project grant (awarded to D.B.P.), the Northcott Devon Medical Foundation and the South West Regional Development Agency for a PhD studentship (awarded to R.D.S.D).

## Supplementary material

Supplementary material is available at *Brain* online.

## References

- Ammoun S, Flaiz C, Ristic N, Schuldt J, Hanemann CO. Dissecting and targeting the growth factor-dependent and growth factor-independent extracellular signal-regulated kinase pathway in human schwannoma. *Cancer Res* 2008; 68: 5236–45.
- Ammoun S, Schmid MC, Ristic N, Zhou L, Hilton D, Ercolano E, et al. The role of insulin-like growth factors signaling in Merlin-deficient human schwannomas. *Glia* 2012; 60: 1721–33.
- Ammoun S, Schmid MC, Zhou L, Ristic N, Ercolano E, Hilton DA, et al. Insulin-like growth factor-binding protein-1 (IGFBP-1) regulates human schwannoma proliferation, adhesion and survival. *Oncogene* 2011; 31: 1710–22.
- Antinheimo J, Sallinen SL, Sallinen P, Haapasalo H, Helin H, Horelli-Kuitunen N, et al. Genetic aberrations in sporadic and neurofibromatosis 2 (NF2)-associated schwannomas studied by comparative genomic hybridization (CGH). *Acta Neurochir (Wien)* 2000; 142: 1099–104; discussion 1104–5.

- Arthur-Farraj P, Wanek K, Hantke J, Davis CM, Jayakar A, Parkinson DB, et al. Mouse schwann cells need both NRG1 and cyclic AMP to myelinate. *Glia* 2011; 59: 720–33.
- Bai Y, Liu YJ, Wang H, Xu Y, Stamenkovic I, Yu Q. Inhibition of the hyaluronan-CD44 interaction by merlin contributes to the tumor-suppressor activity of merlin. *Oncogene* 2007; 26: 836–50.
- Bremer M, Frob F, Kichko T, Reeh P, Tamm ER, Suter U, et al. SOX10 is required for Schwann-cell homeostasis and myelin maintenance in the adult peripheral nerve. *Glia* 2011; 59: 1022–32.
- Britsch S, Goerich DE, Riethmacher D, Peirano RI, Rossner M, Nave KA, et al. The transcription factor SOX10 is a key regulator of peripheral glial development. *Genes Dev* 2001; 15: 66–78.
- Brockes JP, Fields P, Raff MC. Studies on cultured rat Schwann cells. I. Establishment of purified populations from cultures of peripheral nerve. *Brain Res* 1979; 165: 105–18.
- Bruder CE, Ichimura K, Tingby O, Hirakawa K, Komatsuzaki A, Tamura A, et al. A group of schwannomas with interstitial deletions on 22q located outside the NF2 locus shows no detectable mutations in the NF2 gene. *Hum Genet* 1999; 104: 418–24.
- Cooper J, Li W, You L, Schiavon G, Pepe-Caprio A, Zhou L, et al. Merlin/NF2 functions upstream of the nuclear E3 ubiquitin ligase CRL4DCAF1 to Suppress Oncogenic Gene Expression. *Sci Signal* 2011; 4: pt6.
- Curto M, Cole BK, Lallemand D, Liu CH, McClatchey AI. Contact-dependent inhibition of EGFR signaling by Nf2/Merlin. *J Cell Biol* 2007; 177: 893–903.
- Feltri ML, D'Antonio M, Previtali S, Fasolini M, Messing A, Wrabetz L. PO-Cre transgenic mice for inactivation of adhesion molecules in Schwann cells. *Ann N Y Acad Sci* 1999; 883: 116–23.
- Finzsch M, Schreiner S, Kichko T, Reeh P, Tamm ER, Bosl MR, et al. SOX10 is required for Schwann cell identity and progression beyond the immature Schwann cell stage. *J Cell Biol* 2010; 189: 701–12.
- Flaicz C, Ammoun S, Biebl A, Hanemann CO. Altered adhesive structures and their relation to RhoGTPase activation in Merlin-deficient Schwannoma. *Brain Pathol* 2009; 19: 27–38.
- Flaicz C, Kaempchen K, Matthies C, Hanemann CO. Actin-Rich protrusions and nonlocalized GTPase activation in Merlin-deficient schwannomas. *J Neuropathol Exp Neurol* 2007; 66: 608–16. doi: 10.1097/nen.0b013e318093e555.
- Flaicz C, Utermark T, Parkinson DB, Poetsch A, Hanemann CO. Impaired intercellular adhesion and immature adherens junctions in Merlin-deficient human primary schwannoma cells. *Glia* 2008; 56: 506–15.
- Ghislain J, Charnay P. Control of myelination in Schwann cells: a KROX20 cis-regulatory element integrates OCT6, Brn2 and SOX10 activities. *EMBO Rep* 2006; 7: 52–8.
- Giovannini M, Robanus-Maandag E, van der Valk M, Niwa-Kawakita M, Abramowski V, Goutebroze L, et al. Conditional biallelic Nf2 mutation in the mouse promotes manifestations of human neurofibromatosis type 2. *Genes Dev* 2000; 14: 1617–30.
- Hanemann CO. Magic but treatable? Tumours due to loss of merlin. *Brain* 2008; 131(Pt 3): 606–15.
- Hanemann CO, Bartelt-Kirbach B, Diebold R, Kampchen K, Langmesser S, Utermark T. Differential gene expression between human schwannoma and control schwann cells. *Neuropathol Appl Neurobiol* 2006; 32: 605–14.
- Hanemann CO, Claudia R, Sandra K, Susanne W, Florian S, Hans Werner M. Improved culture methods to expand schwann cells with altered growth behaviour from CMT1A patients. *Glia* 1998; 23: 89–98.
- Harrisingh MC, Perez-Nadales E, Parkinson DB, Malcolm DS, Mudge AW, Lloyd AC. The Ras/Raf/ERK signalling pathway drives Schwann cell dedifferentiation. *EMBO J* 2004; 23: 3061–71.
- He TC, Zhou S, da Costa LT, Yu J, Kinzler KW, Vogelstein B. A simplified system for generating recombinant adenoviruses. *Proc Natl Acad Sci USA* 1998; 95: 2509–14.
- He Y, Kim JY, Dupree J, Tewari A, Melendez-Vasquez C, Svaren J, et al. Yy1 as a molecular link between neuregulin and transcriptional modulation of peripheral myelination. *Nat Neurosci* 2010; 13: 1472–80.
- Hennigan RF, Moon CA, Parysek LM, Monk KR, Morfini G, Berth S, et al. The NF2 tumor suppressor regulates microtubule-based vesicle trafficking via a novel Rac, MLK and p38(SAPK) pathway. *Oncogene* 2012; Apr 23. doi: 10.1038/onc.2012.135.
- Jaegle M, Mandemakers W, Broos L, Zwart R, Karis A, Visser P, et al. The POU factor OCT-6 and Schwann cell differentiation. *Science* 1996; 273: 507–10.
- Jagalur NB, Ghazvini M, Mandemakers W, Driegen S, Maas A, Jones EA, et al. Functional dissection of the oct6 schwann cell enhancer reveals an essential role for dimeric sox10 binding. *J Neurosci* 2011; 31: 8585–94.
- Jessen KR, Brennan A, Morgan L, Mirsky R, Kent AR, Hashimoto Y, et al. The Schwann cell precursor and its fate: a study of cell death and differentiation during gliogenesis in rat embryonic nerves. *Neuron* 1994; 12: 509–27.
- Jessen KR, Mirsky R. The origin and development of glial cells in peripheral nerves. *Nat Rev Neurosci* 2005; 6: 671–82.
- Kaempchen K, Mielke K, Utermark T, Langmesser S, Hanemann CO. Upregulation of the Rac1/JNK signaling pathway in primary human schwannoma cells. *Hum Mol Genet* 2003; 12: 1211–21.
- Kanaan HA, Gardner PA, Yeane G, Prevedello DM, Monaco EA 3rd, Murdoch G, et al. Expanded endoscopic endonasal resection of an olfactory schwannoma. *J Neurosurg Pediatr* 2008; 2: 261–5.
- Kao SC, Wu H, Xie J, Chang CP, Ranish JA, Graef IA, et al. Calcineurin/NFAT signaling is required for neuregulin-regulated Schwann cell differentiation. *Science* 2009; 323: 651–4.
- Kipanyula MJ, Woodhoo A, Rahman M, Payne D, Jessen KR, Mirsky R. Calcineurin-nuclear factor of activated t cells regulation of KROX-20 expression in Schwann cells requires elevation of intracellular cyclic AMP. *J Neurosci Res* 2012; 91: 105–15.
- Koutsimpelas D, Felmeden U, Mann WJ, Brieger J. Analysis of cytogenetic aberrations in sporadic vestibular schwannoma by comparative genomic hybridization. *J Neurooncol* 2011; 103: 437–43.
- Lallemand D, Curto M, Saotome I, Giovannini M, McClatchey AI. NF2 deficiency promotes tumorigenesis and metastasis by destabilizing adherens junctions. *Genes Dev* 2003; 17: 1090–100.
- Lallemand D, Manent J, Couvelard A, Watilliaux A, Siena M, Chareyre F, et al. Merlin regulates transmembrane receptor accumulation and signaling at the plasma membrane in primary mouse Schwann cells and in human schwannomas. *Oncogene* 2009; 28: 854–65.
- Le N, Nagarajan R, Wang JY, Araki T, Schmidt RE, Milbrandt J. Analysis of congenital hypomyelinating Egr2Lo/Lo nerves identifies SOX2 as an inhibitor of Schwann cell differentiation and myelination. *Proc Natl Acad Sci USA* 2005; 102: 2596–601.
- Li W, You L, Cooper J, Schiavon G, Pepe-Caprio A, Zhou L, et al. Merlin/NF2 suppresses tumorigenesis by inhibiting the E3 ubiquitin ligase CRL4(DCAF1) in the nucleus. *Cell* 2010; 140: 477–90.
- Li Y, Rao PK, Wen R, Song Y, Muir D, Wallace P, et al. Notch and Schwann cell transformation. *Oncogene* 2004; 23: 1146–52.
- Maitra S, Kulikauskas RM, Gavilan H, Fehon RG. The tumor suppressors Merlin and Expanded function cooperatively to modulate receptor endocytosis and signaling. *Curr Biol* 2006; 16: 702–9.
- Maka M, Stolt CC, Wegner M. Identification of SOX8 as a modifier gene in a mouse model of Hirschsprung disease reveals underlying molecular defect. *Dev Biol* 2005; 277: 155–69.
- Mantripragada KK, Buckley PG, Benetkiewicz M, De Bustos C, Hirvela C, Jarbo C, et al. High-resolution profiling of an 11 Mb segment of human chromosome 22 in sporadic schwannoma using array-CGH. *Int J Oncol* 2003; 22: 615–22.
- McClatchey AI, Fehon RG. Merlin and the ERM proteins—regulators of receptor distribution and signaling at the cell cortex. *Trends Cell Biol* 2009; 19: 198–206.
- McClatchey AI, Giovannini M. Membrane organization and tumorigenesis—the NF2 tumor suppressor, Merlin. *Genes Dev* 2005; 19: 2265–77.
- Monk KR, Naylor SG, Glenn TD, Mercurio S, Perlin JR, Dominguez C, et al. A G protein-coupled receptor is essential for Schwann cells to initiate myelination. *Science* 2009; 325: 1402–5.



- Monk KR, Oshima K, Jors S, Heller S, Talbot WS. Gpr126 is essential for peripheral nerve development and myelination in mammals. *Development* 2011; 138: 2673–80.
- Morgan L, Jessen KR, Mirsky R. The effects of cAMP on differentiation of cultured Schwann cells: progression from an early phenotype (O4+) to a myelin phenotype (P0+, GFAP-, N-CAM-, NGF-receptor-) depends on growth inhibition. *J Cell Biol* 1991; 112: 457–67.
- Morrison H, Sherman LS, Legg J, Banine F, Isacke C, Haipek CA, et al. The NF2 tumor suppressor gene product, merlin, mediates contact inhibition of growth through interactions with CD44. *Genes Dev* 2001; 15: 968–80.
- Morrison H, Sperka T, Manent J, Giovannini M, Ponta H, Herrlich P. Merlin/neurofibromatosis type 2 suppresses growth by inhibiting the activation of Ras and Rac. *Cancer Res* 2007; 67: 520–7.
- Mulvihill JJ, Parry DM, Sherman JL, Pikus A, Kaiser-Kupfer MI, Eldridge R. Neurofibromatosis 1 (Recklinghausen Disease) and Neurofibromatosis 2 (Bilateral Acoustic Neurofibromatosis) An Update. *Ann Intern Med* 1990; 113: 39–52.
- Nagarajan R, Svaren J, Le N, Araki T, Watson M, Milbrandt J. EGR2 mutations in inherited neuropathies dominant-negatively inhibit myelin gene expression. *Neuron* 2001; 30: 355–68.
- Nakai Y, Zheng Y, MacCollin M, Ratner N. Temporal control of Rac in Schwann cell-axon interaction is disrupted in NF2-mutant schwannoma cells. *J Neurosci* 2006; 26: 3390–5.
- National Institutes of Health Consensus Development Conference. Neurofibromatosis: conference statement. *Arch Neurol* 1988; 45: 575–8.
- Nickols JC, Valentine W, Kanwal S, Carter BD. Activation of the transcription factor NF-kappaB in Schwann cells is required for peripheral myelin formation. *Nat Neurosci* 2003; 6: 161–7.
- Ogata T, Iijima S, Hoshikawa S, Miura T, Yamamoto S, Oda H, et al. Opposing extracellular signal-regulated kinase and Akt pathways control Schwann cell myelination. *J Neurosci* 2004; 24: 6724–32.
- Parkinson DB, Bhaskaran A, Arthur-Farraj P, Noon LA, Woodhoo A, Lloyd AC, et al. c-Jun is a negative regulator of myelination. *J Cell Biol* 2008; 181: 625–37.
- Parkinson DB, Bhaskaran A, Droggiti A, Dickinson S, D'Antonio M, Mirsky R, et al. KROX-20 inhibits Jun-NH2-terminal kinase/c-Jun to control Schwann cell proliferation and death. *J Cell Biol* 2004; 164: 385–94.
- Parkinson DB, Dickinson S, Bhaskaran A, Kinsella MT, Brophy PJ, Sherman DL, et al. Regulation of the myelin gene periaxin provides evidence for KROX-20-independent myelin-related signalling in Schwann cells. *Mol Cell Neurosci* 2003; 23: 13–27.
- Parkinson DB, Dong Z, Bunting H, Whitfield J, Meier C, Marie H, et al. Transforming growth factor beta (TGFbeta) mediates schwann cell death *in vitro* and *in vivo*: examination of c-jun activation, interactions with survival signals, and the relationship of tgfbeta-mediated death to schwann cell differentiation. *J Neurosci* 2001; 21: 8572–85.
- Pereira JA, Lebrun-Julien F, Suter U. Molecular mechanisms regulating myelination in the peripheral nervous system. *Trends Neurosci* 2012; 35: 123–34.
- Reiprich S, Kriesch J, Schreiner S, Wegner M. Activation of KROX20 gene expression by SOX10 in myelinating Schwann cells. *J Neurochem* 2010; 112: 744–54.
- Rosenbaum C, Kluwe L, Mautner VF, Friedrich RE, Muller HW, Hanemann CO. Isolation and characterization of Schwann cells from neurofibromatosis type 2 patients. *Neurobiol Dis* 1998; 5: 55–64.
- Rouleau GA, Merel P, Lutchman M, Sanson M, Zucman J, Marineau C, et al. Alteration in a new gene encoding a putative membrane-organizing protein causes neuro-fibromatosis type 2. *Nature* 1993; 363: 515–21.
- Shaw RJ, Paez JG, Curto M, Yaktine A, Pruitt WM, Saotome I, et al. The Nf2 tumor suppressor, merlin, functions in Rac-dependent signaling. *Dev Cell* 2001; 1: 63–72.
- Stolt CC, Lommes P, Sock E, Chaboissier MC, Schedl A, Wegner M. The SOX9 transcription factor determines glial fate choice in the developing spinal cord. *Genes Dev* 2003; 17: 1677–89.
- Topilko P, Schneider-Maunoury S, Levi G, Baron-Van Evercooren A, Chennoufi AB, Seitanidou T, et al. KROX-20 controls myelination in the peripheral nervous system. *Nature* 1994; 371: 796–9.
- Trofatter JA, MacCollin MM, Rutter JL, Murrell JR, Duyao MP, Parry DM, et al. A novel moesin-, ezrin-, radixin-like gene is a candidate for the neurofibromatosis 2 tumor suppressor. *Cell* 1993; 75: 826.
- Utermark T, Kaempchen K, Hanemann CO. Pathological adhesion of primary human schwannoma cells is dependent on altered expression of integrins. *Brain Pathol* 2003; 13: 352–63.
- Wahlbuhl M, Reiprich S, Vogl MR, Bosl MR, Wegner M. Transcription factor SOX10 orchestrates activity of a neural crest-specific enhancer in the vicinity of its gene. *Nucleic Acids Res* 2012; 40: 88–101.
- Wojciak-Stothard B, Potempa S, Eichholtz T, Ridley AJ. Rho and Rac but not Cdc42 regulate endothelial cell permeability. *J Cell Sci* 2001; 114(Pt 7): 1343–55.
- Woods R, Friedman JM, Evans DG, Baser ME, Joe H. Exploring the “two-hit hypothesis” in NF2: tests of two-hit and three-hit models of vestibular schwannoma development. *Genet Epidemiol* 2003; 24: 265–72.
- Xiao GH, Gallagher R, Shetler J, Skele K, Altomare DA, Pestell RG, et al. The NF2 tumor suppressor gene product, merlin, inhibits cell proliferation and cell cycle progression by repressing cyclin D1 expression. *Mol Cell Biol* 2005; 25: 2384–94.
- Zorick TS, Syroid DE, Brown A, Gridley T, Lemke G. KROX-20 controls SCIP expression, cell cycle exit and susceptibility to apoptosis in developing myelinating Schwann cells. *Development* 1999; 126: 1397–406.