

2016-02-01

# Intracellular NAD<sup>+</sup> levels are associated with LPS-induced TNF- release in pro-inflammatory macrophages

Al-Shabany, AJ

<http://hdl.handle.net/10026.1/4406>

---

10.1042/bsr20150247

Bioscience Reports

Portland Press Ltd.

---

*All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.*



## OPEN ACCESS

# Intracellular NAD<sup>+</sup> levels are associated with LPS-induced TNF- $\alpha$ release in pro-inflammatory macrophages

Abbas Jawad Al-Shabany\*, Alan John Moody\*, Andrew David Foey† and Richard Andrew Billington\*<sup>1</sup>

\*School of Biological Sciences, University of Plymouth, Drake Circus, Plymouth, PL4 8AA, U.K.

†School of Biomedical and Healthcare Sciences, University of Plymouth, Drake Circus, Plymouth, PL4 8AA, U.K.

## Synopsis

Metabolism and immune responses have been shown to be closely linked and as our understanding increases, so do the intricacies of the level of linkage. NAD<sup>+</sup> has previously been shown to regulate tumour necrosis factor- $\alpha$  (TNF- $\alpha$ ) synthesis and TNF- $\alpha$  has been shown to regulate NAD<sup>+</sup> homeostasis providing a link between a pro-inflammatory response and redox status. In the present study, we have used THP-1 differentiation into pro- (M1-like) and anti- (M2-like) inflammatory macrophage subset models to investigate this link further. Pro- and anti-inflammatory macrophages showed different resting NAD<sup>+</sup> levels and expression levels of NAD<sup>+</sup> homeostasis enzymes. Challenge with bacterial lipopolysaccharide, a pro-inflammatory stimulus for macrophages, caused a large, biphasic and transient increase in NAD<sup>+</sup> levels in pro- but not anti-inflammatory macrophages that were correlated with TNF- $\alpha$  release and inhibition of certain NAD<sup>+</sup> synthesis pathways blocked TNF- $\alpha$  release. Lipopolysaccharide stimulation also caused changes in mRNA levels of some NAD<sup>+</sup> homeostasis enzymes in M1-like cells. Surprisingly, despite M2-like cells not releasing TNF- $\alpha$  or changing NAD<sup>+</sup> levels in response to lipopolysaccharide, they showed similar mRNA changes compared with M1-like cells. These data further strengthen the link between pro-inflammatory responses in macrophages and NAD<sup>+</sup>. The agonist-induced rise in NAD<sup>+</sup> shows striking parallels to well-known second messengers and raises the possibility that NAD<sup>+</sup> is acting in a similar manner in this model.

**Key words:** immune responses, lipopolysaccharide (LPS), macrophages, pyridine nucleotides, second messenger, TNF- $\alpha$ .

Cite this article as: Bioscience Reports (2016) 36, e00301, doi:10.1042/BSR20150247

## INTRODUCTION

Previous work has revealed a fascinating link between metabolism and the ability of an organism to mount an immune response [1]. This link is bidirectional with examples of changes in metabolism being required in order to mount the appropriate response and with immune mediators being able to modulate metabolism [2]. Indeed, many diseases that were previously considered to be pure metabolic disorders now being reconsidered also as inflammatory diseases and *vice versa* [2]. Immunometabolism has been particularly well studied in macrophages, myeloid derived phagocytes with different phenotypes that can be involved in numerous innate immune functions from bacterial killing to wound healing [3]. In particular, classically activated, pro-inflammatory (M1) macrophages are predominantly glycolytic whereas alternatively activated; anti-inflammatory (M2) macrophages rely on

oxidative metabolism [4]. Forcing M1s to become oxidative, or M2s to become glycolytic, leads to a switch in phenotype. A front-line response to bacterial infection needs to be short lived as the consequent production of large amounts of pro-inflammatory mediators such as tumour necrosis factor- $\alpha$  (TNF- $\alpha$ ) can lead to a pro-inflammatory cascade resulting in potentially fatal cytokine storm [5].

Glycolysis can operate under anaerobic conditions (such as those that may be found at sites of infection) for short periods of time but is unsustainable in the long term due to lactate and ROS production, reflecting the role of M1s. On the other hand, M2s are involved in much longer term anti-inflammatory processes in which cytokine storm is not possible. Thus M2s are much more suited to aerobic oxidative metabolism of glucose and fatty acids which, while slower than glycolysis, produces more ATP per gram of substrate and can be sustained practically indefinitely. Given that NAD<sup>+</sup> is an absolute requirement for

**Abbreviations:** DPI, diphenyliodonium; NAMPT, nicotinamide phosphoribosyl transferase; NMNAT, nicotinamide mononucleotide adenylyl transferase; qPCR, quantitative PCR; TLR4, Toll-like receptor 4; TNF- $\alpha$ , tumour necrosis factor- $\alpha$ .

<sup>1</sup> To whom correspondence should be addressed (email richard.billington@plymouth.ac.uk).

glycolysis, it is perhaps not surprising that  $\text{NAD}^+$  levels have also been linked to immune function in macrophages. Indeed, a key role has been shown for  $\text{NAD}^+$  in the positive control of transcription of the major M1 pro-inflammatory cytokine  $\text{TNF-}\alpha$  via  $\text{NAD}^+$ -dependent sirtuin deacetylase activity [6]. Decreasing  $\text{NAD}^+$  levels, SIRT6 knock out or inhibition, blocks transcription of  $\text{TNF-}\alpha$  suggesting a link between the redox and thus metabolic, state of cell and pro-inflammatory responses. During glycolysis,  $\text{NAD}^+$  is reduced to  $\text{NADH}$  whereas  $\text{NADH}$  can be oxidized back to  $\text{NAD}^+$  during anaerobic lactic fermentation or during mitochondrial oxidative phosphorylation [7], thus,  $\text{NAD}^+/\text{NADH}$  ratios provide readout of glycolytic activity that would appear to be important in the regulation of  $\text{TNF-}\alpha$  transcription. Conversely, it has been found that  $\text{TNF-}\alpha$  itself can modulate intracellular  $\text{NAD}^+$  content in macrophages, by regulating the expression of a number of  $\text{NAD}^+$  homeostasis enzymes, presumably as some sort of feedback regulation [8]. Furthermore, LPS, a classic pro-inflammatory stimulus that leads to  $\text{TNF-}\alpha$  release, has also been shown to mediate the expression of NAMPT (nicotinamide phosphoribosyl transferase, a key enzyme in the  $\text{NAD}^+$  recycling pathway) in activated monocytes and macrophages [8,9].

It is thus clear that  $\text{NAD}^+$  is involved in macrophage pro-inflammatory responses involving  $\text{TNF-}\alpha$ . Previous interest in the pharmacological modulation of  $\text{NAD}^+$  homeostasis has led to a number of drugs entering clinical trials as chemotherapeutic agents for cancer [10]. Given the link between  $\text{NAD}^+$  and  $\text{TNF-}\alpha$ , it is also possible that such agents could be used to modulate pro-inflammatory responses. Indeed, FK866, an inhibitor of the  $\text{NAD}^+$  recycling pathway, decreased  $\text{TNF-}\alpha$  secretion in THP-1 monocytes and human monocyte during LPS stimulation [10,11]. We have investigated whether  $\text{NAD}^+$  levels could also be involved in the LPS-stimulated secretion of  $\text{TNF-}\alpha$  in monocytic THP-1 cells differentiated to generate M1-like and M2-like macrophage models. The generated subsets differed in their relative levels of expression of  $\text{NAD}^+$ -homeostasis enzymes and in their resting levels of  $\text{NAD}^+$  suggesting that  $\text{NAD}^+$  homeostasis may be phenotype specific. LPS induced robust  $\text{TNF-}\alpha$  production that was accompanied by large changes in intracellular  $\text{NAD}^+$  levels in M1-like cells, whereas this was not the case in M2-like cells. LPS also induced changes in the expression of  $\text{NAD}^+$ -homeostasis enzymes that differed between subsets. These data extend the intimate relationship between  $\text{NAD}^+$  levels and homeostasis and  $\text{TNF-}\alpha$ . Furthermore, to our knowledge, this is the first report of agonist stimulated  $\text{NAD}^+$  rises and bringing  $\text{NAD}^+$  a step closer to being classified as a second messenger.

## MATERIALS AND METHODS

### Cell culture maintenance, macrophage differentiation and stimulation

The human monocytic leukaemia cell line THP-1 (ECACC) was used routinely between passage 8 and 24 and maintained in

complete growth RPMI 1640 medium (Lonza) supplemented with 10% (v/v) FBS (Sera lab), 2 mM L-glutamine, 100 unit/ml penicillin and 100  $\mu\text{g/ml}$  streptomycin (P/S) (Lonza). Cells were kept at  $37^\circ\text{C}$ , under 5%  $\text{CO}_2$  in a humidified incubator.

For THP-1 differentiation into macrophage-like subsets, the cells were plated (at density of  $1 \times 10^6$  cell/ml) in complete medium and treated with PMA (M1-like; 25 ng/ml; 3 days followed by 5 days resting in fresh medium) or  $1,25\text{-(OH)}_2\text{-Vitamin D}_3$  (Sigma-Aldrich; M2-like; 10 nM; 7 days) [12].

Cells were stimulated with 100 ng/ml LPS derived from *Escherichia coli* K12-LPS (Autogen-Bioclear Ltd.) for the indicated time intervals. Dead cells were aspirated and the live cells and supernatant were harvested by centrifugation at 200 g for 5 min, and stored at  $-20^\circ\text{C}$  until the day of analysis. For cell counting, adherent cells (M1-like) were de-attached by trypsinization using 0.5% (v/v) of trypsin with 0.2% (v/v) EDTA (Sigma-Aldrich).

### $\text{NAD}^+$ cycling assay for the measurement of intracellular $\text{NAD}^+$ levels

For the determination of  $\text{NAD}^+$  content, the washed cell pellets were lysed with 0.2 M HCl and heated to  $100^\circ\text{C}$  for 10 min. The intracellular  $\text{NAD}^+$  content were measured using a modified enzymatic cycling assay [13]. Briefly, the acid extract was mixed with 151  $\mu\text{l}$  of cycling buffer: 98 mM bicine pH 8.0, containing 0.2 M sodium hydroxide, 1.63 mM PES (phenazine ethosulfate), 0.412 mM MTT (3-(4,5-Dimethyl-2-thiazolyl)-2,5-diphenyl-2H-tetrazolium bromide), 9.8% (v/v) absolute ethanol, 3.92 mM EDTA and 5  $\mu\text{l}$  of yeast alcohol dehydrogenase (ADH) (400 unit/mg) were incubated for 30 min in the dark at room temperature. The absorbance was read at 565 nm using a plate reader (Versa Max, Molecular Devices).

### Quantitative analysis of cytokine levels

To quantify extracellular  $\text{TNF-}\alpha$  levels, cell culture supernatants were analysed using sandwich ELISA according to the manufacturer's instructions. Supernatants were harvested from THP-1-derived macrophage subsets at the indicated time points, and protocols were followed and compared with standard curves between the range of 7 and 5000 pg/ml using the international standards available from NIBSC, Potter's Bar, UK. Briefly, 96-well ELISA plates (Nunc, Fisher Scientific) were coated with (4  $\mu\text{g/ml}$ ) of mouse anti-human  $\text{TNF-}\alpha$  (BD Pharmingen), in PBS as a coating buffer, and incubated overnight at  $4^\circ\text{C}$ .

Thereafter, samples were added into the plate, and incubated overnight at  $4^\circ\text{C}$ . For  $\text{TNF-}\alpha$  detection, biotinylated mouse anti- $\text{TNF-}\alpha$  at 1  $\mu\text{g/ml}$  in 2% BSA (w/v in PBS) buffer were applied, and horseradish peroxidase streptavidin conjugate were added, followed by incubation with TMB reagent (Insight Biotechnology Ltd.). Colorimetric development was assayed spectrophotometrically by an OPTI Max tuneable microplate reader set to 450 nm and analysed by Soft max Pro version 2.4.1 software (Versa Max, Molecular Devices).

**Table 1** Primer sequences used for qPCR

Gene	Sequence (5' → 3')	Size (bp)	Gene	Sequence (5' → 3')	Size
GAPDH	For: CCCACTCCTCCACCTTTGAC	20	IDO	For: GCCTGCGGGAAGCTTATG	18
	Rev: CTGTTGCTGTAGCCAAATTCGT	22		Rev: TGGCTTGCAGGAATCAGGAT	20
CD38	For: GCACCACCAAGCGCTTTC	18	NMNAT	For: TCATTCAATCCCACCAACA	22
	Rev: TCCCATACACTTTGGCAGTCTACA	24		Rev: AGGAGAGATGATGCCTTTGACAA	23
CD157	For: GGGGAAGGCAGCATGAAAGTC	20	NAMPT	For: TCCGGCCCCGAGATGAAT	17
	Rev: GGTCCACGCACTGTAAGAGCTT	22		Rev: TGCTTGTGTTGGGTGGATATTG	22

### RNA extraction, reverse transcription and qPCR

Total RNA was extracted from approximately  $1 \times 10^6$  cultured cells, and isolated using Gen Elute Mammalian Total RNA kit (Sigma–Aldrich) with a DNase I Digestion treatment for 15 min according the manufacturer's instructions. The amount and purity of RNA was measured using a Nano-drop Plus spectrophotometer (Bioscience, Cambridge Bioscience Ltd). Total RNA (25 ng/ml) was subsequently reverse transcribed in 20  $\mu$ l total volume containing: 2.5  $\mu$ M random nonamers, 10 unit/ $\mu$ l RT and 500  $\mu$ M deoxynucleotide mixture, for 1 h at 42 °C before quantitative PCR (qPCR) was performed. The primers used for qPCR are listed in Table 1.

The qPCR was performed to quantify specific mRNA using SYBR green detection kit according to the manufacturer's instructions (Sigma–Aldrich). The cDNA product was utilized in the reaction mixture with 0.4  $\mu$ M of specific forward/reverse primer sets alongside with (0.025 unit/ $\mu$ l reaction) of SYBR-Green Jumpstart Tag ready mixture (Sigma–Aldrich). The reactions were performed under the following conditions: 94 °C for 2 min followed by 40 cycles of 94 °C for 30 s, 60 °C for 30 s and 72 °C for 30 s. The quality of the RNA samples was confirmed by 1.5 % (w/v) agarose gel electrophoresis. The relative amounts of mRNA, for each target transcript, were normalized according to the house-keeping gene (GAPDH) glyceraldehyde-3-phosphate dehydrogenase [14].

### Statistical analysis


Statistical calculations were performed using Statgraphics Centurion XVI (Stat Point Technologies Inc.). All data were tested for normality using the Anderson–Darling test ( $P < 0.05$  was used as the cut off for non-normal data). Levene's test was used to check for differences in variance between each group. For normally distributed data, one-way ANOVA followed by Tukey's test was performed. Data are reported as means  $\pm$  S.E.M. for triplicate samples from three independent experiments unless otherwise indicated. Differences were considered significant if  $P < 0.05$ .

## RESULTS

The link between NAD<sup>+</sup> and TNF- $\alpha$  is now well established in macrophages. Briefly, NAD<sup>+</sup> regulates production of TNF- $\alpha$

in a sirtuin-dependent manner [6] and TNF- $\alpha$  has been shown to modulate the expression of a number of enzymes that are involved in NAD<sup>+</sup> homeostasis [8]. In order to investigate this link further, intracellular NAD<sup>+</sup> level changes have been investigated during LPS stimulation of THP-1 monocytic cell line-derived macrophages. This system has been shown to generate cells that express markers and are functionally very close to pro-(M1) or anti-(M2) inflammatory macrophages [12,15]. Due to the heterogeneity of macrophage function, distinct M1-like pro-inflammatory and M2-like anti-inflammatory phenotypes were investigated to elucidate any potential differences. Upon LPS stimulation, M1-like cells showed a distinctive change in NAD<sup>+</sup> levels that consisted of an initial fast increase, peaking at 30 min followed by a lower but sustained high level of NAD<sup>+</sup> that lasted for several hours, starting to return to unstimulated levels at 24 h post-stimulation. Unstimulated cells showed a constant low level of NAD<sup>+</sup> throughout the experiment (Figure 1A). In contrast, LPS stimulation of M2-like cells failed to increase intracellular NAD<sup>+</sup> levels and both stimulated and unstimulated cells showed similar levels of NAD<sup>+</sup> throughout the experiment. Basal levels of NAD<sup>+</sup> in unstimulated M1-like and M2-like cells were different with M1-like cells appearing to have three-fold higher level of NAD<sup>+</sup> than M2-like cells (Figures 1A and 1B). During stimulation experiments, media was collected and secreted TNF- $\alpha$  assayed over the same time course. As expected, M1-like cells showed significant TNF- $\alpha$  secretion that peaked at 8 h and decayed back to basal levels by 24 h (Figure 1C). In contrast, M2-like cells secreted a relatively small amount of TNF- $\alpha$  which corresponded to less than 10% of that secreted by M1-like cells (Figure 1D). In M2-like cells, the TNF- $\alpha$  peak was short lived and had returned almost to basal levels by 8 h however there was a low level of TNF- $\alpha$  that persisted to beyond 18 h.

These results suggest that LPS affects NAD<sup>+</sup> homeostasis differently in the two phenotypes. While conducting experiments to assess the effect of the NAD<sup>+</sup>-recycling (NAMPT) inhibitor FK866 in the two phenotypes in the absence of LPS stimulation, we noticed opposing effects on NAD<sup>+</sup> levels. In M2-like cells, FK866 reduced the levels of NAD<sup>+</sup> over time as would be expected due to inhibition of NAD<sup>+</sup> synthesis and constitutive NAD<sup>+</sup> consumption (Figure 2B). In M1-like cells, on the other hand, FK866 gradually increased NAD<sup>+</sup> levels over time (Figure 2A). We are unsure as to how FK866 can cause an increase in NAD<sup>+</sup> levels but this does suggest that there is a fundamental difference in how the two phenotypes perform NAD<sup>+</sup> homeostasis. To check whether the two phenotypes show different




**Figure 1** Intracellular NAD<sup>+</sup> and TNF-α levels in THP-1-derived macrophages subsets

LPS effect on NAD<sup>+</sup> levels in pro-inflammatory (A; M1-like) and anti-inflammatory (B; M2-like) macrophages. TNF-α levels in M1-like (C) and M2-like (D) macrophages after LPS challenge. Data shown are mean  $\pm$  S.E.M. for three separate experiments ( $n = 3$ ).


expression of some key regulators of NAD<sup>+</sup> homeostasis, real-time qPCR (RT-qPCR) was performed on unstimulated M1-like and M2-like cells and compared with expression in the monocytic progenitor THP-1 cell line (Figure 3). Both phenotypes showed increased expression of IDO compared with THP-1. The NADase CD38, that mediates NAD<sup>+</sup> degradation and generates cyclic ADP ribose/NAADP [16], was significantly up-regulated in M1-like and down-regulated in M2-like cells whereas the opposite was true for CD157, a homologue of CD38 [17]. Finally, NMNAT (nicotinamide mononucleotide adenylyl transferase) that catalyses the final step in NAD<sup>+</sup> biosynthesis [18], was slightly, but not significantly up-regulated in both phenotypes whereas NAMPT was significantly up-regulated only in M1-like cells.

To further investigate this and the source of the LPS-induced rise in NAD<sup>+</sup> in M1-like cells, LPS stimulation of M1-like cells

was performed in the presence of diphenyliodonium (DPI), an NAD(P)H oxidase (flavoenzyme) inhibitor, FK866 a NAMPT inhibitor or 1-MT, an IDO inhibitor. Concurrently TNF-α secretion was measured. In M1s, DPI was able to significantly attenuate both the initial NAD<sup>+</sup> peak and the sustained rise caused by LPS suggesting that one source of the NAD<sup>+</sup> is the oxidation of NADH during the respiratory burst as might be expected (Figure 4A). Surprisingly, both FK866 and 1-MT were also able to attenuate both the initial peak and the sustained rise in NAD<sup>+</sup> perhaps suggesting that multiple pathways lead to LPS-induced NAD<sup>+</sup> level rises (Figures 4B and 4C). Most striking was the effect of the inhibitors on TNF-α production. DPI was able to almost completely abolish TNF-α secretion although a small amount was secreted over 8–24 h but significantly less than in the absence of DPI (Figure 5A). FK866 was able to partially



**Figure 2** Effect of FK866 (100 nM) on NAD<sup>+</sup> levels in M1-like (A) and M2-like (B) macrophages. Data shown are mean  $\pm$  S.E.M. for three separate experiments (n = 3).



**Figure 3** Gene expression profiles of NAD<sup>+</sup> homoeostasis enzymes in M1- and M2-like macrophages relative to THP-1 monocytes.


Data shown are mean  $\pm$  S.E.M. of fold change for three separate experiments (n = 3–5). The data were analysed by one-way ANOVA. \*P < 0.05.

attenuate TNF- $\alpha$  secretion whereas 1-MT had no effect (Figures 5B and 5C). These data suggest that the link between NAD<sup>+</sup> and TNF- $\alpha$  secretion is perhaps complex and may require both NAD<sup>+</sup> and the activity of NAD(P)H oxidase.

Finally, it was assessed whether LPS stimulation and the concurrent changes in NAD<sup>+</sup> levels and TNF- $\alpha$  secretion were able

to feedback and affect the expression of some key regulators of NAD<sup>+</sup> homoeostasis over 24 h. In M1-like cells, LPS stimulation caused a significant increase in IDO mRNA with smaller increases in CD38, CD157 and NAMPT expression whereas NMNAT expression was unaffected (Figure 6A). The increase was only significant for CD38. Despite LPS not being able to






**Figure 4** Pharmacological modulation of intracellular NAD<sup>+</sup> levels by DPI (A), FK866 (B) and 1-MT (C) in M1-like cells. Data shown are mean  $\pm$  S.E.M. for three separate experiments ( $n = 3-4$ ).

stimulate NAD<sup>+</sup> changes in M2-like cells, LPS stimulation caused a similar but significant increase in IDO and in NAMPT expression (Figure 6B). CD38, CD157 and NMNAT expression were largely unaffected although there was a modest but significant increase in NMNAT and CD38 expression.

## DISCUSSION

It is becoming clear that cellular metabolic status is intrinsically linked to immune system function, particularly in macrophages [4]. One of the best studied links is that between NAD<sup>+</sup>/NADH and TNF- $\alpha$  with NAD(H) levels known to regulate TNF- $\alpha$  synthesis [6] and TNF- $\alpha$  levels known to regulate expression of NAD(H) homeostasis enzymes [8]. In the present study, we have further extended the knowledge on this link by showing

that TNF- $\alpha$  release is also correlated to NAD<sup>+</sup> levels in a pro-inflammatory (M1-like) macrophage model. Our data show that LPS stimulation of M1-like cells induces a biphasic rise in NAD<sup>+</sup> levels. LPS is well known to activate NAD(P)H oxidase [19], in a Toll-like receptor 4 (TLR4)-dependent manner (oxidative burst) which catalyses the oxidation of NAD(P)H to NAD<sup>+</sup> [20]. This may explain, at least in part, the rapid peak that was observed in NAD<sup>+</sup> levels as this was partially inhibited by the NAD(P)H oxidase inhibitor DPI. However, both FK866 (an NAD<sup>+</sup> recycling inhibitor) and 1-MT (an IDO inhibitor) were also able to partially inhibit this initial rise. This may suggest that the NAD<sup>+</sup> is also derived from a number of different synthesis and recycling pathways. The sustained rise in NAD<sup>+</sup> levels also suggests that NAD<sup>+</sup> synthesis occurs as it would be expected that the cell would return the redox balance after the oxidative burst relatively rapidly so as to maintain the NAD<sup>+</sup>/NADH ratio. Thus, it would appear that LPS has a direct effect on NAD<sup>+</sup> levels in M1-like cells. LPS was unable to alter NAD<sup>+</sup> levels in M2-like




**Figure 5** Pharmacological modulation of TNF- $\alpha$  release by DPI (A), FK866 (B) and 1-MT (C) in M1-like cells  
Data shown are mean  $\pm$  S.E.M. for three separate experiments ( $n = 4$ ).

cells. NAD<sup>+</sup> levels also appear to be correlated with secretion of TNF- $\alpha$ . In M1-like cells, LPS induced a robust secretion of TNF- $\alpha$  which was almost completely blocked by DPI and partially inhibited by FK866 suggesting that NAD<sup>+</sup> levels are, in some way, related to TNF- $\alpha$  secretion. The exact mechanism is unclear but it has been shown that TNF- $\alpha$  secretion requires active glycolysis [21]. Glycolysis is inhibited by high NADH levels as NAD<sup>+</sup> is absolutely required as a coenzyme. Both oxidation of NADH to NAD<sup>+</sup> and a simple increase in NAD<sup>+</sup> synthesis could both provide these conditions. Indeed, lactic fermentation exists to oxidize NADH to NAD<sup>+</sup> under anaerobic conditions such that glycolysis may continue. M2-like cells responded to LPS by releasing a relatively small amount of TNF- $\alpha$ . Crucially, this was not associated with any changes in NAD<sup>+</sup> levels, suggesting that the signalling between TLR4 and TNF- $\alpha$  release is via a different pathway compared with M1-like cells. TLR4 is known to signal via different intracellular pathways, one of which

is NAD(P)H oxidase dependent which causes the oxidative burst and ROS production [22] and converts NADH to NAD<sup>+</sup> [20].

One interesting outcome from our data is that the two different phenotypes appear to have quite different mechanisms and levels of machinery for NAD<sup>+</sup> homeostasis, raising the exciting possibility that NAD<sup>+</sup> homeostasis may be phenotype-specific and even an underlying cause of phenotypic differences. Indeed, previous work has shown that NAD<sup>+</sup> and TNF- $\alpha$  are inextricably linked suggesting that high NAD<sup>+</sup> may be 'pro-inflammatory' [6,8,23]. We found that resting NAD<sup>+</sup> levels were higher in M1-like than M2-like cells and this may be linked with the finding that M1s are predominantly glycolytic [24] and thus, require relatively more NAD<sup>+</sup>. M1-like cells, in resting state, displayed higher expression of the recycling pathway enzymes NAMPT than both M2-like cells and the parent THP-1 cells as well as higher levels of the NADase CD38. The CD38 homologue and NADase CD157, however, displayed higher expression in





**Figure 6** Gene expression profile of NAD<sup>+</sup> homoeostasis enzymes in M1-like (A) and M2-like (B) macrophages after LPS challenge

Gene expression is expressed as fold change relative to the control. Data shown are mean  $\pm$  S.E.M. for three separate experiments ( $n = 4$ ). The data were analysed by one-way ANOVA. \* $P < 0.05$ .

M2-like cells, the converse of CD38. NAMPT is the rate limiting enzyme in the NAD<sup>+</sup> recycling pathway [25], this pathway is the most active of the three known pathways that lead to NAD<sup>+</sup> synthesis; the others being the *de novo* (IDO) pathway [26] and the nicotinamide riboside kinase (NRK) pathway [27]. The greatest increase in expression, compared with parent THP-1 cells, was in IDO in both phenotypes and it might be expected that the *de novo* pathway would, thus, be the biggest contributor to NAD<sup>+</sup> levels. However, this pathway is known to contribute little to the intracellular NAD<sup>+</sup> pools with as little as 1 in 60 molecules of NAD<sup>+</sup> being derived from tryptophan [28]. As NAD<sup>+</sup> levels represent a steady state between synthesis and degradation, it is perhaps surprising that the NADase, CD38, was also up-regulated in M1-like cells. This may reflect greater turnover of NAD<sup>+</sup> in these cells or that the increase in CD38 is related to its receptor functions. CD38 is unusual in that it can act as both a cell surface receptor (binding to adhesion molecule-1, CD31) and as an enzyme [29] but our knowledge of how exactly the two functions interact is scant [30]. CD38 is known to be a marker of classically activated (M1-like) macrophages [31] and our results suggest that CD38 might contribute to differentiated macrophage responses. For the M2-like cells, aside from modest increases in IDO and CD157 expression, NAD<sup>+</sup> homoeostasis enzymes were largely unchanged compared with the parent THP-1 cells.

Finally the gene expression of the same enzymes was investigated after LPS challenge. In M1-like cells, mRNA expression of all enzymes except NMNAT increased over 20 h after LPS addition with the largest effects on IDO and CD38. Iqbal and Zaidi [8] have shown that TNF- $\alpha$  application to macrophages increased the expression of the same enzymes although they also showed an increase in NMNAT expression. Furthermore, Fedele et al. [29] also showed that LPS increases CD38 expression in monocyte-derived dendritic cells in an NF- $\kappa$ B-

dependent manner. It is likely that the TNF- $\alpha$  released during the experiment is partially responsible for the expression observed and thus why the results largely follow the same trend. However, the data presented here were generated after LPS stimulation and early signalling events may have initiated expression pathways that were then augmented or modulated by autocrine/paracrine signals upon TNF- $\alpha$  secretion. M2-like cells, that secreted a lesser amount of TNF- $\alpha$ , showed subtle differences in expression profiles of the enzymes. Whereas IDO and CD38 both increased, CD157 remained unchanged. Most striking was the differential regulation of NMNAT and NAMPT mRNA expression. NAMPT expression increased whereas NMNAT expression remained unchanged. Large increases in NAMPT in M1 would suggest that the cells are priming themselves for NAD<sup>+</sup> production as this is the rate limiting enzyme of the recycling pathway. Although we have been unable to measure long-term effects after LPS stimulation due to limitations in the cell model, it is tempting to postulate that changes in NAD<sup>+</sup> homoeostasis caused by these changes in gene expression, might contribute to changes in cell phenotype post-infection.

These data serve to reinforce the link between NAD<sup>+</sup>, TNF- $\alpha$  and inflammatory responses. Recent focus has shifted to the links between metabolism and innate immune responses and NAD<sup>+</sup> is a key player both as readout of the metabolic status of a cell and as a signalling molecule. Of particular interest is that the two phenotypes appear to have different mechanisms for handling NAD<sup>+</sup> and this may reflect different needs in terms of NAD<sup>+</sup>. Furthermore, their basal and stimulated expression profiles of a number of NAD<sup>+</sup> homoeostasis enzymes also differ suggesting that they can further model how NAD<sup>+</sup> homoeostasis is carried out depending on the circumstances of the cell. Although NAD<sup>+</sup> is central to the links between metabolism and immunology, there has been precious little work on NAD<sup>+</sup> homoeostasis in

this context and this may prove to be an interesting avenue to pursue in different aspects of immune responses.

Finally, that LPS induces rises in NAD<sup>+</sup> levels, suggests that NAD<sup>+</sup> may be acting in a similar manner to a classical second messenger. Indeed, of the five criteria suggested by Sutherland [32], we have shown NAD<sup>+</sup> fulfilling four in this work namely: agonist stimulated changes in NAD<sup>+</sup> levels, NAD<sup>+</sup> synthesis and degradation, antagonism of NAD<sup>+</sup> action blocking the cellular response and intracellular NAD<sup>+</sup> targets. This cell/response system is not ideal for intracellular application of NAD<sup>+</sup> as cell damage may release TNF- $\alpha$  making any observations hard to interpret but should intracellular NAD<sup>+</sup> application be shown to mimic agonist responses in another system, NAD<sup>+</sup> could be regarded as a true second messenger, albeit a slightly unusual one.

#### AUTHOR CONTRIBUTION

Abbas Al-Shabany performed experiments, performed data analysis and wrote the manuscript, Alan John Moody performed data analysis, Andrew Foey and Richard Billington conceived experiments, analysed data and wrote the manuscript.

#### ACKNOWLEDGEMENTS

We thank the College of Science, Al-Qadisiyah University, Ministry of Higher Education and Scientific Research (MOHESR), Iraq for a PhD studentship to Abbas Jawad Al-Shabany.

## REFERENCES

- Hotamisligil, G.S. and Erbay, E. (2008) Nutrient sensing and inflammation in metabolic diseases. *Nat. Rev. Immunol.* **8**, 923–934 [PubMed](#)
- Schertzer, J.D. and Steinberg, G.R. (2014) Immunometabolism: the interface of immune and metabolic responses in disease. *Immunol. Cell Biol.* **92**, 303 [PubMed](#)
- Murray, R.J. and Wynn, T.A. (2011) Protective and pathogenic functions of macrophage subsets. *Nat. Rev. Immunol.* **11**, 723–737 [PubMed](#)
- Galvan-Pena, S. and O'Neill, L.A. (2014) Metabolic reprogramming in macrophage polarization. *Front. Immunol.* **5**, 420 [PubMed](#)
- Finlay, B.B. and McFadden, G. (2006) Anti-immunology: evasion of the host immune system by bacterial and viral pathogens. *Cell* **124**, 767–782 [PubMed](#)
- Van Gool, F., Galli, M., Gueydan, C., Kruys, V., Prevot, P.P., Bedalov, A., Mostoslavsky, R., Alt, F.W., De Smedt, T. and Leo, O. (2009) Intracellular NAD<sup>+</sup> levels regulate tumor necrosis factor protein synthesis in a sirutin-dependent manner. *Nat. Med.* **15**, 206–210 [PubMed](#)
- Maciver, N.J., Jacobs, S.R., Wieman, H.L., Wofford, J.A., Coloff, J.L. and Rathmell, J.C. (2008) Glucose metabolism in lymphocytes is a regulated process with significant effects on immune cell function and survival. *J. Leukoc. Biol.* **84**, 949–957 [PubMed](#)
- Iqbal, J. and Zaidi, M. (2006) TNF- $\alpha$  regulates cellular NAD<sup>+</sup> metabolism in primary macrophages. *Biochem. Biophys. Res. Commun.* **342**, 1312–1318 [PubMed](#)
- Rongvaux, A., Shea, R.J., Mulks, M.H., Gigot, D., Urbain, J., Leo, O. and Andris, F. (2002) Pre-B-cell colony-enhancing factor, whose expression is up-regulated in activated lymphocytes, is a nicotinamide phosphoribosyltransferase, a cytosolic enzyme involved in NAD biosynthesis. *Eur. J. Immunol.* **32**, 3225–3234 [PubMed](#)
- Holen, K., Saltz, L.B., Hollywood, E., Burk, K. and Hanauske, A.R. (2008) The pharmacokinetics, toxicities, and biologic effects of FK866, a nicotinamide adenine dinucleotide biosynthesis inhibitor. *Invest. New Drugs* **26**, 45–51 [PubMed](#)
- Schilling, E., Wehrhahn, J., Klein, C., Raulien, N., Ceglarek, U. and Hauschildt, S. (2012) Inhibition of nicotinamide phosphoribosyltransferase modifies LPS-induced inflammatory responses of human monocytes. *Innate Immun.* **18**, 518–530 [PubMed](#)
- Daigneault, M., Preston, J.A., Marriott, H.M., Whyte, M.K. and Dockrell, D.H. (2010) The identification of markers of macrophage differentiation in PMA-stimulated THP-1 cells and monocyte-derived macrophages. *PLoS One* **5**, e8668 [PubMed](#)
- Leonardo, M.R., Dailly, Y. and Clark, D.P. (1996) Role of NAD<sup>+</sup> in regulating the adhE gene of *Escherichia coli*. *J. Bacteriol.* **178**, 6013–6018 [PubMed](#)
- Livak, K.J. and Schmittgen, T.D. (2001) Analysis of relative gene expression data using real-time quantitative PCR and the 2<sup>−</sup> $\Delta\Delta C(T)$  Method. *Methods* **25**, 402–408 [PubMed](#)
- Schwende, H., Fitzke, E., Ambis, P. and Dieter, P. (1996) Differences in the state of differentiation of THP-1 cells induced by phorbol ester and 1,25-dihydroxyvitamin D<sub>3</sub>. *J. Leukoc. Biol.* **59**, 555–561 [PubMed](#)
- Howard, M., Grimaldi, J.C., Bazan, J.F., Lund, F.E., Santos-Argumedo, L., Parkhouse, R.M., Walseth, T.F. and Lee, H.C. (1993) Formation and hydrolysis of cyclic ADP-ribose catalyzed by lymphocyte antigen CD38. *Science* **262**, 1056–1059 [PubMed](#)
- Ferrero, E. and Malavasi, F. (1997) Human CD38, a leukocyte receptor and ectoenzyme, is a member of a novel eukaryotic gene family of nicotinamide adenine dinucleotide<sup>+</sup>-converting enzymes: extensive structural homology with the genes for murine bone marrow stromal cell antigen 1 and aplysian ADP-ribosyl cyclase. *J. Immunol.* **159**, 3858–3865 [PubMed](#)
- Magni, G., Amici, A., Emanuelli, M., Raffaelli, N. and Ruggieri, S. (1999) Enzymology of NAD<sup>+</sup> synthesis. *Adv. Enzymol. Relat. Areas Mol. Biol.* **73**, 135–182 [PubMed](#)
- Remer, K.A., Brice, M. and Jungi, T.W. (2003) Toll-like receptor-4 is involved in eliciting an LPS-induced oxidative burst in neutrophils. *Immunol. Lett.* **85**, 75–80 [PubMed](#)
- Riebel, B.R., Gibbs, P.R., Wellborn, W.B. and Bommarium, A.S. (2002) Cofactor regeneration of NAD<sup>+</sup> from NADH: novel water-forming NADH oxidases. *Adv. Synth. Catal.* **344**, 1156–1168
- Dietl, K., Renner, K., Dettmer, K., Timischl, B., Eberhart, K., Dorn, C., Hellerbrand, C., Kastenberger, M., Kunz-Schughart, L.A., Oefner, P.J. et al. (2010) Lactic acid and acidification inhibit TNF- $\alpha$  secretion and glycolysis of human monocytes. *J. Immunol.* **184**, 1200–1209 [PubMed](#)
- Park, H.S., Jung, H.Y., Park, E.Y., Kim, J., Lee, W.J. and Bae, Y.S. (2004) Cutting edge: direct interaction of TLR4 with NAD(P)H oxidase 4 isozyme is essential for lipopolysaccharide-induced production of reactive oxygen species and activation of NF- $\kappa$ B. *J. Immunol.* **173**, 3589–3593 [PubMed](#)
- Bruzzone, S., Moreschi, I., Guida, L., Usai, C., Zocchi, E. and De Flora, A. (2006) Extracellular NAD<sup>+</sup> regulates intracellular calcium levels and induces activation of human granulocytes. *Biochem. J.* **393**, 697–704 [PubMed](#)
- Tavakoli, S., Zamora, D., Ullevig, S. and Asmis, R. (2013) Bioenergetic profiles diverge during macrophage polarization: implications for the interpretation of 18F-FDG PET imaging of atherosclerosis. *J. Nucl. Med.* **54**, 1661–1667 [PubMed](#)



- 25 Revollo, J.R., Grimm, A.A. and Imai, S. (2004) The NAD biosynthesis pathway mediated by nicotinamide phosphoribosyltransferase regulates Sir2 activity in mammalian cells. *J. Biol. Chem.* **279**, 50754–50763 [PubMed](#)
- 26 Magni, G., Amici, A., Emanuelli, M., Orsomando, G., Raffaelli, N. and Ruggieri, S. (2004) Enzymology of NAD<sup>+</sup> homeostasis in man. *Cell Mol. Life Sci.* **61**, 19–34 [PubMed](#)
- 27 Bieganski, P. and Brenner, C. (2004) Discoveries of nicotinamide riboside as a nutrient and conserved NRK genes establish a Preiss–Handler independent route to NAD<sup>+</sup> in fungi and humans. *Cell* **117**, 495–502 [PubMed](#)
- 28 Horwitt, M.K., Harvey, C.C., Rothwell, W.S., Cutler, J.L. and Haffron, D. (1956) Tryptophan-niacin relationships in man: studies with diets deficient in riboflavin and niacin, together with observations on the excretion of nitrogen and niacin metabolites. *J. Nutr.* **60**, 1–43 [PubMed](#)
- 29 Fedele, G., Frasca, L., Palazzo, R., Ferrero, E., Malavasi, F. and Ausiello, C.M. (2004) CD38 is expressed on human mature monocyte-derived dendritic cells and is functionally involved in CD83 expression and IL-12 induction. *Eur. J. Immunol.* **34**, 1342–1350 [PubMed](#)
- 30 Al-Abady, Z.N., Durante, B., Moody, A.J. and Billington, R.A. (2013) Large changes in NAD<sup>+</sup> levels associated with CD38 expression during HL-60 cell differentiation. *Biochem. Biophys. Res. Commun.* **442**, 51–55 [PubMed](#)
- 31 Shubinsky, G. and Schlesinger, M. (1997) The CD38 lymphocyte differentiation marker: new insight into its ectoenzymatic activity and its role as a signal transducer. *Immunity* **7**, 315–324 [CrossRef](#) [PubMed](#)
- 32 Sutherland, E.W., Robins, G.A. and Butcher, R.W. (1968) Some aspects of the biological role of adenosine 3',5'-monophosphate (cyclic AMP). *Circulation* **37**, 279–306 [CrossRef](#)

---

**Received 25 September 2015/9 December 2015; accepted 4 January 2016**

---

**Accepted Manuscript online 13 January 2016, doi 10.1042/BSR20150247**

---