

# Journal of International Medical Research

<http://imr.sagepub.com/>

---

## Physical therapy in multiple sclerosis differs across Europe: Information regarding an ongoing study

Kamila Rasová, Patricia Martinkova, Davide Cattaneo, Johanna Jonsdottir, Thomas Henze, Ilse Baert, Paul Van Asch, Carme Santoyo, Tori Smedal, Stachowiak Malgorzata, Jenny Freeman, Anders Romberg and Peter Feys

*Journal of International Medical Research* published online 12 August 2014

DOI: 10.1177/0300060514540249

The online version of this article can be found at:

<http://imr.sagepub.com/content/early/2014/08/12/0300060514540249>

---

Published by:



<http://www.sagepublications.com>

Additional services and information for *Journal of International Medical Research* can be found at:

**P<P**

Published online 12 August 2014 in advance of the print journal.

**Email Alerts:** <http://imr.sagepub.com/cgi/alerts>

**Subscriptions:** <http://imr.sagepub.com/subscriptions>

**Reprints:** <http://www.sagepub.com/journalsReprints.nav>

**Permissions:** <http://www.sagepub.com/journalsPermissions.nav>

>> [OnlineFirst Version of Record](#) - Aug 12, 2014

[What is This?](#)

## Physical therapy in multiple sclerosis differs across Europe: Information regarding an ongoing study

This paper provides information about an ongoing multi-staged online web questionnaire survey established by a core group within the Special Interest Group in Mobility, Rehabilitation in Multiple Sclerosis ([www.eurims.org](http://www.eurims.org)). The aims of the survey were to document the following across Europe: the availability and setting in which physical therapy for multiple sclerosis (MS) is delivered; the level of awareness and knowledge of different physical therapy approaches; the actual content of physical therapy and applied evaluation tools.

In 2011, the MS Barometer<sup>1</sup> identified the aspects of MS that were well managed in various European countries, and in which areas there was a need to improve policies and practices. The MS Barometer confirmed substantial differences between European countries in terms of access to care and treatment, as well as in the health-related quality-of-life of people with MS. Eliminating these disparities is a key aim of the MS Code of Good Practice,<sup>2</sup> which considers equal access to treatment (including rehabilitation) to be the right of every European citizen/patient.

As opposed to the MS Code of Good Practice, other studies have found relatively low rates of access to physical therapy for people with MS that vary widely across Europe. In Croatia, 41.3% of patients had accessed outpatient physical therapy in the previous 2 years.<sup>3</sup> A similar frequency of

access (38%) was shown in the UK in one study,<sup>4</sup> where 28.5% of patients participated in inpatient rehabilitation. In Croatia, only 17.4% had used inpatient rehabilitation.<sup>3</sup> In Sweden in 2008, 23% of patients accessed inpatient and 34% accessed outpatient physical therapy.<sup>5</sup> In the Netherlands, 30–36% of patients visited a physical therapist at least once per week.<sup>6</sup> On the other hand, a comparative study showed relatively high use of physical therapy services in some countries (Belgium 100%, Italy 87%, Greece 56% and UK 62%) compared with Estonia (15%), where service provision was sparse.<sup>7</sup>

The questionnaire survey was established in order to address the lack of information regarding the approach to physical therapy in European countries (e.g. how physical therapy is offered, length and intensity of typical inpatient and outpatient programmes, how [and who] plans the overall physical therapy process, the type of evidence-based physical therapy implemented in routine clinical practice). A total of 28 European countries participated in the survey. The resulting data will help to identify needs and barriers related to indicators in physical therapy and rehabilitation. In addition, the data will help to develop a roadmap towards an improved, harmonized, evidence-based system of physical therapy in Europe, including the development of guidelines and education packages for health care professionals.



## Declaration of conflicting interest

The authors declare that there are no conflicts of interest.

## Funding

The survey is supported by a grant from RIMS, an educational grant from Novartis Pharma AG, 260045/SVV/2014 and PRVOUK P34.

## References

1. MS Barometer. The European multiple sclerosis platform, <http://www.emsp.org/projects/ms-id/160-ms-barometer-2011%5D> (2011, accessed 28 April 2014).
2. European Multiple Sclerosis Platform. Recommendation on rehabilitation services for persons with multiple sclerosis in Europe, <http://www.emsp.org/projects/eu-code-of-good-practice/184-recommendations-on-rehabilitation-services-for-persons-with-multiple-sclerosis-in-europe%5D> (2012, accessed 28 April 2014).
3. Milivojević I, Adamec I and Habek M. Utilization of physical rehabilitation among people with multiple sclerosis. *Ir J Med Sci* 2013; 182: 429–432.
4. Somerset M, Campbell R, Sharp DJ, et al. What do people with MS want and expect from health-care services? *Health Expect* 2001; 4: 29–37.
5. Gottberg K, Einarsson U, Ytterberg C, et al. Use of health care services and satisfaction with care in people with multiple sclerosis in Stockholm County: a population-based study. *Mult Scler* 2008; 14: 962–971.
6. Moorer P, Suurmeijer TH and Zwanikken CP. Health care utilization by people with multiple sclerosis in The Netherlands: results of two separate studies. *Disabil Rehabil* 2000; 22: 695–701.
7. Kersten P, McLellan DL, Gross-Paju K, et al. A questionnaire assessment of unmet needs for rehabilitation services and resources for people with multiple sclerosis: results of a pilot survey in five European countries. Needs Task group of MARCH (Multiple Sclerosis and Rehabilitation, Care and Health Services Research in Europe). *Clin Rehabil* 2000; 14: 42–49.

Kamila Řasová

Corresponding author:

Kamila Řasová, Department of Rehabilitation, Third Faculty of Medicine, Charles University in Prague, Ruska 87, Prague, 10000, Czech Republic.  
Email: kamila.rasova@gmail.com

- Patricia Martinkova<sup>1</sup>, Davide Cattaneo<sup>2</sup>, Johanna Jonsdottir<sup>3</sup>, Thomas Henze<sup>4</sup>, Ilse Baert<sup>5</sup>, Paul Van Asch<sup>6</sup>, Carme Santoyo<sup>7</sup>, Tori Smedal<sup>8</sup>, Stachowiak Małgorzata<sup>9</sup>, Jenny Freeman<sup>10</sup>, Anders Romberg<sup>11</sup> and Peter Feys<sup>12</sup>
- <sup>1</sup>Institute of Computer Science, Academy of Sciences of the Czech Republic, Pod Vodarenskou vezi 2, Prague 8, 182 07, Czech Republic
- <sup>2</sup>Don Gnocchi Foundation, Larice Lab, Santa Maria Nascente, Via Capecelatro 66, 20148 Milan, Italy
- <sup>3</sup>Neurorehabilitation at the Don Gnocchi Foundation, Larice Lab, Santa Maria Nascente, Via Capecelatro 66, 20148 Milan, Italy
- <sup>4</sup>PASSAUER WOLF Reha-Zentrum Nittenau, Rehabilitationsklinik für Neurologie-Geriatrie-Urologie, Eichendorffstr. 21 D-93149 Nittenau, Germany
- <sup>5</sup>Hasselt University, Campus Diepenbeek, REVAL Research Institute (BIOMED), Agoralaan building A, B-3590 Diepenbeek, Belgium
- <sup>6</sup>Fit Up, Fitness- and Physiotherapy Center, Mechelsesteenweg 192a, 2550 Kontich, Belgium
- <sup>7</sup>CEMCat. Neurorehabilitation Unit. Passeig de la Vall d'Hebron, 119, 08035 Barcelona, Spain
- <sup>8</sup>Norwegian Multiple Sclerosis Competence Centre, Department of Neurology/ Department of Physiotherapy, Haukeland University Hospital, Jonas Liesvei 65, 5021 Bergen, Norway

<sup>9</sup>*MS Rehabilitation Centre, Szpitalna 5,  
78-449 Borne Sulinowo, Poland*

<sup>10</sup>*Reader in Physiotherapy, Plymouth  
University, Devon PL6 8BH, UK*

<sup>11</sup>*Masku Neurological Rehabilitation Centre,  
Physiotherapy, Vaihemäentie 10, PL 15,  
21251 Masku, Finland*

<sup>12</sup>*Hasselt University, Campus Diepenbeek,  
REVAL Research Institute (BIOMED),  
Agoralaan building A,  
B-3590 Diepenbeek, Belgium*