

2015

# Foraminifera of the Fal Estuary (Cornwall), including taxa associated with maerl beds

Hart, Malcolm

<http://hdl.handle.net/10026.1/8430>

---

Geoscience in South-West England

---

*All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.*

# FORAMINIFERA OF THE FAL ESTUARY (CORNWALL), INCLUDING TAXA ASSOCIATED WITH MAERL BEDS

M.B. HART<sup>1,2</sup>, K. CARTWRIGHT<sup>3</sup>, B. FISK<sup>3</sup>, C.W. SMART<sup>1,2</sup>, C. CONSOLARO<sup>1,2</sup> AND J.M. HALL-SPENCER<sup>2,4</sup>



Hart, M.B., Cartwright, K., Fisk, B., Smart, C.W., Consolaro, C. and Hall-Spencer, J.M. 2015. Foraminifera of the Fal Estuary (Cornwall), including taxa associated with Maerl Beds. *Geoscience in South-West England*, **13**, 483-490.

The Fal Estuary (Cornwall) contains a nationally important accumulation of calcareous red seaweeds commonly referred to as maerl. Maerl beds are often associated with high benthic diversity but there has been little research done on their associated microfaunas. This investigation has studied the foraminifera that are found within samples of maerl and the adjacent sediments. Our samples were preserved and then stained with rose Bengal, in order to ascertain the 'living' (stained) assemblage of foraminifera. Only <1% of the taxa associated with the maerl appeared to be living at the time of collection in October 2012, and the assemblage of foraminifera was a mixture of taxa that are characteristic of open marine environments and those characteristic of estuarine and sea grass communities. The presence of pelagic ostracods and centric diatoms supports the notion that at least some of the high foraminiferal diversity reported from the maerl assemblages is the result of transported material trapped within the intricate maerl habitat. Foraminifera from other areas of the Fal Estuary are typical of saltmarsh, estuarine and near-shore marine assemblages reported elsewhere in South-West England.

<sup>1</sup> School of Geography, Earth & Environmental Sciences, Plymouth University, Drake Circus, Plymouth, PL4 8AA, U.K.

<sup>2</sup> Marine Institute, Plymouth University, Drake Circus, Plymouth, PL4 8AA, U.K.


<sup>3</sup> Exeter College, Hele Road, Exeter, EX4 4JS, Devon, U.K.

<sup>4</sup> School of Marine Science & Engineering, Plymouth University, Drake Circus, Plymouth, PL4 8AA, U.K.  
(E-mail: M.Hart@plymouth.ac.uk)


**Keywords:** Fal Estuary, foraminifera, maerl, charophytes, marine conservation.

## INTRODUCTION

The designation of Marine Conservation Zones (MCZ) in England is being undertaken by Defra through Natural England (NE) and the Joint Nature Conservation Committee (JNCC). The first two tranches of MCZs were approved in 2013 and 2016, although a further 50+ potential locations have been identified through a nationwide process (Sadri *et al.*, 2011; Lieberknecht *et al.*, 2011; Bureck *et al.*, 2013) to augment an existing network of marine conservation areas around England. Under the Habitats Directive of the European Union (Natura 2000), a number of areas have been listed as Special Areas of Conservation (SAC). One such area is the 6,387.8 ha of the Fal and Helford SAC which encompasses the Helford River, part of Falmouth Bay and the Fal Estuary (Figure 1). This area includes sea inlets, tidal rivers, estuaries, mud flats, sand flats, salt marsh, salt pasture, sand dunes, beaches, machair, cliffs and islets (Figure 2). This SAC is one of the most important ria systems in South-West England, with a central, sinuous, relatively deep (20–30 m) channel (Sheehan *et al.*, 2015). The low fresh water input from a number of small rivers (e.g., Pencuil, Fal, Truro, Carnon, etc.) has allowed the development of a range of fully marine habitats from the extremely sheltered to the wave-exposed, tide-swept open coastline. Of particular importance are the maerl beds that are found on St Mawes Bank and extensive areas of maerl 'gravel' (Sheehan *et al.*, 2015) which extend within an area of the Carrick Roads, Falmouth Bank and Falmouth Bay (Figures 1, 2). These are the largest known maerl beds in South-West England and they provide habitat for an extremely high diversity of algae and a great many infaunal and epifaunal species (Bosence and Wilson, 2003; Peña *et al.*, 2014). The Fal Estuary contains both living and dead maerl deposits,



**Figure 1.** The boundaries of the Fal and Helford Special Area of Conservation.



**Figure 2.** Distribution of sediments and facies in the Fal Estuary, Carrick Roads, St Mawes Bank and tidal estuaries.

with a salinity of >30‰ or just below (Bryan and Hummerston, 1973). The maerl was dredged for agricultural use until 2005, when the Board of Falmouth Harbour Commissioners ceased to licence its extraction, following UK Government advice from English Nature and the Department for the Environment, Fisheries and Rural Affairs (Defra): see Hall-Spencer (2005).

## MAERL

Maerl is the collective name for a number of species of red seaweeds (Rhodophyta) that develop hard, calcareous skeletons (Corallinaceae). Maerl has twig-like, branching forms and, as the living alga, requires sunlight to grow. They are found down to depths of 51 m in the NE Atlantic Ocean but are usually restricted to depths of <20 m (Peña *et al.*, 2014). Some Corallinaceae encrust rocks and shells but attached forms can often form rhodolith or maerl habitats in near-shore environments (Martin and Hall-Spencer, 2016). In tropical regions (e.g. Brazil) rhodoliths (Figure 3a) can be 10 cm in diameter and have a characteristic internal structure (Pascelli *et al.*, 2013). Rhodoliths are also well-known in the geological record, including the Tata Limestone Formation (mid-Cretaceous) of Hungary (Figure 3b). The maerl requires sufficient water movement (tidal flow or wave action) to remove fine sediment that may restrict growth. Dead maerl accumulates on the sea floor, often with a thin layer of purple/pink-coloured living maerl on the top (Figure 3c). In English waters there are approximately 15 km<sup>2</sup> of maerl habitat, which represents just 0.03% of the near-shore marine environment. The maerl beds of the Fal are, therefore, a significant proportion of the total. The maerl in the Fal Estuary is composed of two species: *Phymatolithon calcareum* (Pallas) and *Lithothamnion corallioides* (P. & H. Crouan) and has been described by Bosence (1976), Farnham and Bishop (1985), Irvine and Chamberlain (1994), Birkett *et al.* (1998) and Hall-Spencer *et al.* (2010). Beds of maerl are concentrated along the western coastline of Europe in the NE Atlantic Ocean (Peña *et al.*, 2014, fig. 1; Dutertre *et al.*, 2015), including South-West England, Brittany, Western Ireland, the Inner Hebrides and Iceland.


Maerl beds are particularly important near-shore habitats as they support a high diversity of benthic, marine, invertebrates

(Bosence, 1979). Areas of maerl are a 'Priority Habitat' within the UK Biodiversity Action Plan (see <http://jncc.defra.gov.uk/page-6023> and links to UK BAP priority species and habitats). As a result of devolution, the country-level biodiversity strategies in the UK Post-2010 Biodiversity Framework include 'Habitats of Principal Importance in England' (Natural Environment and Rural Communities (NERC) Act, 2006) which includes maerl beds as being of national importance. Dead maerl accumulates very slowly over thousands of years and can also contain small patches of living maerl such as near the channel into Falmouth Docks. Austin and Cage (2010) showed, in an investigation of two samples, that a maerl bed offshore the Isle of Bute (Scotland) contained a high diversity of foraminifera. A high diversity of foraminifera in maerl beds has also been reported in Recent deposits by Boillot (1964), Blanc-Vernet (1969), Rosset-Moulinier (1972) and Freiwald and Henrich (1994) as well as in Holocene sediments in the south of France (Morhange *et al.*, 2003).

Here, maerl-rich samples from the Fal Estuary have been investigated for their associated foraminifera and, unlike samples studied by Austin and Cage (2010), those investigated in this pilot study were preserved in buffered formalin before being stained with the protoplasmic stain, rose Bengal, in an attempt to distinguish the 'living' (stained) assemblage and the 'dead' (un-stained) component of the assemblage, some of which may have been transported onto the maerl after death.

## MATERIAL AND METHODS

Aside from the samples of maerl from the Fal Estuary (see Sheehan *et al.*, 2015), a number of other samples were collected from the headwaters of the Tresillian River near Pencalenick [SW 860 454], Fal River near Lamorran [SW 877 417], Percuil River near Trethem Mill [SW 862 363], Calenick Creek east of Calenick [SW 827 432], Tallack's Creek east of Devoran [SW 802 389], and in a boatyard near St Just-in-Roseland [SW 847 357]. The methods employed were comparable to those used by Stubbles (1993, 1999), Olugbode *et al.* (2005) and Hart *et al.* (2014). All the samples collected in the Fal Estuary system were stored directly in buffered formalin (10%), having been taken from the uppermost 1 cm of the inter-tidal sediments enclosed in a 10 cm diameter ring (Figure 4a). All samples were



**Figure 3.** Examples of rhodoliths, including maerl, from the Fal Estuary. (a) A large modern rhodolith from the coastline of Sergipe (Brazil), scale bar = 1 cm. (b) Fossil rhodolith from the mid-Cretaceous of Hungary, scale bar = 1 cm. (c) Maerl from the Fal Estuary with the specimen on the left (un-stained) which was dead at the time of collection while the specimen on the right (stained) may have been living at the time of collection, scale bar = 1 cm.





**Figure 4.** Collection of field samples at Trethem Mill at the tidal limit of the Percuil River. (a) Sample collection within the 10 cm diameter plastic ring. (b) Looking along the Percuil River, downstream towards St Mawes.

subsequently washed on a 63  $\mu\text{m}$  large diameter sieve to remove most of the fine sediment. The residue was then placed in rose Bengal organic stain (1 g per litre de-ionized water) for three hours before being re-washed on a 63  $\mu\text{m}$  large diameter sieve. Once the water ran clear of mud and stain, the samples were filtered and dried in a cool oven ( $<40^{\circ}\text{C}$ ). Dried residues were studied in a range of size fractions ( $>500\ \mu\text{m}$ ,  $500\text{--}250\ \mu\text{m}$ ,  $250\text{--}150\ \mu\text{m}$  and  $150\text{--}63\ \mu\text{m}$ ). Images of the foraminifera and other microfauna/microflora were generated using a JEOL 7100FE Field Emission scanning electron microscope.

The two bulk maerl samples collected from the Fal Estuary in October 2012 were preserved in buffered formalin and then washed on a 2 mm sieve (to remove the maerl and other shell fragments). The finer material was then washed on a large diameter 63  $\mu\text{m}$  sieve, after which they were soaked in the protoplasmic stain, rose Bengal, for three hours before being washed again on the 63  $\mu\text{m}$  sieve, filtered and then dried at  $<40^{\circ}\text{C}$  in a cool oven. Some of this maerl was stained bright pink and presumed to have been 'living' at the time of collection. The other components of the assemblage were studied in the range of size fractions described above.

The maerl was inspected at low magnification in order to determine if any epifauna was present. Serpulid worms, some of which were quite small, were frequently encountered but no adherent foraminifera were found. Even after quite vigorous washing, however, some foraminifera were still found in the interstices of the maerl, but these were not adherent forms.


## FORAMINIFERA

Foraminifera were recorded in every sample, often in large numbers. Almost all the taxa are well known in the estuarine and coastal waters of Cornwall and South Devon (Millett, 1885; Heron-Allen and Earland, 1930; Murray, 1965, 1970, 1971; Castignetti, 1997; Stubbles, 1993, 1999; Olughode *et al.*, 2005; Hart *et al.*, 2014). The distribution of foraminifera is directly comparable to that described in the Fowey Estuary, which is located only 30 km to the east of the Fal Estuary. The Fal Estuary is, however, a much larger system with many more rivers entering from a much larger catchment, which was extensively mined in the 19th Century (Pirrie *et al.*, 2003). In 1991, the Carnon River, Restronguet Creek and Carrick Roads suffered a major flood of acidic mine water from the former Wheal Jane Mine (Stubbles *et al.*, 1996) which caused a significant loss of biota (especially in Restronguet Creek). This is the primary reason for the exclusion of Restronguet Creek from the SAC, as concentrations of heavy metals are still present within the estuarine sediments (Pirrie *et al.*, 2003) despite a natural cleaning and water filtration process being installed at Wheal Jane. Samples collected in 2004/2005 from Restronguet Creek still recorded significantly elevated numbers of deformed foraminifera, as reported by Olughode *et al.* (2005). In this investigation deformed foraminifera have been recorded although the numbers, away from Restronguet Creek, are comparable to those reported in the Fowey Estuary (Hart *et al.*, 2014). Examples of deformed taxa are shown in Figure 5 (12, 15), and these species have been selected for illustration as they are typical of those recorded in the 2015 sampling campaign.

Some of the foraminifera collected from the Fal Estuary are illustrated in Figure 5, although many of the taxa have already been illustrated in Hart *et al.* (2014, figs 4, 5). The species encountered are listed in the Appendix and their numbers presented in Table 1 which gives the total counts of each species, together with the percentage of the total assemblage. In this pilot study the live/dead separation is not presented as this work is still on-going. In the study of the Fowey Estuary, Hart *et al.* (2014, fig. 6) identified five assemblages, each characterized by the dominance of one species or the co-dominance of two species. These assemblages can also be identified for the sample locations around the Fal Estuary. As all the collected samples were located in the upper estuary areas of the tributaries feeding Carrick Roads and the Fal Estuary, it was expected that the majority of samples would yield Assemblages 2 and 3, with Calenick Creek providing the closest assemblage to the saltmarsh seen at Shirehall Moor near Lostwithiel (Assemblage 1). *Miliammina fusca* is the dominant species found in Calenick Creek and at Trethem Mill, although *Haynesina germanica* and *Elphidium williamsoni* are the dominant species in the Tresillian River, Fal River (Lamorrán) and Tallacks' Creek (Restronguet Creek). Since the recovery from the Wheal Jane pollution incident in 1991 (Stubbles, 1993, 1999; Olughode *et al.*, 2005), Restronguet Creek has been dominated by *E. williamsoni*, *H. germanica* and *Ammonia* sp. cf. *A. aberdoveyensis* and the Tallack's Creek sample confirmed that this is still the case. *E. williamsoni* and *H. germanica* also dominated the boatyard sample at St Just-in-Roseland, although this location also recorded more normal marine species (*Quinqueloculina* spp., *Cibicides lobatulus* and *Reophax moniliformis*). The St Just-in-Roseland sample is, therefore, comparable to Assemblage 4 of the Fowey Estuary and, especially, the samples from the boatyard area near Golant.

The assemblage from the maerl samples was, as expected, quite different in being dominated by *Elphidium crispum*, *Astigerinata mamilla*, *Quinqueloculina* spp. and *Rosalina globularis*. Despite the sample being stained with rose Bengal following normal procedures, few individuals took up the stain, even though some of the maerl appears to have been living when collected (bright pink in colour). This may be due to the time of sampling as, by October in most of the estuaries in South-West England (e.g. Plymouth Sound; see Castignetti, 1997), the





**Figure 5.** Foraminifera and a charophyte from the Fal Estuary. (1, 2) *Ammonia batava* (Hofker, 1951). (3, 4) *Astigerinata mamilla* (Williamson, 1858). (5-8) *Cibicides lobatulus* (Walker & Jacob, 1798). (9, 10) *Elphidium crispum* (Linné, 1758). (11) *Elphidium williamsoni* Haynes, 1973. (12) *Elphidium earlandi* Cushman, 1936. (13, 14, 15) *Haynesina germanica* (Ehrenberg, 1840), with (15) showing deformed growth pattern in final whorl. (16) *Nitella opaca* (C. Agardh ex. Bruzelius) C. Agardh (1824) from the Percuil River at Tretham Mill. This charophyte oospore is badly preserved, indicating that it has been transported prior to sedimentation in the upper estuarine muds of the sample location (see Figure 4a). Scale bars all 100  $\mu$ m.

	Calenick Creek	Tresillian River (U)	Tresillian River (L)	Trethem Mill	Fal River Lamorran	Tallack's Creek	St Just Boatyard	Maerl
<i>J.macrescens</i>	32(11%)	1(0.5%)	7(2.0%)	9(2.5%)	2(0.5%)	-	-	1(0.25%)
<i>T.inflata</i>	-	-	-	1(0.3%)	-	-	1(0.5%)	-
<i>M.fusca</i>	247(84%)	7(2.5%)	3(1.0%)	168(49%)	34(9.5%)	-	2(1%)	-
<i>H.germanica</i>	11(4%)	267(92%)	279(86%)	102(29.8%)	194(53%)	191(73.75%)	106(46%)	7(1.5%)
<i>E.williamsoni</i>	3(1%)	14(5%)	14(4.0%)	40(11.8%)	72(20%)	63(24.25%)	73(31.5%)	13(3%)
<i>Quinque. spp.</i>	-	-	14(4.0%)	15(4.3%)	25(7%)	-	20(8.5%)	84(20%)
<i>A.mamilla</i>	-	-	1(0.5%)	-	-	-	-	109(26%)
<i>C.lobulatus</i>	-	-	2(1.0%)	1(0.3%)	-	-	1(0.5%)	22(5%)
<i>E.crispum</i>	-	-	1(0.5%)	-	-	-	-	128(30.5%)
<i>R.globularis</i>	-	-	1(0.5%)	-	-	-	-	18(4.25%)
<i>O.squamosa</i>	-	-	1(0.5%)	-	-	-	-	-
<i>A.aberdovey.</i>	-	-	-	7(2.0%)	31(8.5%)	5(2%)	12(5%)	-
<i>R.moniliformis</i>	-	-	-	-	5(1.5%)	-	16(7%)	-
<i>A.batava</i>	-	-	-	-	-	-	-	6(1.5%)
<b>Other marine</b>	-	-	-	-	-	-	-	8%
<b>Assemblage</b>	2	2/3	2/3	2/3	2/3	2/3	4	(mixed)

**Table 1.** Foraminifera assemblage counts of samples from the Fal Estuary and adjacent creeks. Assemblage numbers in the lowest row relate to those used by Hart et al. (2014). Abbreviations: the two samples from the Tresillian River came from upper (U) mud-flats and lower (L) mud-flats, *Quinque. spp.* = *Quinqueloculina spp.* and *A. aberdovey.* = *A. aberdoveyensis*. The assemblage numbers given in the bottom row refer to those established for the Fowey Estuary (Hart et al., 2014).

living assemblage of foraminifera is normally quite reduced.

The species present in the maerl samples included a mixture of saltmarsh (*Jadammina macrescens*), estuarine (*H. germanica*, *E. williamsoni*) and open marine species (*Ammonia batava*, *Trochammina ochracea*, *Cribratomoides jeffreysii*, *Asigerinata mamilla*, *Fissurina* sp., *Parafissurina* spp., *Oolina* spp. *Quinqueloculina clarensis*, *Q. bicornis*, *Brizalina* sp., *Globocassidulina* sp., and *Sejunctella earlandi*). There were also rare centric diatoms indicating an open marine source as well as a few elliptical forms more typical of estuarine assemblages. Some species of foraminifera had specimens of the coccolith *Emiliania huxleyi* sticking to their chambers: again indicating an open marine source although coccolithophores are known from some estuaries in South-West England (e.g. Plymouth Sound). The most common foraminiferan in the counted material was *E. crispum* (30.5%), closely followed by *A. mamilla* (26%). In a comparable count of a maerl sample from Stravanan Bay, Isle of Bute (Scotland), Austin and Cage (2010) give figures for *A. mamilla* (16.5%), *E. crispum* (1.5%) and *A. batavus* (1%) that look different to the samples collected in the Fal Estuary. Stravanan Bay is not estuarine and although there are sea grass meadows in the Firth of Clyde, none are thought to be close to Stravanan Bay. Sadri et al. (2011) recorded high numbers of living *E. crispum* on the fronds of sea grass (*Zostera marina*) in Tor Bay during the summer months. These data from Tor Bay suggest that, by October, when the maerl samples from the Fal Estuary were collected, specimens of *E. crispum* that had been living in association with the nearby sea grass meadows during the summer had been transported into the area of the maerl and trapped with a mixture of other marine and estuarine taxa. Any maerl sample provides only a snapshot of the accumulated sediment but it is clear that the recorded assemblage appears to be an admixture of transported marine, estuarine and sea grass communities. Sampling in May-July would provide a more definitive view of a maerl-associated living assemblage which may include adherent forms, such as *Patellina corrugata*, as recorded on maerl in the Firth of Clyde (Hall-Spencer, 1994).

## CHAROPHYTES

In a sample from the Percuil River near Trethem Mill (Figure 4a) a number of charophyte oospores have been recorded (Figure 5 (16)). Charophytes are freshwater 'green algae', colloquially known as stoneworts. This name is derived from the observation that the stems of these algae, and the oospores,

are often encrusted with calcium carbonate. This calcification has given the stoneworts a long geological record extending back to the Devonian Rhynie Chert environment. The Characea and Nitellacea have a 200 million year fossil record, with modern *Chara* represented by 16 species and *Nitella* represented by eight species. Stoneworts require a clear, low-nutrient, still, freshwater environment and their presence can be an excellent indicator of water quality (Stewart, 2004).

It is, therefore, significant that the sample collected from the Percuil River contained oospores identified as *Nitella opaca* (C. Agardh ex. Bruzelius) C. Agardh (1824). This is the type species of the genus *Nitella*, and the species identification has been confirmed by N.F. Stewart (pers. comm., 2015). Further details on this species are available in Bryant and Stewart (2011) and Guiry (2015). This is the first record of this species in the Roseland area (Stewart and Church, 1992), although its occurrence in the estuarine muds of the Percuil River raises the question of where these algae are living. The normal habitat would be lakes or brooks with clean, fresh water (Stewart, 1996), though there are records from slightly brackish waters.

Specimens of *Nitella opaca* are normally 5-50 cm high with an axis (stem) ~1 mm in diameter. The fresh oospores are usually dark (chestnut to black) in colour and the normal occurrence is in lakes and brooks. Stewart (2004) records six important locations in South-West England: Braunton Burrows, Newton Abbot clay-pits and heaths, Slapton Ley National Nature Reserve, Tinhay Quarry (South Hams), Widdicombe Ley (South Hams) in Devon and heaths and quarries on the Lizard Peninsula in Cornwall. The alga was not observed in the sampling area and so is presumed to be living upstream in the Percuil River. It would be interesting to determine the location, in case any conservation measures are required to protect the habitat.

## SUMMARY

The distribution of foraminifera in the upper reaches of the Fal Estuary follow predicted patterns of distribution and reflect the local water depth, salinity and temperature. The communities from St Just-in-Roseland appear more 'marine' than those reported in the Fowey Estuary, probably reflecting the much wider opening of the estuary to the open sea and the resulting higher salinities. Our maerl samples contained assemblages of foraminifera that were different in character from that recorded by Austin and Cage (2010) in a comparable



environment off the Isle of Bute in Scotland. The dominance of *Elphidium crispum* appears to reflect a derivation from fleshy seaweeds or the fronds of sea grasses, as none of these individuals were alive at the time of sampling. The presence of rare marine species (e.g. *Sejunctella earlandi*) and some estuarine taxa (e.g. *Haynesina germanica* and *Elphidium williamsoni*) indicates that the species recorded from the maerl represents an admixture of environments and that it includes transported foraminifera, trapped within the irregular surface of the algae. The presence of *Nitella opaca* in the Percuil River is interesting as this is the first record of this species in the Roseland area of Cornwall.

## ACKNOWLEDGEMENTS

All the samples from the Fal were collected as part of an investigation of the maerl beds being undertaken by the Marine Institute (Plymouth University) and the authors thank those involved in this work for providing these samples. The authors wish to thank Dr N.F. Stewart for his identification of *Nitella opaca*, and his guidance on possible further work relating to these algae. The authors also wish to thank Tim Absalom, who assisted with the production of some of the figures, while Dr Roy Moate (Electron Microscope Unit, Plymouth University) assisted with the SEM work. Kai Cartwright and Ben Fisk both acknowledge receipt of a Nuffield Research Placement from the Nuffield Foundation. The authors acknowledge the helpful comments on an earlier version of this paper provided by Dr Jonathan Larwood (Natural England).

## REFERENCES

- AGARDH, C.A. 1824. *Systema Algarum*. Lundae (Lund), Literis Berlingianis (Berling), i-xxxvii, 1-312.
- AUSTIN, W.E.N. and CAGE, A.G. 2010. High benthic foraminiferal species counts in a Clyde Sea maerl bed, western Scotland. In: HOWE, J.A., AUSTIN, W.E.N., FORWICK, M. and PAETZEL, M. (Eds), *Fjord Systems and Archives*. Geological Society, London, Special Publications, 344, 83-88.
- BIRKETT, D.A., MAGGS, C.A. and DRING, M.J. 1998. *Maerl (volume V). An overview of dynamic and sensitivity characteristics for conservation management of marine SACs*. Scottish Association for Marine Science (UK Marine SACs Project).
- BLANC-VERNET, L. 1969. Contribution à l'étude des foraminifères de Méditerranée. *Recueil des Travaux de la Station Marine d'Endoume*, **64-48**, 1-279.
- BOILLLOT, G. 1964. Géologie de la Manche occidentale. *Annals de l'Institut Océanographique*, **42**, 1-219.
- BOSENCE, D.W.J. 1976. *Ecological and sedimentological studies on some carbonate sediment producing organisms*. Unpublished PhD thesis, University of Reading.
- BOSENCE, D.W.J. 1979. Live and dead faunas from coralline algal gravels, Co. Galway. *Palaeontology*, **22**, 449-478.
- BOSENCE, D. and WILSON, J. 2003. Maerl growth, carbonate production rates and accumulation rates in the northeast Atlantic. *Aquatic Conservation, Marine and Freshwater Ecosystems*, **13**, S21-S31.
- BRYAN, G.W. and HUMMERSTONE, L.G. 1973. Brown seaweed as an indicator of heavy metals in estuaries in south-west England. *Journal of the Marine Biological Association of the United Kingdom*, **53**, 705-720.
- BRYANT, J.A. and STEWART, N.F. 2011. Phylum Chlorophyta. Order Charales. In: JOHN, D.M., WHITTON, B.A. and BROOK, A.J. (Eds), *The freshwater algal flora of the British Isles*, 2nd Edition. Cambridge University Press, Cambridge, 742-765.
- BURECK, C.V., ELLIS, N.V., EVANS, D.H., HART, M.B. and LARWOOD, J.G. 2013. Marine Geoconservation in the United Kingdom. *Proceedings of the Geologists' Association, London*, **124**, 581-592.
- CASTIGNETTI, P. 1997. *Population dynamics and facies association of recent foraminifera from a nearshore marginal marine environment: Plymouth Sound*. Unpublished PhD thesis, Plymouth University.
- DUTERTRE, M., GRALL, J., EHRHOLD, A. and HAMON, D. 2015. Environmental factors affecting maerl bed structure in Brittany. *European Journal of Phycology*. DOI 10.1080/09670262.2015.

- FARNHAM, W.F. and BISHOP, G.M. 1985. *Survey of the Fal Estuary, Cornwall*. 18th Symposium of the Underwater Association for Scientific Research, 53-63.
- FREIWALD, A. and HENRICH, R. 1994. Reefal coralline algal build-ups within the Arctic Circle: morphology and sedimentary dynamics under extreme environmental seasonality. *Sedimentology*, **41**, 963-984.
- GUIRY, M.D. 2015. In: GUIRY, M.D. and GUIRY, G.M. *Algaebase*. World-wide electronic publication, National University of Ireland, Galway. <http://www.algaebase.org>, last accessed on 15th October 2015.
- HALL-SPENCER, J.M. 1994. *Biological studies of nongeniculate Corallinaceae*. Unpublished PhD thesis, University of London.
- HALL-SPENCER, J. 2005. Ban on maerl extraction, in News. *Marine Pollution Bulletin*, **50**, 121.
- HALL-SPENCER, J.M., KELLY, J. and MAGGS, C.A. 2010. Background document for maerl beds. London: OSPAR Commission: 2010, **491/2010**.
- HART, M.B., STUBBLES, S.J., SMART, C.W., FISHER, J.K., HODDINOTT, C., MARSHALL-PENN, I. and YEO, A. 2014. Foraminifera from the Fowey Estuary, Cornwall. *Geoscience in South-West England*, **13**, 304-315.
- HERON-ALLEN, E. and EARLAND, A. 1930. The Foraminifera of the Plymouth District. *Journal of the Royal Microscopical Society, London*, **50**, 46-84, 161-199.
- IRVINE, L.M. and CHAMBERLAIN, Y.M. 1994. *Seaweeds of the British Isles: Volume 1, Rhodophyta, Part 2B, Corallinales, Hildenbrandiales*. British Phycological Society and The Natural History Museum, London, HMSO.
- LIEBERKNECHT, L.M., HOOPER, T.H., MULLIER, T.M., MURPHY, A., NEILLY, M., CARR, H., HAINES, R., LEWIN, S. and HUGHES, E. 2011. *Finding Sanctuary final report and recommendations*. A report submitted by the Finding Sanctuary stakeholder project to Defra, the Joint Nature Conservation Committee, and Natural England. Available at [www.finding-sanctuary.org](http://www.finding-sanctuary.org) and The UK National Archives [http://na.europearchive.org/\\*http://finding-sanctuary.org/](http://na.europearchive.org/*http://finding-sanctuary.org/), last accessed on 3rd November 2014.
- MARTIN, S. and HALL-SPENCER, J.M. 2016. Effects of ocean warming and acidification on rhodolith/maerl beds. In: RIOSMENA-RODRIGUEZ, R., NELSON, W. and AQUIRRE, J. (Eds), *Rhodolith/maerl beds: A Global Perspective*. Springer, Coastal Research Library.
- MILLETT, F.W. 1885. The recent foraminifera of Mounts Bay. *Report and Transactions of the Penzance Natural History and Antiquarian Society*, **1885**, 26-28.
- MORHANGE, C., BLANC, F., SCHMITT-MERCURY, S., BOURCIER, M., CARBONEL, P., OBERLIN, C., PRONE, A., VIVENT, D. and HESNARD, A. 2003. Stratigraphy of late-Holocene deposits of the ancient harbour of Marseilles, southern France. *The Holocene*, **13**, 593-604.
- MURRAY, J.W. 1965. On the Foraminifera of the Plymouth Region. *Journal of the Marine Biological Association of the United Kingdom*, **45**, 481-505.
- MURRAY, J.W. 1970. Living foraminifera of the Western Approaches to the English Channel. *Micropaleontology*, **16**, 471-485.
- MURRAY, J.W. 1971. *An Atlas of British Recent Foraminiferids*. Heinemann Educational Books, London.
- OLUGBODE, O.I., HART, M.B. and STUBBLES, S.J. 2005. Foraminifera from Restronguet Creek: monitoring recovery from the Wheal Jane pollution incident. *Geoscience in south-west England*, **11**, 82-92.
- PASCELLI, C., RIUL, P., RIOSMENA-RODRÍGUEZ, R., SCHERNER, F., NUNES, M., HALL-SPENCER, J.M., CABRAL DE OLIVEIRA, E. and HORTA, P. 2013. Seasonal and depth-driven changes in rhodolith bed structure and associated macroalgae off Arvoredo Island (southeastern Brazil). *Aquatic Botany*, **111**, 62-65.
- PEÑA, V., BÁRBARA, I., GRALL, J., MAGGS, C.A. and HALL-SPENCER, J.M. 2014. The diversity of seaweeds on maerl in the NE Atlantic. *Marine Biodiversity*. DOI 10.1007/s12526-014-0214-7.
- PIRRIE, D., POWER, M.R., ROLLINSON, G., CAMM, G.S., HUGHES, S.H., BUTCHER, A.R. and HUGHES, P. 2003. The spatial distribution of arsenic, copper, tin and zinc within the surface sediments of the Fal Estuary, Cornwall, UK. *Sedimentology*, **50**, 579-595.
- ROSSET-MOULINIER, M. 1972. Etude des foraminifères des côtes nord et ouest de Bretagne. *Travaux du Laboratoire de Géologie, Ecole Normale Supérieure, Paris*, **6**, 1-225.
- SADRI, S., HART, M.B. and SMART, C.W. 2011. Foraminifera from the sea grass communities of the proposed Marine Conservation Zone in Tor Bay. *Geoscience in South-West England*, **12**, 269-277.
- SHEEHAN, E.V., BRIDGER, D., COUSENS, S.L. and ATTRILL, M.J. 2015. Testing the resilience of dead maerl infaunal assemblages to the experimental removal and re-lay of habitat. *Marine Ecology Progress Series*, **535**, 117-128.



- STEWART, N.F. 1996. Stoneworts – connoisseurs of clean water. *British Wildlife*, **8**, 92-99.
- STEWART, N.F. 2004. *Important Stonewort Areas. An assessment of the best areas for stoneworts in the United Kingdom (summary)*. Plantlife International, Salisbury.
- STEWART, N.F. and CHURCH, J.M. 1992. *Red Data Books of Britain and Ireland: stoneworts*. Joint Nature Conservation Committee, Peterborough.
- STUBBLES, S.J. 1993. Recent benthic foraminifera as indicators of pollution in Restronguet Creek, Cornwall. *Proceedings of the Ussher Society*, **8**, 200-204.
- STUBBLES, S.J. 1999. *Responses of recent benthic foraminifera to metal pollution in South West England estuaries: a study of impact and change*. Unpublished PhD thesis, Plymouth University.
- STUBBLES, S.J., GREEN, J.C., HART, M.B. and WILLIAMS, C.L. 1996. Response of foraminifera to the presence of heavy metal contamination and acidic mine drainage. In: *Minerals, Metals and the Environment II, Prague*. Institute of Mining and Mineralogy, London, Special Publication, 217-235.

## APPENDIX: TAXONOMIC NOTES ON FORAMINIFERA

The species mentioned in the text are well-known from UK near-shore marine environments and a full taxonomy is not presented. The species are listed in alphabetical (not taxonomic) order. Note that the taxonomic references are not given in the reference list (above).

- Ammonia batava* (Hofker) = *Streblus batavus* Hofker, 1951.
- Ammonia* sp. cf. *A. aberdoveyensis* Haynes = *Ammonia aberdoveyensis* Haynes, 1973.
- Astigerinata mamilla* (Williamson) = *Rotalina mamilla* Williamson, 1858.
- Cibicides lobatulus* (Walker and Jacob) = *Nautilus lobatulus* Walker and Jacob, 1798.
- Cribrostomoides jeffreysii* (Williamson) = *Nonionina jeffreysii* Williamson, 1858.
- Elphidium crispum* (Linné) = *Nautilus crispus* Linné, 1758.
- Elphidium williamsoni* Haynes, 1973.
- Haynesina germanica* (Ehrenberg) = *Nonionina germanica* Ehrenberg, 1840.
- Jadammina macrescens* (Brady) = *Trochammina inflata* (Montagu) var. *macrescens* Brady, 1870.
- Miliammina fusca* (Brady) = *Quinqueloculina fusca* Brady, 1870.
- Oolina squamosa* (Montagu, 1803) = *Vermiculum squamosum* Montagu, 1803.
- Patellina corrugata* Williamson, 1858
- Quinqueloculina bicornis* (Walker and Jacob) = *Serpula bicornis* Walker and Jacob, 1798.
- Quinqueloculina cliarensis* (Heron-Allen and Earland) = *Miliolina cliarensis* Heron-Allen and Earland, 1930.
- Reopbax moniliformis* Siddal, 1886.
- Rosalina globularis* d'Orbigny, 1826.
- Sejunctella earlandi* Loeblich and Tappan, 1957
- Trochammina inflata* (Montagu) = *Nautilus inflatus* Montagu, 1808.
- Trochammina ocracea* (Williamson, 1858) = *Rotalina ocracea* Williamson, 1858.