# University of Plymouth

## PEARL

01 University of Plymouth Research Outputs

https://pearl.plymouth.ac.uk

University of Plymouth Research Outputs

2017-03-22

1 Characterisation of microsatellites for *Litoria nannotis* (Amphibia: Hylidae), an

2 endangered waterfall frog endemic to the Australian Wet Tropics

3

4 Robert Puschendorf<sup>1,2</sup>, Erica V. Todd<sup>3</sup> and Michael G. Gardner<sup>4</sup>

5 School of Biological Sciences, Plymouth University, Drake Circus, Plymouth, Devon

6 PL48AA

7 <sup>2</sup>College of Marine and Environmental Sciences, James Cook University, Townsville, QLD

- 8 4811, Australia
- 9 3Department of Anatomy, University of Otago, Dunedin, New Zealand
- 10 4School of Biological Sciences, Flinders University, GPO Box 2100, Adelaide 5001 South

11 Australia, Australia; Australian Centre for Evolutionary Biology and Biodiversity, University

12 of Adelaide, 5005 Adelaide, South Australia, Australia; and Evolutionary Biology Unit,

13 South Australian Museum, North Terrace Adelaide 5000 South Australia, Australia.

14 Corresponding author. Email: robert.puschendorf@plymouth.ac.uk

15

### 16 Abstract

17 Litoria nannotis is an endangered waterfall frog from the wet tropics region in north 18 Queensland which has suffered significant population declines due to the emerging fungal disease known as chytridiomycosis. The species has two deeply divergent lineages, and we 19 20 used 454 shotgun sequencing of DNA extracted from one individual of the northern lineage to identify and design PCR primers for 576 microsatellite loci. Thirty markers were tested for 21 amplification success and variability in a population sample from each lineage. Of these, 17 22 23 were found to be polymorphic in the northern lineage and 10 loci were polymorphic in the 24 southern lineage. Numbers of alleles per locus ranged from 2 to 14 (mean 6.47, SD 4.02) for the northern lineage (17 polymorphic loci), and from 2 to 8 (mean 5.40, SD 2.55) in the 25 southern lineage (10 polymorphic loci). Levels of heterozygosity were high in both lineages 26 (northern mean  $H_E = 0.63$ , SD 0.21, range 0.27-0.89; southern mean  $H_E = 0.57$ , SD 0.25, 27 28 range 0.18-0.81). These loci will be useful in understanding the genetic variation and connectivity amongst populations of this species recovering from mass population declines 29

30 due to disease.

31 Keywords: *Litoria nannotis*; waterfall frog; Australian Wet Tropics; microsatellites; 454
32 GSFLX; shotgun sequencing; populations declines

33

34 The waterfall frog (Litoria nannotis) is an endangered species from the Australian Wet 35 Tropics. High elevation populations declined significantly in the early 1990's due to 36 the emergence of the fungal disease known as chytridiomycosis caused by the fungus 37 Batrachochytrium dendrobatidis (Berger et al. 1998), but lowland populations persisted 38 (Richards et al. 1993). Litoria nannotis is part of the torrent frog group comprised of four species, two of which were feared extinct during the declines (Richards et al. 1993). All 39 40 species in this group seem to have a similar biology (Cunningham 2001), and understanding 41 population dynamics and potential gene flow between high and low elevations as well as 42 between dry and wet forest sites is crucial when designing conservation strategies for these 43 amphibians in this system. This species is comprised of at least two distinct lineages, product 44 of historical climatic shifts and expansions and contractions in their habitat (Schneider et al. 45 1998; Cunningham 2002; Bell et al. 2011). Knowledge of current and recent historical 46 population structure, gene flow and levels of genetic diversity is especially pertinent for L. *nannotis*, as some higher elevation populations are showing some signs of recovery 47 48 (Puschendorf et al. 2011).

49

We isolated genomic DNA (1 µg) from liver of one individual *Litoria nannotis* from the northern lineage (16.466291°N, 145.152538°W, WGS84, 668 m elev) using a DNeasy spin column tissue extraction kit (Qiagen) and following manufacturers instructions. DNA was then sent to the Australian Genomic Research Facility (AGRF) in Brisbane Australia for shotgun sequencing on a Titanium GS-FLX (454 Life Sciences/Roche FLX) following 55 Gardner et al. (2011). The sample occupied 12.5 % of a plate and produced 110,205 56 individual sequences, with an average fragment size of 314.2 (Stdev 132.2). Raw sequences are available on DRYAD (doi: 10.5061/dryad.jd183; Meglécz et al. 2012). We 57 58 used the program QDD v. 1.3 (Meglecz et al., 2010) to screen the raw sequences for> eight di-, tetra- or penta-base repeats, and to remove redundant sequences and design primers for 59 60 PCR amplification of products 80-480 base pairs (automated in QDD using Primer3; Rozen & Skaletsky 2000). We identified 576 in silico microsatellite loci and ordered primer pairs 61 62 for 30 of these. Initially, the loci were trialed for amplification success in eight individuals 63 four from each lineage using the Type-it microsatellite PCR kit (Qiagen). We performed amplifications in 10 µl reactions, containing 20–50 ng template, 1x Type-it Multiplex PCR 64 65 Master Mix (Qiagen) and 0.2 µM each primer (forward and reverse). Indirectly labelled 66 reactions contained a tailed forward primer and a reporter primer (5' labelled with fluorescent dye modification HEX, TET or FAM) at a 1:4 ratio (total =  $0.2 \mu$ M). PCR 67 cycling conditionswere as follows: initial 5 min denaturation at 95°C, followed by 28 68 69 cycles of 95°C for 30 s (denaturation)/58°C for 90 s (annealing)/72°C for 30 s (extension), 70 with a final extension 30 min at 60°C. Following visualization by electrophoresis through 71 a 1.5% agarose gel, loci exhibiting reliable amplification of a single product of expected size 72 were assessed for polymorphism. We separated DNA fragments on a MegaBACE 1000 capillary sequencer and sized with GeneMarker v 2.2 software (SoftGenetics) using a 400 73 74 base pair DNA ladder as internal size standard.

75

For all polymorphic loci, forward primers were synthesised with a 5' flourescent tag: FAM
(GeneWorks), NED, PET or VIC (Applied Biosystems). Loci were then screened for
variation in 44 individuals from a single locality within the northern *L. nannotis* lineage
(16.236250 °N; 144.935690°W, WGS84, 959 m asl) and 40 individuals collected from a

single locality representing the southern lineage (18.992422°N, 146.191184°W, WGS84, 80 742nm asl; Table 1). We used the same PCR conditions and allele scoring software described 81 82 above, with allele binning to ensure consistent scoring across genotyping runs. Due to 83 consistent differences in allele profiles among lineages, independent scoring panels were used 84 for each lineage. Multiplex PCR combinations (Table 1) were later designed *in silico* with the 85 aid of MULTIPLEX MANAGER 1.0 software (Holleley and Geerts 2009), and tested using 86 PCR conditions described above. Characteristics of each locus in each lineage are 87 summarised in Table 1. Data are presented for 19 loci that amplified consistently in the 88 northern lineage, and similarly for 17 loci in the southern lineage. Basic summary statistics (number of alleles, observed and expected heterozygosities) were calculated in GENALEX 89 6.5 (Peakall and Smouse 2012), which was also used to test for deviations from Hardy-90 91 Weinberg Equilibrium (HWE). Polymorphic Information Content (PIC) values were 92 calculated for each locus in CERVUS (Kalinowski et al. 2007). Potential linkage 93 disequilibrium (LD) between pairs of loci was investigated using GENEPOP 4.2 online, with 94 10,000 iterations (http://genepop.curtin.edu.au/; Raymond and Rousset 1995; Rousset 2008) (Table 1). P values from HWE and LD tests were adjusted for multiple tests of significance 95 using the false discovery rate (FDR) correction and included in Table 1. (Benjamini and 96 97 Hochberg 1995). We used MICROCHECKER 2.2.3 (Van Oosterhout et al. 2004) to check 98 each locus for evidence of null alleles, scoring error due to stuttering, and large allele drop 99 out, using a 95% confidence level and 10,000 iterations. 100 In the northern lineage, 17 of 19 polymorphic loci conformed to HWE expectations and are 101 considered suitable for population genetic studies (bold in table 1). In the southern lineage, 10 102 of 17 polymorphic loci met HWE expectations. Of those loci not in HWE, there was evidence 103 for null alleles at locus Lnan15 in the northern lineage, and Lnan17 and Lnan25 in the 104 southern lineage. There was no evidence of large allele drop out at any locus. Following FDR

105 correction, all loci were found to be inherited independently (North P >0.002, FDR value 106 0.0003; South P > 0.02, FDR value 0.0006). Overall, the markers exhibit high levels of polymorphism in northern and southern L. nannotis lineages suitable for studies of 107 108 relatedness, population genetic structure and connectivity. For polymorphic loci also in HWE, numbers of alleles per locus ranged from 2 to 14 (mean 6.47, SD 4.02) for the northern 109 110 lineage (17 polymorphic loci), and from 2 to 8 (mean 5.40, SD 2.55) in the southern lineage (10 polymorphic loci). Levels of heterozygosity were high in both lineages (northern mean 111 112  $H_E = 0.63$ , SD 0.21, range 0.27-0.89; southern mean  $H_E = 0.57$ , SD 0.25, range 0.18-0.81). 113 Overall, the markers exhibit high levels of polymorphism in northern and southern L. 114 *nannotis* lineages suitable for studies of relatedness, population genetic structure and 115 connectivity. 116 117 These markers will be used to document patterns of gene flow, population structure and 118 genetic diversity in *L. nannotis* and to investigate their recovery from the amphibian 119 population declines linked to chytridiomycosis documented since the early 1990's (Berger et al. 1998). More recently, high elevation populations seem to be recovering, and 120 121 larger seemingly healthy populations have been described in the western slopes of the wet 122 tropics region, including one sister species, *Litoria lorica* which was previously thought to be 123 extinct (Puschendorf et al. 2011). How these populations are interconnected and the 124 source of the recovering populations is a key aspect of frog conservation in this region.

125

## 126 Acknowledgements

127 David Blair, Martin van der Meer, Blanche Danastas and Giana Gomez provided help in the
128 laboratory as well as in the design of these primers . We acknowledge the assistance and

129	support of Alison Fitch from Flinders University. This project was funded by the Australian
130	Research Council (DP1094540).
131	References
132	
133	Benjamini Y., and Hochberg Y. (1995). Controlling the false discovery rate: a practical and
134	powerful approach to multiple testing. Journal of the Royal Statistical Society. Series B
135	(Methodological) <b>57</b> , 289–300.
136	
137	Bell, R. C., MacKenzie, J. B., Hickerson, M. J., Chavarría, Krystle L., Cunningham, M.,
138	Williams, S. and Moritz, C. (2012). Comparative multi-locus phylogeography confirms
139	multiple vicariance events in co-distributed rainforest frogs. Proceedings of the Royal Society
140	B: Biological Sciences 279: 991-999. doi: 10.1098/rspb.2011.1229
141	
142	Berger, L., Speare, R., Daszak, P., Green, D. E., Cunningham, A. A., Goggin, C. L.,
143	Slocombe, R., Ragan, M. A., Hyatt, A. D., McDonald, K. R., Hines, H. B., Lips, K. R.,
144	Marantelli, G, and Parkes, H. (1998). Chytridiomycosis causes amphibian mortality
145	associated with population declines in the rainforests of Australia and Central America.
146	Proceedings of the National Academy of Science USA 95: 9031-9036. doi:
147	<u>10.1073/pnas.95.15.9031</u>
148	
149	Cunningham, M. (2002). Identification and evolution of Australian torrent treefrogs (Anura:
150	Hylidae: Litoria nannotis group). Memoirs of the Queensland Museum 48: 93-102.
151	
152	Gardner M.G., Fitch A.J., Bertozzi T, and Lowe, A.J. (2011). Rise of the machines -
153	recommendations for ecologists when using next generation sequencing for microsatellite
154	development. Molecular Ecology Resources 11, 1093-1101. doi: 10.1111/j.1755-
155	0998.2011.03037.x
156	
157	Holleley C.E., and Geerts P.G. (2009). MULTIPLEX MANAGER 1.0: a cross-platform
158	computer program that plans and optimizes multiplex PCR. Biotechniques 46:511-517.doi:
159	10.2144/000113156
160	

161	Kalinowski S., Taper M., and Marshall T. (2007) Revising how the computer program
162	CERVUS accommodates genotyping error increases success in paternity assignment.
163	<i>Molecular Ecology</i> <b>16</b> , 1099-1006.doi: 10.1111/j.1365-294x.2007.03089.x
164	
165	Meglécz E., (2007). Microfamily (version 1): a computer program for detecting flanking-
166	region similarities among different microsatellite loci. Molecular Ecology Notes 7, 18-20.
167	doi: 10.1111/j.1471-8286.2006.01537.x
168	
169	Meglecz E., Costedoat C., Dubut V., Gilles A., Malausa T., Pech N., and Martin JF. (2010).
170	QDD: a user-friendly program to select microsatellite markers and design primers from large
171	sequencing projects. Bioinformatics, 26, 403-404.doi: 10.1093/bioinformatics/btp670
172	
173	Meglécz E., Nève G., Biffin E. and Gardner M.G. (2012) Breakdown of Phylogenetic Signal:
174	A Survey of Microsatellite Densities in 454 Shotgun Sequences from 154 Non Model
175	Eukaryote Species. PLoS ONE 7, e40861.doi: 10.1371/journal.pone.0040861
176	
177	Peakall, R., and Smouse P.E. (2012). GenAlEx 6.5: genetic analysis in Excel. Population
178	genetic software for teaching and research-an update. Bioinformatics 28, 2537-2539.doi:
179	10.1093/bioinformatics/bts460
180	
181	Puschendorf, R., Hoskin, C.J., Cashins, S.D., McDonald, K., Skerratt, L. F., Vanderwal, J.,
182	and Alford, R. A. (2011). "Environmental Refuge from Disease-Driven Amphibian
183	Extinction." Conservation Biology 25, 956-964.doi: 10.1111/j.1523-1739.2011.01728.x
184	
185	Raymond M., and Rousset F. 1995. GENEPOP (version 1.2): Population genetics software
186	for exact tests and ecumenicism. Journal of Heredity, 86, 248-249.
187	
188	Richards, S. J., McDonald, K. R., Alford, R. A (1993). Declines in populations of Australia's
189	endemic tropical rainforest frogs. Pacific Conservation Biology 1: 66-77. doi:
190	10.1071/pc930066
191	
192	Rousset, F., 2008. Genepop'007: a complete reimplementation of the Genepop software for
193	Windows and Linux. Molecular Ecology Resources 8, 103-106.doi: 10.1111/j.1471-
194	8286.2007.01931.x

195

- 196 Rozen S., and Skaletsky H.J. (2000). Primer3 on the WWW for general users and for
- 197 biologist programmers. In 'Bioinformatics Methods and Protocols: Methods in Molecular
- 198 Biology' (eds. Krawetz S, Misener S), pp. 365-386. (Humana Press: Totowa). doi: 10.1385/1-
- 199 <u>59259-192-2:365</u>
- 200
- 201 Schneider, C. J., Cunningham, M. and Moritz, C. (1998). Comparative phylogeography and
- 202 the history of endemic vertebrates in the Wet Tropics rainforests of Australia. *Molecular*
- 203 *Ecology* **7**: 487-498. doi: 10.1046/j.1365-294x.1998.00334.x
- 204
- 205 Van Oosterhout C., Hutchinson B., Wills D. and Shipley P. (2004). MICRO-CHECKER:
- software for identifying and correcting genotyping errors in microsatellite data. *Molecular*
- 207 Ecology Notes 4, 535–538. doi: 10.1111/j.1471-8286.2004.00684.x
- 208
- 209
- 210 Table 1. Details for 19 *Litoria nannotis* microsatellite loci developed from 454 shotgun
- 211 sequence data. Loci in bold are in Hardy-Weinberg equilibrium.

NORTHERN

			Primer										
			conc.									Multiplex	Genbank
Locus	Primer sequence 5' ti 3'	Repeat Motif	(μM)	Ta(°C)	Ν	Allele size range	Na	HO	HE	PIC	P HWE*	group	accession no.
Lnan03	F:GCCATGCACATGAGCTTTTA	(AT)8	0.2	58	44	140-142	2	0.568	0.500	0.375	0.364	4	KX518722
	R: CCAATACGCGCCAATTTTAC												
Lnan04	F: GGTGGACATCATGTGGATCA	(AT)8	0.2	58	44	190-192	2	0.068	0.107	0.101	0.016	5	KX518723
	R: CCAATACGCGCCAATTTTAC												
Lnan06	F: GAGTTTCCTTCCCAAAAGCA	(TG)9	0.2	58	44	100-106	3	0.250	0.271	0.24	0.118	5	KX518724
	R: GCATCAATCCCTGTCTCCAA												
Lnan08	F: GTATAACAGGGCGGAACTGC	(GT)9	0.2	58	44	131-139	4	0.727	0.667	0.611	0.644	2	KX518725
	R: GTGTAACTCGCCTTCCTTGC												
Lnan10	F: TGTGTAAATTGCTCCAGGCA	(AT)11	0.2	58	44	140-184	10	0.750	0.761	0.734	0.654	4	KX518726
	R: TGAATGATGCCAGACCAAGA												
Lnan14	F: GCAACCAATATGGGTGACATT	(AT)12	0.2	58	44	210-216	4	0.591	0.582	0.504	0.285	5	KX518728
	R: GCACTTATGTTGCGATGCAC												
Lnan15	F: TGCAGATCCATGCAATACTGA	(AAT)8	0.2	58	44	149-167	7	0.636	0.774	0.74	0.021	1	KX518729
	R: TCAACGTTCAATGGTCAAGG												
Lnan16	F: ACTTTGTTAGGTGCTGCGGA	(AAT)8	0.2	58	43	103-109	2	0.419	0.381	0.308	0.514	3	KX518730
	R: GCACCCTTAATGTGTTCCTGA												
Lnan17	F: GCGGTTACAGGGTACAGCAT	(TTA)8	0.2	58	44	207-219	4	0.432	0.440	0.377	0.960	1	KX518731
	R: TGTACTTTGTTAGGCGCTGC												
Lnan18	F: CCAAAACCGCTTTTCTGTTG	(CTA)8	0.2	58	44	136-142	2	0.386	0.363	0.297	0.675	2	KX518721
	R: TGGGTTAATAACATGAGGAAGAGTT												
Lnan20	F: AAGTGCTCCGGATACCAATG	(TAT)11	0.2	58	43	285-294	4	0.721	0.653	0.589	0.466	3	KX518720
	R: TTGTTGATGAATCTGGTGCC												
Lnan21	F:TACTTTGTTAGTCGCTGCGG	(ATT)12	0.2	58	44	124-136	4	0.386	0.326	0.296	0.866	4	KX857664
	R:CTCTTGTTGGCCTCCCATAA												
Lnan22	F: CAAGGTTGACACCAAGCAGA	(TTA)12	0.2	58	44	107-134	7	0.864	0.808	0.781	0.519	1	KX518732
	R: TGTAACTTTGTTAGGCGCTGC												
Lnan24	F: GCCATTTAAGACACCTGGGA	(ATCT)12	0.2	58	43	136-170	9	0.884	0.858	0.841	0.771	3	KX518733
	R: CCATTGTGTGCTGCAGTGAT												
Lnan25	F:TAAGGGGATTGGTATGCTGG	(CTAT)13	0.2	58	44	155-187	9	0.818	0.793	0.771	0.441	5	KX857663
	R:GAAGTGCCACTACCATTCTTTTG												
Lnan26	F: CTTTCACGTCATAGGAACCCA	(GATA)13	0.2	58	43	133-171	12	0.837	0.839	0.822	0.997	3	KX518734
	R: CAACAGGGCTTTCAACCATT												
Lnan27	F: CCACTCCTGTTGGGGAGATA	(GATA)14	0.2	58	44	81-159	9	0.886	0.839	0.821	0.081	1	KX518719
	R: AAATGTGGGAAAAGTGAAGCA												
Lnan29	F: CTATGCGGCCATCTTCTCTC	(ATCT)17	0.2	58	44	178-249	13	0.909	0.894	0.885	0.499	4	KX518735
	R: GTGACTTGCAGCCTGTTGAG												
Lnan30	F: GTGAAAAGCAATGCCACCTT	(ATCT)17	0.2	58	43	127-210	14	0.791	0.860	0.847	0.266	2	KX518736
	R: TCAGTAGACCACAAAGAGCGTT												

#### SOUTHERN

			Primer										
			conc.									Multiplex	Genbank
Locus	Primer sequence 5' ti 3'	Repeat Motif	(μM)	Ta(°C)	Ν	Allele size range	Na	HO	HE	PIC	P HWE*	group	accession no.
Lnan03	F:GCCATGCACATGAGCTTTTA	(AT)8	0.2	58	40	140-142	2.000	0.200	0.180	0.164	0.482	4	KX518722
	R: CCAATACGCGCCAATTTTAC												
Lnan04	F: GGTGGACATCATGTGGATCA	(AT)8	0.2	58	39	192	1.000	NA	NA	NA	NA	5	KX518723
	R: CCAATACGCGCCAATTTTAC												
Lnan08	F: GTATAACAGGGCGGAACTGC	(GT)9	0.2	58	40	131	1.000	NA	NA	NA	NA	2	KX518725
	R: GTGTAACTCGCCTTCCTTGC												
Lnan10	F: TGTGTAAATTGCTCCAGGCA	(AT)11	0.2	58	40	139-162	4.000	0.575	0.641	0.574	0.115	4	KX518726
	R: TGAATGATGCCAGACCAAGA												
Lnan12	F: TCAAATCCATTGTGGTGGTG	(TA)11	0.2	58	40	191-221	8.000	0.700	0.681	0.631	0.997	2	KX518727
	R: CCACATGTTGCCTACTCCCT												
Lnan14	F: GCAACCAATATGGGTGACATT	(AT)12	0.2	58	39	206-232	6.000	0.718	0.673	0.624	0.198	5	KX518728
	R: GCACTTATGTTGCGATGCAC												
Lnan15	F: TGCAGATCCATGCAATACTGA	(AAT)8	0.2	58	39	148	1.000	NA	NA	NA	NA	1	KX518729
	R: TCAACGTTCAATGGTCAAGG												
Lnan16	F: ACTTTGTTAGGTGCTGCGGA	(AAT)8	0.2	58	39	112-127	5.000	0.538	0.617	0.583	0.228	3	KX518730
	R: GCACCCTTAATGTGTTCCTGA												
Lnan17	F: GCGGTTACAGGGTACAGCAT	(TTA)8	0.2	58	40	210-213	3.000	0.100	0.184	0.174	<0.001	1	KX518731
	R: TGTACTTTGTTAGGCGCTGC												
Lnan18	F: CCAAAACCGCTTTTCTGTTG	(CTA)8	0.2	58	40	133-136	2.000	0.200	0.180	0.164	0.482	2	KX518721
	R: TGGGTTAATAACATGAGGAAGAGTT												
Lnan20	F: AAGTGCTCCGGATACCAATG	(TAT)11	0.2	58	39	273-283	3.000	0.359	0.325	0.296	0.710	3	KX518720
	R: TTGTTGATGAATCTGGTGCC												
Lnan21	F:TACTTTGTTAGTCGCTGCGG	(ATT)12	0.2	58	40	121	1.000	NA	NA	NA	NA	4	KX857664
	R:CTCTTGTTGGCCTCCCATAA												
Lnan24	F: GCCATTTAAGACACCTGGGA	(ATCT)12	0.2	58	39	123-145	6.000	0.718	0.739	0.705	0.023	3	KX518733
	R: CCATTGTGTGCTGCAGTGAT												
Lnan25	F:TAAGGGGATTGGTATGCTGG	(ATCT)12	0.2	58	37	142-224	13.000	0.676	0.874	0.861	0.005	5	KX857663
	R:GAAGTGCCACTACCATTCTTTTG												
Lnan26	F: CTTTCACGTCATAGGAACCCA	(GATA)13	0.2	58	39	121-151	8.000	0.744	0.811	0.787	0.508	3	KX518734
	R: CAACAGGGCTTTCAACCATT												
Lnan27	F: CCACTCCTGTTGGGGAGATA	(GATA)14	0.2	58	39	106-138	8.000	0.769	0.812	0.786	0.862	1	KX518719
	R: AAATGTGGGAAAAGTGAAGCA												
Lnan30	F: GTGAAAAGCAATGCCACCTT	(ATCT)17	0.2	58	40	123-153	8.000	0.775	0.814	0.789	0.414	2	KX518736
	R: TCAGTAGACCACAAAGAGCGTT												

\*Lnan17, Lnan25 significant after FDR correction, FDR value 0.012