



UNIVERSITY OF
PLYMOUTH



School of Nursing and Midwifery
Faculty of Health

2020-03-31

A systematic review of effectiveness of interventions applicable to radiotherapy that are administered to improve patient comfort, increase patient compliance, and reduce patient distress or anxiety

S. Goldsworthy

S. Palmer

J. M. Latour *School of Nursing and Midwifery*

H. McNair

M. Cramp

Let us know how access to this document benefits you

General rights

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

Take down policy

If you believe that this document breaches copyright please [contact the library](#) providing details, and we will remove access to the work immediately and investigate your claim.

Follow this and additional works at: <https://pearl.plymouth.ac.uk/nm-research>

Recommended Citation

Goldsworthy, S., Palmer, S., Latour, J., McNair, H., & Cramp, M. (2020) 'A systematic review of effectiveness of interventions applicable to radiotherapy that are administered to improve patient comfort, increase patient compliance, and reduce patient distress or anxiety', *Radiography*, . Available at: <https://doi.org/10.1016/j.radi.2020.03.002>

This Article is brought to you for free and open access by the Faculty of Health at PEARL. It has been accepted for inclusion in School of Nursing and Midwifery by an authorized administrator of PEARL. For more information, please contact openresearch@plymouth.ac.uk.

A systematic review of interventions administered to improve patient comfort, assist with completion of a clinical procedure and reduce patient distress or anxiety applicable to radiotherapy: an evaluation of effectiveness.

Simon Goldsworthy MSc^{1,2}, Shea Palmer PhD², Jos M. Latour PhD, RN³, Helen McNair PhD⁴, Mary Cramp PhD²

Journal Title: Radiography

Acceptance Date: 5 March 2020

1. Radiotherapy, Beacon Centre, Musgrove Park Hospital, Taunton and Somerset NHS Foundation Trust, Taunton, United Kingdom.
2. Faculty of Health and Applied Sciences, University of the West of England, Bristol, United Kingdom.
3. Faculty of Health and Human Sciences, University of Plymouth, Plymouth, United Kingdom
4. Royal Marsden NHS Foundation Trust and Institute of Cancer Research, Sutton, United Kingdom.

Correspondence: simon.goldsworthy@tst.nhs.uk

Conflict of interest: None to declare

Funding: College of Radiographers Doctoral Fellowship 008

Key words: Comfort interventions, radiotherapy, randomised controlled trial, systematic review, clinical significance

Acknowledgements:

We would like to acknowledge three librarians; Pauline Shaw at the University of the West of England, Carol-Ann Regan and Jessica Pawley at Taunton and Somerset NHS Trust for supporting the search strategy development. We would also like to acknowledge the College of Radiographers Industrial Partnerships Doctoral Fellowship Scheme for supporting the primary author to deliver this research as part of his Ph.D.

ABSTRACT

Objectives

The aim of this review was to search existing literature to identify comfort interventions that can be used to assist an adult patient to undergo complex radiotherapy requiring positional stability for periods greater than 10 minutes. The objectives of this review were to; 1) identify comfort interventions used for clinical procedures that involve sustained inactivity similar to radiotherapy; 2) define characteristics of comfort interventions for future practice; and 3) determine the effectiveness of identified comfort interventions. The Preferred Reporting Items for Systematic Reviews and meta-analyses statement and the Template-for-Intervention-Description-and Replication guide were used.

Key findings

The literature search was performed using PICO criteria with five databases (AMED, CINAHL EMBASE, MEDLINE, PsycINFO) identifying 5,269 titles. After screening, 46 randomised controlled trials met the inclusion criteria. Thirteen interventions were reported and were grouped into four categories: Audio-visual, Psychological, Physical, and Other interventions (education/information and aromatherapy). The majority of aromatherapy, one audio-visual and one educational intervention were judged to be clinically significant for improving patient comfort based on anxiety outcome measures (effect size ≥ 0.4 , mean change is greater than the Minimal-Important-Difference and low-risk-of-bias). Medium to large effect sizes were reported in many interventions where differences did not exceed the Minimal-Important-Difference for the measure. These interventions were deemed worthy of further investigation.

Conclusion

Several interventions were identified that may improve comfort during radiotherapy assisting patients to sustain and endure the same position over time. This is crucial for the continual growth of complex radiotherapy requiring a need for comfort to ensure stability for targeted treatment.

Implications for practice

Further investigation of comfort interventions is warranted, including tailoring interventions to patient choice and determining if multiple interventions can be used concurrently to improve effectiveness.

INTRODUCTION

Positioning and immobilisation of patients are crucial for reproducible and accurate delivery of radiotherapy in both radical and palliative settings to ensure tumour control while avoiding healthy tissue toxicity¹⁻². Recent studies have shown that comfort in patients receiving radiotherapy for prostate cancer can be determined by treatment position³ and a strong association was observed between comfortable patient positioning and improved treatment accuracy in patients' receiving radiotherapy for breast cancer⁴. As more complex treatment techniques like stereotactic ablative body radiotherapy (SABR) becomes standard, and treatment times are extended above 10 minutes, the comfort of patients is an important consideration⁵⁻⁶. It is also hypothesized that there is an association between patient comfort and radiotherapy treatment time⁷ and one limitation to technical radiotherapy advancements is managing the patient's tolerability of immobilisation to complete the procedure while also achieving comfort⁸. Hypothetically, not providing a comfort intervention might increase the treatment time in radiotherapy.

To assist with identification and development of suitable comfort interventions, there is a need to consider what patient comfort is and means. Patient comfort is defined holistically as a state of having met the basic human needs for ease, relief, and transcendence in four contexts⁹⁻¹¹. In radiotherapy procedures the role and purpose of holistic comfort interventions aim to make the procedure more tolerable to patients and ensure compliance reducing discomfort, anxiety, distress and claustrophobia. Comfort has been explored in a few studies including a focus group of patients with head and neck cancer receiving radiotherapy². Their experiences reflected the definition of holistic comfort⁹⁻¹¹ and indicated that therapeutic radiographers may not fully appreciate their level of discomfort. A survey of 100 head and neck cancer patients who had received radiotherapy found that a quarter were anxious and that interventions were required including better patient preparation/ education¹². In UK and European guidelines, recommendations on how to manage patient comfort during radiotherapy are limited^{2,13-14}. Greater evidence of comfort intervention effectiveness is required to inform national radiotherapy practice and guidelines.

Interventions such as communication with professionals and music were reported to reduce distress in up to 86% of patients receiving radiotherapy for head and neck cancer¹⁵. A previous systematic review explored the efficacy of holistic comfort interventions during invasive paediatric nursing procedures such as venepuncture, port access and intramuscular injection¹⁶. The review grouped comfort interventions into four categories: music, amusement and entertainment, caregiver facilitation and a multifaceted approach and supported the use of various distraction methods to reduce anxiety, distress, fear and pain during procedures¹⁸. Further studies have investigated interventions ranging from music to self-hypnosis and deep breathing exercises¹⁷⁻¹⁸. Thus, there are promising procedural

comfort interventions that may be applicable to radiotherapy. A limited number of interventions have been investigated to manage patient comfort during radiotherapy¹⁹⁻²⁰. The aim of this review was to search existing literature to identify comfort interventions that can be used to support an adult patient to undergo clinical procedures that requires a patient to sustain the same position over a period greater than 10 minutes. The current estimated time cut off set at 10 minutes was deployed to capture procedures that would replicate the radiotherapy phase after positioning when patients must remain still during pre-treatment verification and treatment delivery such as SABR or palliative radiotherapy. The focus above 10 minutes was set to ensure a breadth of clinical procedures were included that would be more representative of radiotherapy. The objectives of this review were to: 1) identify comfort interventions that are used for clinical procedures that involve sustained inactivity similar to radiotherapy; 2) record the characteristics of the comfort interventions for future practice; and 3) determine the effectiveness of the comfort interventions.

METHODS

Protocol and registration

A review protocol was developed and prospectively published in PROSPERO (CRD42017059688) in line with the Centre of Reviews and Dissemination²¹.

Information sources

The review was structured and reported according to the preferred reporting items for systematic reviews and meta-analyses (PRISMA) statement²² and the Template for Intervention Description and Replication (TIDieR)²³ Guide.

Search

Five databases, AMED, CINAHL EMBASE, MEDLINE and PsycINFO, were searched to identify relevant text in titles, abstracts and key words to develop search terms. The literature search used the same databases and refined terms (electronic supplement A). The search was restricted to title and abstract fields to avoid retrieving non-related papers from the subject headings.

Selection criteria for eligible primary research was defined according to the Participant(P), Intervention(I), Control(C), Outcome(O) and Studies(S)²⁴ framework:

(P) Adult patients (≥ 18 years) undergoing a clinical procedure that required alignment, stabilisation, immobilisation and having to sustain, endure or tolerate the procedure while conscious over a period greater than 10 minutes. Clinical procedures included those where patients must remain stable and unwanted movement is critical. In the surgical and radiotherapy setting, unwanted movement could result in collateral damage such as the laceration or irradiation of surrounding normal tissue respectively and potentially poorer outcomes for patients.

- (I) Interventions to aid comfort; ease, relieve, relief, reduce distress/anxiety, relax, calm, alleviate, distract, or transcend a patient/service user immediately before or within a clinical procedure which requires alignment, stabilisation, or immobilised and has to sustain, endure or tolerate the procedure while conscious.
- (C) Usual standard of care or comparator (another intervention)
- (O) Assessments of patient comfort, psychological well-being, patient satisfaction and quality of life outcomes.
- (S) Randomised controlled trials (RCTs) and controlled clinical trials (CCTs).

Studies published in English between 2000 and January 2019 were included to focus on contemporaneous practice. The searches were initially performed in August 2017 and updated in January 2019. Following removing of duplicates, two researchers independently reviewed the titles and abstracts initially and then full texts to identify papers that met the eligibility criteria. A consensus meeting was held, and concordance was achieved on 95% of the full texts. A third reviewer arbitrated on inclusion of the remaining 5% (n=4) of full texts.

Data extraction

Data was extracted from each paper by one researcher using a data extraction form based on the TIDieR checklist and guidelines²³ and reviewed by a second researcher. The data extraction form included: authors, year of publication, study design, setting, participants, clinical procedures, outcome measures, main outcomes (measured before and after clinical procedure delivery, or as a mean change), and delivery characteristics of the comfort interventions.

Risk of bias

The Cochrane Risk of Bias Checklist (Version 5.1.0) was used to assess risk of bias (RoB) in RCTs²⁵. Six areas of ROB were assessed (random sequence generation; allocation concealment; blinding of participants and personnel; blinding of outcome assessment; incomplete outcome data; and selective reporting) with each area given either “low,” “high,” or “unclear” risk of bias²⁵. To reduce the effect of human factors in assessing RoB²⁷⁻²⁸ an online software RoBotReviewer™ which aims to semi-automate evidence synthesis using machine learning was used²⁸⁻²⁹ alongside review by the researchers. International clinical trials registers were accessed to determine selective reporting bias; if not registered then studies were judged unclear for RoB. For CCTs the RoB was assessed using Risk-of-Bias-In-Non-Randomised-Studies-of Interventions. For this review, studies were judged not acceptable if there was high risk for selection bias in both domains because randomisation is a crucial attribute of well-designed RCTs. Studies judged high risk in one area of selection bias, or another RoB component were deemed acceptable but treated with caution, and not included in the data synthesis.

Data synthesis

Only validated outcome measures were included in the synthesis and were reported separately for intervention and comparator groups. Where available, the change in outcome measures from before to after clinical procedures was calculated as mean differences, percentage change, Cohen's D effect size (normalised distribution) or r-effect size (non-normalised distribution) with 95% confidence intervals (CI)³⁰⁻³¹. Studies were selected for the Cohen's D or r-effect size analysis dependant on whether the data followed a normal distribution³²⁻³³, confirmed by the reported use of Kolmogorov-Smirnov (KS) or Shapiro-Wilk (SW) test for normality or assumed based on the use of parametric tests³⁷⁻³⁸. Where mean and standard deviations (SD) were not reported, an estimation from either inter quartile range or p-value was calculated³⁴⁻³⁷. Meta-analysis was not conducted because of the clinical heterogeneity in the study populations, healthcare settings, interventions and comparator types.

To determine whether comfort interventions make an important difference to the patients, the clinical significance of studies was assessed to supplement statistical significance³⁰⁻³². In this review, clinical significance was determined using effect size and the minimal importance difference (MID). Effect sizes were interpreted using the following criteria: small effect (≤ 0.4), medium effect ($\geq 0.5 \leq 0.7$) or large effect (≥ 0.8)³⁸. Minimal important differences (MID) of validated outcome measures were identified from the literature³⁹⁻⁴². A comfort intervention was considered to demonstrate clinical significance when the effect size exceeded 0.4, mean differences were greater than the MID and RoB was acceptable.

RESULTS

Study selection

Database searches initially identified 5269 titles (Figure 1). After removing duplicates (n=191), 5078 titles and abstracts were screened, and 4994 papers were removed leaving 84 papers for full review. Of these, 38 papers were excluded for reasons listed in Figure 1. One CCT was excluded because it used a parallel cross over design with potential for cross contamination between intervention and comparator groups. In total 46 papers were included in the review⁴³⁻⁸⁸.

Study characteristics (electronic supplement B & C).

The studies included consisted of 46 RCTs with a total of 5782 patients⁴³⁻⁸⁸. The age of participants ranged between 18 and 80 years. The study design of RCTs included; two-arm parallel, multiple arm parallel, and mixed factorial multiple/ parallel arm study designs.

Clinical procedures

Nineteen different clinical procedures were identified. The two most common clinical procedures were observational investigations such as bronchoscopy/hysteroscopy (n=14) and interventional radiology (n=13).

Outcome measures

Most studies reported an anxiety outcome measure (n=44) and 29 studies used the State-Trait Anxiety-Inventory (STAI) aligning to psychological wellbeing. The STAI examines feelings 'at the present moment' and gives a score between 20 and 80, with a higher score indicating greater anxiety levels⁸⁹. One study used a 6-item short STAI which is stated to be more sensitive to fluctuations in anxiety⁹⁰. One study used the anxiety Visual-Analogue-Scale (VAS-A)⁴¹⁻⁴², and another study used the Beck-Anxiety-Inventory (BAI) and Hamilton-Anxiety-Scale (HAS) and non-validated numeric rating scales for comfort, satisfaction, willingness to repeat and experience of the environment⁹¹. Only validated anxiety measures including the STAI, the VAS-A, the BAI and the HAS, reported before and after clinical procedures, were included in the data synthesis. For the STAI, the MID was set at 10³⁹⁻⁴⁰. The MID was set at 46 for the VAS-A⁴¹, 8.8 for the BAI and 8.2 for the HAS⁴².

Comfort interventions (electronic supplement B).

Thirteen comfort interventions were identified and grouped into the four categories (Table 1): Audio-visual, Psychological, Physical, and Other Interventions (education/information and aromatherapy). Comfort interventions were delivered before the clinical procedure in 10 studies, during the clinical procedure in 19 studies and both before and during the clinical procedure in 17 studies.

- Audio-visual technology interventions include audio only (n =20)^{43-48,55,60-61,64,66, 68,70-74,77,82,84,86}, audio-visual (n= 6)^{50,51,53,60,69,88}, virtual reality (n =2)^{67,85} and visual only (n=1)⁸⁸. The interventions were used for the purpose of improving (dis)comfort, reducing anxiety, distraction, improving well-being and relaxation. A wide range of music genres were used ranging from classical to easy listening popular music, chants and nature sounds. The delivery features ranged from music or video players, loudspeakers or earphones to headsets and goggles for virtual reality. Interventions were delivered by professionals and/or self-administered by patients.
- Psychological interventions include breathing techniques (n=1)⁸⁰, cognitive behavioural therapy (n=1)⁷⁹, distraction (n=1)⁶⁴, empathetic attention (n=4)^{49,60,65,76} and hypnosis (n=4)^{57,65,76,81}. The interventions were used for the purpose of reducing discomfort, anxiety and pain, or improving satisfaction and relaxation. The delivery features ranged from face to face to audio players. Interventions were delivered by therapists or self-administered by patients via audio players.

- Physical interventions includes massage (n=2)^{75,80}, therapeutic touch (n=1)⁵⁴, reflexology (n=2)^{56,78} and stress balls (n=1)⁶⁰. The interventions were used for the purpose of reducing discomfort, anxiety, distress and pain, or improving satisfaction. The delivery was face to face with professionals.
- Other interventions includes education/information (n=4)^{43,62-63,87} and aromatherapy (n=5)^{52, 58, 59,78,83}. The interventions were used for the purpose of improving experience and satisfaction or reducing anxiety and psychophysiological arousal/parameters. Interventions were delivered by a range of personnel and methods.

Some studies with multiple arm parallel designs investigated interventions that crossed the above categories (n=5)^{45,64,73,75,80}.

Cochrane Risk of bias for included studies

Each of the included RCTs had areas where the ROB was high, low, and unclear (Fig.2). 38% of RCTs had a low overall risk of bias. Low risk for random sequence generation and concealment was reported in 77% and 32% of studies respectively. Blinding of professionals or participants to the allocated comfort intervention was reported in 6% of studies, whilst blinding of outcome assessment was completed in 36%. 81% of RCTs were judged unclear for selective reporting because trials were not registered. 3 RCTs were deemed unacceptable due to high risk of selection bias and were not included in the data synthesis^{59, 67,69}.

Effectiveness of comfort interventions

Only anxiety outcomes were synthesised as the outcome measures were validated and reported before and after clinical procedure (Table 2). This resulted to exclude another 17 RCTs^{45-46,49,54,58,61-62, 65,70,72,74,76,78-79, 84-86}. 26 RCTs were included in the data synthesis.

Audio-visual technology interventions includes studies of audio alone^{44, 47-48, 50, 55, 60, 64, 66, 68, 71,73,77, 82, 88}, audio-visual^{50-51, 53, 60} and visual⁸⁸ interventions with data available for synthesis.

Audio: six out of eleven studies of audio interventions reported statistical significance favouring the intervention ($p < 0.05$)^{44, 47, 55, 60, 73}. The mean difference in anxiety exceeded the MID in one intervention and with a medium effect size was judged clinically significant⁷¹. Medium to large effect sizes were observed in all eleven studies.

Audio-visual: three out of four audio-visual interventions studies reported statistically significance favouring the intervention ($p < 0.05$)^{51, 53, 60}. The mean difference in anxiety exceeded the MID in two studies⁵⁰⁻⁵¹; one had a small effect size⁵¹ and one favoured the comparator group⁵⁰. Medium to large effect sizes were observed in all other studies^{50, 51, 53, 60, 88}.

Visual: one visual intervention study favoured the intervention statistically ($p < 0.05$)⁸⁸. The mean difference in anxiety did not exceed the MID but had a large effect size⁸⁸. Only one study investigating music interventions was deemed clinically significant⁷¹.

Psychological interventions with data available for synthesis included distraction⁶⁴, empathetic attention⁶⁰ and hypnosis^{57, 81} interventions.

Distraction: one study did not show a statistically significant effect for distraction intervention⁶⁴. The difference in mean anxiety did not exceed the MID⁶⁴, and the effect size favoured the comparator group.

Empathetic attention: one study reported statistical significance favouring the intervention⁶⁰ ($p < 0.05$). The mean difference in anxiety did not exceed the MID, and while it had a large effect size, it was deemed not clinically significant.

Hypnosis: two studies reported statistical significance favouring hypnosis interventions^{57, 81} ($p < 0.05$). Both had large effect sizes but the mean difference in anxiety did not exceed the MID in either study^{57, 81}.

No intervention in this category was considered clinically significant.

Physical interventions were used in three studies with data available for synthesis and involved physical touch: reflexology⁵⁶, massage⁷⁵, and stress balls⁶⁰.

Two out of three studies reported statistical significance favouring the intervention ($p < 0.05$)^{56, 60}. The mean difference in anxiety exceeded the MID in one study⁷⁵ with large effect sizes in the other two^{56, 60}. None of the physical interventions were judged clinically significant^{56, 60, 75}.

Other intervention studies with data available for synthesis involved education/information^{63, 83} and aromatherapy^{52, 83} interventions.

Education/ information: three studies evaluated the effects of education/information interventions^{43, 63, 87}. After the clinical procedure one studies reported statistical significance favouring the intervention ($p < 0.05$)⁶³. The mean difference in anxiety did not exceed the MID in two studies^{43, 63}, and small to large effects sizes favouring the comparator were observed. One study investigating a multi-media information and instruction intervention deemed to be clinically significant⁸⁷.

Aromatherapy: two studies evaluated the effects of aromatherapy essential oil interventions with different methods of diffusion^{52, 83}. One study reported statistical significance favouring the intervention ($p < 0.05$)⁵² and the other did not⁸³. The difference in mean anxiety exceeded the MID in both studies^{52, 83}. Medium to large effect sizes were observed in both studies and were deemed clinically significant^{52, 83}. These two studies investigating Lavandula angustifolia, Citrus aurantium L, Lavender-sandalwood, and Orange-peppermint aromatherapy were deemed clinically significant^{52, 83}.

DISCUSSION

The aim of this review was to identify effective comfort interventions to support patients undergoing clinical procedures that require a patient to sustain the same position over a

period greater than 10 minutes. Thirteen comfort interventions were identified which ranged from aromatherapy to virtual reality delivered before and during nineteen different clinical procedures in 46 studies. Anxiety outcomes were synthesised as the outcome measures were validated and reported before and after clinical procedure in 26 studies.

The findings of the review showed that many comfort interventions produced statistically significant improvement in anxiety outcomes but did not demonstrate clinical significance as defined for this study. Aromatherapy^{52, 59, 83} used in colonoscopy, interventional radiology and minor surgery demonstrated both statistical and clinical significance and could be used in radiotherapy with careful consideration of application. Aromatherapy using vaporising systems may be contraindicated because of the potential for skin irritation or allergies linked to radiation induced skin toxicity or for vapour damage to radiotherapy equipment. A clothing tab infused with aromatherapy oils, found to be favourable in previous clinical trials⁹², may be more appropriate in radiotherapy. Audio and audio-visual interventions demonstrated medium to large effect sizes^{44,47,48,51,53,55,60,67,68,71,73,77,82,88} with several showing clinical significance that warrant further investigation in radiotherapy. A number of radiotherapy departments have audio-visual technology available to support their patients and audio interventions have been successfully tested in radiotherapy. For example, Chen et al⁹³ reported that music therapy reduced pre-radiotherapy anxiety only but did not focus on the effect during the clinical procedure and for this reason, was not included in this review. Audio interventions may be contraindicated in radiotherapy at times where constant communication between radiographers and patients is required such as verbal instructions to patients on performing deep inspiration breath hold or where an audio device such as earphones or audio pillows attenuate the radiation beam. Devices may be impractical due to an immobilisation mask. Visual interventions may not be so easily accommodated during some radiotherapy techniques but some interventions such as decorative wall colour or murals may be a pragmatic option.

Three psychological interventions and two physical interventions provided immediately before or during the clinical procedure demonstrated medium to large effect sizes^{57,60,81}. Psychological interventions provided as part of the preparation for radiotherapy have been studied and cognitive behavioural therapy and hypnosis have been shown to significantly ($p = .0035$) improve breast cancer patient general experiences⁹⁴, and likewise was not included in this review. Similarly, to improve mood, massage was provided during a course of radiotherapy treatment reduced anger, anxiety and depression in patients with breast cancer receiving radiotherapy ($P < 0.001$)⁹⁵. This review focused on interventions that could be delivered within radiotherapy sessions. Psychological interventions could be readily adopted if self-administered using an audio player. Use of empathetic interventions encouraging social interaction could be challenging to deliver. However Gibbon et al⁹⁶ found that patient

orientated communications skills training for the radiotherapy multi-disciplinary team resulted in significantly more empathetic interaction ($p = 0.037$).

Distraction using physical devices such as stress balls could be implemented with care taken not to disrupt the desired position for accurate radiotherapy. One intervention providing educational information via DVD demonstrated clinical significance⁸⁷ and could be implemented in a radiotherapy department. These interventions could also be applicable to clinical procedures including brachytherapy where there is need to develop non-pharmacological interventions⁹⁷ and paediatric radiotherapy where general anaesthesia could be reduced⁹⁸.

One gap observed from the studies is the effect of combining interventions as a 'comfort package' to enhance effectiveness. Simmons et al⁸⁰ investigated four interventions to support patients undergo cataract surgery with favourable results for combined interventions. Similarly, a systematic review by Bice et al¹⁹ found statistically significant differences favouring multifaceted (more than one intervention) interventions in most studies included in their review. Further research investigating a comfort intervention package (multiple interventions) may provide greater effectiveness for patients during radiotherapy treatment. Some methodological aspects of the systematic literature review and reviewed studies warrant further consideration. Firstly, anxiety outcome measures may not be the most suitable measure of comfort. The current review included studies with interventions that aimed to comfort, or to alleviate or reduce discomfort, anxiety and distress of clinical procedures. Comfort can be viewed holistically within physical, sociocultural, psychospiritual and environmental contexts that are not reflected in anxiety measures. There are limited comfort outcome measures, however the recently validated Radiotherapy-Experience-Questionnaire could be considered for measuring comfort in radiotherapy⁹⁹. Going forward, use of comfort outcome measures within all specialties is required for generating new evidence and confirming treatment effects of comfort interventions.

For the purposes of this review, clinical significance of the anxiety measures was demonstrated with a medium or above effect size (≥ 0.4) and mean differences greater than the MID. However, the availability of information about MID specific to the outcome measures reported in this review was limited. The MID level of 10 for the STAI was based on a population of smokers; in a non-smoking population the MID maybe higher or lower⁴⁰. Similarly, the MID for the BAI and HAS was based on a sample of patients with Parkinson's⁴². Further work is required for MID development in appropriate populations to assist with determining clinically effective interventions.

The research quality of the reviewed studies was an issue and a meta-analysis was not conducted due to this factor and because of the challenges of defining the nuances of comfort, clinical procedures and interventions. 8 RCTs were deemed unacceptable due to a

high risk of selection bias and were not included in the data synthesis. Many studies did not register with an international clinical trial register which affected the assessment of selective reporting; these studies were therefore judged as having unclear RoB. Although there were some methodological challenges, a rigorous review process was followed and a semi-automated machine learning programme, RoBotReviewerTM²⁸⁻²⁹, was used for Cochrane RoB to increase the rigour of this review by reducing the impact of human factors during data extraction. Combining the use of semi-automated extraction with manual assessment was useful and future reviews should consider using machine or deep learning systems to improve the rigour and quality of data extraction¹⁰⁰.

To our knowledge, this is the first systematic review that could support the further investigation of comfort interventions in radiotherapy. Given the limited recommendation of how to manage patient comfort during radiotherapy from national and European guidelines^{2,9}, the findings of this review and further investigation of comfort interventions will provide the evidence required for future guidelines. Given the perpetual increase in new effective treatment options and technology available in radiotherapy, it is essential that the community embraces and implements comfort interventions ensuring the best outcomes for patients.

CONCLUSION

The majority of aromatherapy interventions were clinically significant; and they can be potentially considered for radiotherapy that require patients to sustain and endure the same position over time similar to these clinical procedures. There was limited evidence for other comfort interventions, although most effect sizes favoured the intervention, suggesting important benefit to patients. Further investigation of these comfort interventions is warranted, including tailoring interventions to patient choice and determining if multiple interventions could be used concurrently to improve their effectiveness. This is crucial for complex radiotherapy that necessitates more demand and attention to patient comfort to ensure stability for targeted treatment.

REFERENCES

- 1) Folkert MR & Timmerman RD. Stereotactic ablative body radiosurgery (SABR) or Stereotactic body radiation therapy (SBRT). *Adv Drug Deliv Rev* 2017 Jan; 15:109:3-14.
- 2) The Royal College of Radiologists (RCR), Society and College of Radiographers, Institute of Physics and Engineering in Medicine. 2008. On target: ensuring geometric accuracy in radiotherapy. London: RCR. P. 11–14
- 3) Boda-Heggemann J, Mai S, Fleckenstein J, Siebenlist K, Simeonova A, Ehmann M, et al. Flattening-filter-free intensity modulated breath-hold image-guided SABR (Stereotactic ABlative Radiotherapy) can be applied in a 15-min treatment slot. *Radiother Oncol.* 2013 Dec;109(3):505-9
- 4) Bartlett FR, Colgan RM, Donovan EM, McNair HA, Carr K, Evans PM, Et al. The UK HeartSpare Study (Stage IB): randomised comparison of a voluntary breath-hold technique and prone radiotherapy after breast conserving surgery. *Radiother Oncol.* 2015 Jan;114(1):66-72
- 5) Goldsworthy S, Mundy K, Latour JM. A focus group consultation round exploring patient experiences of comfort during radiotherapy for head and neck cancer; *Journal of J Radiother Pract.* 2016 Jan; 15 (2):143-149
- 6) Bayley AJ, Catton CN, Haycocks T, Kelly V, Alasti H, Bristow R, et al . A randomized trial of supine vs. prone positioning in patients undergoing escalated dose conformal radiotherapy for prostate cancer. *Radiotherapy and Oncology.* 2004 Jan; 70 (1):37-44
- 7) Dawson LA, & Balter JM. Interventions to Reduce Organ Motion Effects in Radiation Delivery; *Seminars in Radiation Oncology.* 2004 Jan 1;14 (1): 76-80
- 8) Osztavics A, & Kirchheiner K. Immobilising the patient to be as comfortable as possible. A general overview. *RA. Radiother Oncol.* 2017 May;123. S54
- 9) Kolcaba KY. Holistic comfort: operationalizing the construct as a nurse-sensitive outcome. *ANS Adv Nurs Sci.* 1992 Sep;15(1):1-10
- 10) Kolcaba KY. A theory of holistic comfort for nursing. *J Adv Nurs.* 1994 Jun;19(6):1178-84
- 11) Kolcaba KY, & Fox C. The effects of guided imaginary on comfort of women with early stage breast cancer undergoing radiation therapy. *Oncology Nursing Forum.* 1991; 26(1): 67–92
- 12) Nixon JL, Cartmill B, Turner J, Pigott AE, Brown E, Wall LR, et al. Exploring the prevalence and experience of mask anxiety for the person with head and neck cancer undergoing radiotherapy. *J Med Radiat Sci.* 2018 Dec;65(4):282-290

- 13) Leech M, Mary Coffey M, Mast M, Moura F, Osztavics A, Pasinie D, et al. ESTRO ACROP guidelines for positioning, immobilisation and position verification of head and neck patients for radiation therapists. *Technical Innovations & Patient Support in Radiation Oncology*. 2017 March; 1: 1-7
- 14) Malinowski A, & Stamler LL. Comfort: exploration of the concept in nursing. *J Adv Nurs*. 2002 Sep;39(6):599-606.
- 15) Nixon JL, Brown B, Pigott AE, Turner J, Brown E, Bernard A, et al. A prospective examination of mask anxiety during radiotherapy for head and neck cancer and patient perceptions of management strategies. *J Med Radiat Sci*. 2019 Sep;66(3):184-190
- 16) Bice AA, & Wyatt TH. Holistic Comfort Interventions for Paediatric Nursing Procedures: A Systematic Review. *J Holist Nurs*. 2017 Sep;35(3):280-295
- 17) Hudson BF, & Ogden J. Exploring the Impact of Intraoperative Interventions for Pain and Anxiety Management During Local Anesthetic Surgery-A Systematic Review and Meta-Analysis. *J Perianesth Nurs*. 2016 Apr;31(2):118-33
- 18) Rudin D, Kiss A, Wetz RV, Sottile VM. Music in the endoscopy suite: a meta-analysis of randomized controlled studies. *Endoscopy*. 2007 Jun;39(6):507-10.
- 19) Dahele M, Verbakel W, Cuijpers J, Slotman B, Senan S, An analysis of patient positioning during stereotactic lung radiotherapy performed without rigid external immobilization. *Radiother Oncol*. 2012 Jul;104(1):28-32
- 20) Dogramadzi S, Griffiths G, Cross P, Goldsworthy S, Winstone B. Motion Capture Pillow for Head and- Neck Cancer Radiotherapy Treatment. 7th IEEE RAS/EMBS International Conference on Biomedical Robotics and Biomechanics (BioRoB 2018), 26-29 August 2018, Enschede, Netherlands.
- 21) Centre for Reviews and Dissemination (CRD). *Systematic reviews: CRD's guidance for undertaking reviews in health care*. York Publishing Services Ltd. 2009 Jan. 1-294. Department of Health. *Cancer Reform Strategy*; 2007
- 22) Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Ioannidis JP, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *J Clin Epidemiol*. 2009 Oct;62(10): e1-34
- 23) Hoffmann TC, Glasziou PP, Boutron I, Milne R, Perera R, Moher D, et al. Better Reporting of Interventions: Template for Intervention Description and Replication (TIDieR) Checklist and Guide. *Gesundheitswesen*. 2016 Mar;78(3): e174
- 24) Huang X, Lin J, Demner-Fushman D. Evaluation of PICO as a knowledge representation for clinical questions. *AMIA Annu Symp Proc*. 2006:359-63.

- 25) Higgins JPT, Green S, Cochrane Handbook for Systematic Reviews of Interventions. Version 5.1.0 Edition. London, The Cochrane Collaboration, 2011.
- 26) Mathes T, Klößen P, Pieper D. Frequency of data extraction errors and methods to increase data extraction quality: a methodological review. *BMC Med Res Methodol*. 2017 Nov 28;17(1):152
- 27) Armijo-Olivo S, Ospina M, da Costa BR, Egger M, Saltaji H, Fuentes J, et al. Poor reliability between Cochrane reviewers and blinded external reviewers when applying the Cochrane risk of bias tool in physical therapy trials. *PLoS One*. 2014 May 13;9(5):e96920
- 28) Marshall IJ, Kuiper J, Wallace BC. RoBotReviewer: evaluation of a system for automatically assessing bias in clinical trials. *J Am Med Inform Assoc*. 2016 Jan;23(1):193-201
- 29) Marshall IJ, Noel-Storr A, Kuiper J. Machine learning for identifying Randomized Controlled Trials: An evaluation and practitioner's guide. *Res Synth Methods*. 2018 Dec;9(4):602-61
- 30) Bothe AK, & Richardson JD. Statistical, practical, clinical, and personal significance: definitions and applications in speech-language pathology. *Am J Speech Lang Pathol*. 2011 Aug;20(3):233-42
- 31) Wright A, Hannon J, Hegedus EJ, Kavchak AE. Clinimetrics corner: a closer look at the minimal clinically important difference (MCID). *J Man Manip Ther*. 2012 Aug;20(3):160-6
- 32) Jakobsen JC, Gluud C, Winkel P, Lange T, Wetterslev J. The thresholds for statistical and clinical significance - a five-step procedure for evaluation of intervention effects in randomised clinical trials. *BMC Med Res Methodol*. 2014 Mar 4; 14:34
- 33) Grissom RJ, & Kim JJ. *Effect Sizes for Research: Univariate and Multivariate Applications*, Second Edition. New York, Routledge, Taylor & Francis Group., 2011
- 34) Leech NL, & Onwuegbuzie AJ. A Call for Greater Use of Nonparametric Statistics. Annual Meeting of the Mid-South Educational Research Association. 2002 Nov: 1-25
- 35) Wan X, Wang W, Liu J, Tong T. Estimating the sample mean and standard deviation from the sample size, median, range and/or interquartile range. *BMC Med Res Methodol*. 2014 Dec 19; 14:135
- 36) Luo D, Wan X, Liu J, Tong T. Optimally estimating the sample mean from the sample size, median, mid-range, and/or mid-quartile range. *Stat Methods Med Res*. 2018 Jun;27(6):1785-1805
- 37) Altman DG, & Bland JM. How to obtain the P value from a confidence interval. *BMJ*. 2011;343: d2304.

- 38) Rice ME, & Harris GT. Comparing effect sizes in follow-up studies: ROC Area, Cohen's d, and r. *Law Hum Behav.* 2005 Oct; 29(5):615-20.
- 39) Corsaletti BF, Proença MDGL, Bisca GKW, Leite JC, Bellinetti LM, Pitta F. Minimal important difference for anxiety and depression surveys after intervention to increase daily physical activity in smokers. *Fisioter Pesq.* 2014;21(4):359-364
- 40) Taghizadeh N, Tremblay A, Cressman S, Peacock S, McWilliams AM, MacEachern P, et al. Health-related quality of life and anxiety in the PAN-CAN lung cancer screening cohort. *BMJ Open.* 2019 Jan 17;9(1): e024719
- 41) Facco E, Stellini E, Bacci C, Manani G, Pavan C, Cavallin F, et al. Validation of visual analogue scale for anxiety (VAS_A) in preanesthesia evaluation. *Minerva Anestesiologica.* 2013; 79 (12): 1389-1395
- 42) Leentjens AF, Dujardin K, Marsh L, Richard IH, Starkstein SE, Martinez-Martin P. Anxiety rating scales in Parkinson's disease: a validation study of the Hamilton anxiety rating scale, the Beck anxiety inventory, and the hospital anxiety and depression scale. *Mov Disord.* 2011 Feb 15;26(3):407-15
- 43) Ahlander BM, Engvall J, Maret E, Ericsson E. Positive effect on patient experience of video information given prior to cardiovascular magnetic resonance imaging: A clinical trial. *J Clin Nurs.* 2018; 27:1250–1261.
- 44) Angioli R, De Cicco Nardone C, Plotti F, Cafà EV, Dugo N, Damiani P, et al. Use of music to reduce anxiety during office hysteroscopy: prospective randomized trial. *J Minim Invasive Gynecol.* 2014 May-Jun;21(3):454-9
- 45) Argstatter H, Haberbosch W, Bolay HV. Study of the effectiveness of musical stimulation during intracardiac catheterization. *Clin Res Cardiol.* 2006 Oct;95(10):514-22
- 46) Björkman I, Karlsson F, Lundberg A, Frisman GH. Gender differences when using sedative music during colonoscopy. *Gastroenterol Nurs.* 2013 Jan-Feb;36(1):14-20
- 47) Buffum MD, Sasso C, Sands LP, Lanier E, Yellen M, Hayes A. A music intervention to reduce anxiety before vascular angiography procedures. *J Vasc Nurs.* 2006 Sep;24(3):68-73
- 48) Chlan L, Evans D, Greenleaf M, Walker J. Effects of a single music therapy intervention on anxiety, discomfort, satisfaction, and compliance with screening guidelines in outpatients undergoing flexible sigmoidoscopy. *Gastroenterol Nurs.* 2000 Jul-Aug;23(4):148-56
- 49) Choi SM, Lee J, Park YS, Lee CH, Lee SM, Yim JJ, Effect of Verbal Empathy and Touch on Anxiety Relief in Patients Undergoing Flexible Bronchoscopy: Can Empathy Reduce Patients' Anxiety? *Respiration.* 2016;92(6):380-388

- 50) Diette GB, Lechtzin N, Haponik E, Devrotes A, Rubin HR. Distraction therapy with nature sights and sounds reduces pain during flexible bronchoscopy: a complementary approach to routine analgesia. *Chest*. 2003 Mar;123(3):941-8
- 51) Drahota A, Galloway E, Stores R, Ward D, Severs M, Dean T. Audiovisual distraction as an adjunct to pain and anxiety relief during minor surgery. *Foot (Edinb)*. 2008 Dec;18(4):211-9
- 52) Eslami J, Ebrahimi A, Hosseinkhani A, Khazaei Z, Darvishi I. The effect of aromatherapy using Lavender (*Lavandula angustifolia* Miller) and Citrus aurantium L. extracts to treat anxiety of patients undergoing laparoscopic cholecystectomy: A randomized clinical trial in Iran *Biomed Res Ther* 2018; 5(3): 2096-2110
- 53) Fang AS, Movva L, Ahmed S, Waldman D, Xue J. Clinical Efficacy, Safety, and Feasibility of Using Video Glasses during Interventional Radiologic Procedures: A Randomized Trial. *J Vasc Interv Radiol*. 2016 Feb;27(2):260-7
- 54) Frank LS, Frank JL, March D, Makari-Judson G, Barham RB, Mertens WC. Does therapeutic touch ease the discomfort or distress of patients undergoing stereotactic core breast biopsy? A randomized clinical trial. *Pain Med*. 2007 Jul-Aug;8(5):419-24
- 55) Hayes A, Buffum M, Lanier E, Rodahl E, Sasso C. A music intervention to reduce anxiety prior to gastrointestinal procedures. *Gastroenterol Nurs*. 2003 Jul-Aug;26(4):145-9.
- 56) Heidaria F, Rejeha N, Heravi-Karimooia M, Tadrissi SD, Vaismoradic M. Effect of short-term hand reflexology on anxiety in patients before coronary angiography: A randomized placebo-controlled trial. *European Journal of Integrative Medicine*; 2017;16: 1–7
- 57) Hızlı F, Özcan O, Selvi İ, Eraslan P, Köşüş A, Baş O, et al. The effects of hypnotherapy during transrectal ultrasound-guided prostate needle biopsy for pain and anxiety. *Int Urol Nephrol*. 2015 Nov;47(11):1773-7
- 58) Hozumi H, Hasegawa S, Tsunenari T, Sanpei N, Arashina Y, Takahashi K, et al. Aromatherapies using *Osmanthus fragrans* oil and grapefruit oil are effective complementary treatments for anxious patients undergoing colonoscopy: A randomized controlled study. *Complement Ther Med*. 2017 Oct; 34:165-169
- 59) Hu PH, Peng YC, Lin YT, Chang CS, Ou MC. Aromatherapy for reducing colonoscopy related procedural anxiety and physiological parameters: a randomized controlled study. *Hepatogastroenterology*. 2010 Sep-Oct;57(102-103):1082-6
- 60) Hudson BF, Ogden J, Whiteley MS. Randomized controlled trial to compare the effect of simple distraction interventions on pain and anxiety experienced during conscious surgery. *Eur J Pain*. 2015 Nov;19(10):1447-55

- 61) Jiménez-Jiménez M, García-Escalona A, Martín-López A, De Vera-Vera R, De Haro J. Intraoperative stress and anxiety reduction with music therapy: a controlled randomized clinical trial of efficacy and safety. *J Vasc Nurs.* 2013 Sep;31(3):101-6
- 62) Kekecs Z, Jakubovits E, Varga K, Gombos K. Effects of patient education and therapeutic suggestions on cataract surgery patients: a randomized controlled clinical trial. *Patient Educ Couns.* 2014 Jan;94(1):116-22
- 63) Kola S, Walsh JC, Hughes BM, Howard S. Matching intra-procedural information with coping style reduces psychophysiological arousal in women undergoing colposcopy. *J Behav Med.* 2013 Aug;36(4):401-12
- 64) Kwekkeboom KL. Music versus distraction for procedural pain and anxiety in patients with cancer. *Oncol Nurs Forum.* 2003 May-Jun;30(3):433-40.
- 65) Lang EV, Benotsch EG, Fick LJ, Lutgendorf S, Berbaum ML, Berbaum KS, et al. Adjunctive non-pharmacological analgesia for invasive medical procedures: a randomised trial. *Lancet.* 2000 Apr 29;355(9214):1486-90
- 66) Lee WL, Sung HC, Liu SH, Chang SM. Meditative music listening to reduce state anxiety in patients during the uptake phase before positron emission tomography (PET) scans. *Br J Radiol.* 2017 Feb;90(1070):20160466
- 67) McSherry T, Atterbury M, Gartner S, Helmold E, Searles DM, Schulman C. Randomized, Crossover Study of Immersive Virtual Reality to Decrease Opioid Use During Painful Wound Care Procedures in Adults. *J Burn Care Res.* 2018 Feb 20;39(2):278-285
- 68) Ng MY, Karimzad Y, Menezes RJ, Wintersperger BJ, Li Q, Forero J, et al. Randomized controlled trial of relaxation music to reduce heart rate in patients undergoing cardiac CT. *Eur Radiol.* 2016 Oct;26(10):3635-42
- 69) Navidian A, Moulaei N, Ebrahimi Tabas E, Solaymani S. The effect of audiovisual distraction on the tolerability of flexible bronchoscopy: a randomized trial. *Clin Respir J.* 2018 Jan;12(1):76-83
- 70) Nilsson U1, Lindell L, Eriksson A, Kellerth T. The effect of music intervention in relation to gender during coronary angiographic procedures: a randomized clinical trial. *Eur J Cardiovasc Nurs.* 2009 Sep;8(3):200-6
- 71) Nilsson U. Effectiveness of music interventions for women with high anxiety during coronary angiographic procedures: a randomized controlled. *Eur J Cardiovasc Nurs.* 2012 Jun;11(2):150-3
- 72) Packiam VT, Nottingham CU, Cohen AJ, Eggener SE, Gerber GS. No Effect of Music on Anxiety and Pain During Transrectal Prostate Biopsies: A Randomized Trial. *Urology.* 2018 Jul; 117:31-35

- 73) Padam A, Sharma N, Sastri OSKS, Mahajan S, Sharma R, Sharma D. Effect of listening to Vedic chants and Indian classical instrumental music on patients undergoing upper gastrointestinal endoscopy: A randomized control trial. *Indian J Psychiatry*. 2017 Apr-Jun;59(2):214-218
- 74) Ripley L, Christopoulos G, Michael TT, Alomar M, Rangan BV, Roesle M, et al. Randomized controlled trial on the impact of music therapy during cardiac catheterization on reactive hyperemia index and patient satisfaction: the Functional Change in Endothelium After Cardiac Catheterization, With and Without Music Therapy (FEAT) study. *J Invasive Cardiol*. 2014 Sep;26(9):437-42.
- 75) Rosen J, Lawrence R, Bouchard M, Doros G, Gardiner P, Saper R, et al. Massage for perioperative pain and anxiety in placement of vascular access devices. *Adv Mind Body Med*. 2013 Winter;27(1):12-23.
- 76) Schupp CJ, Berbaum K, Berbaum M, Lang EV. Pain and anxiety during interventional radiologic procedures: effect of patients' state anxiety at baseline and modulation by nonpharmacologic analgesia adjuncts. *J Vasc Interv Radiol*. 2005 Dec;16(12):1585-92.
- 77) Shabanloei R, Golchin M, Esfahani A, Dolatkah R, Rasoulia M. Effects of music therapy on pain and anxiety in patients undergoing bone marrow biopsy and aspiration. *AORN J*. 2010 Jun;91(6):746-51.
- 78) Shahsavaria H, Abadb MEE, Yekaninejad MS. The effects of foot reflexology on anxiety and physiological parameters among candidates for bronchoscopy: A randomized controlled trial. *Eur J Integr Med*. 2017 June; 12:177–181.
- 79) Shenefelt PD. Anxiety reduction using hypnotic induction and self-guided imagery for relaxation during dermatologic procedures. *Int J Clin Exp Hypn*. 2013;61(3):305-18.
- 80) Simmons D, Chabal C, Griffith J, Rausch M, Steele B. A clinical trial of distraction techniques for pain and anxiety control during cataract surgery. *Insight*. 2004 Oct-Dec;29(4):13-6.
- 81) Snow A, Dorfman D, Warbet R, Cammarata M, Eisenman S, Zilberfein F et al. A randomized trial of hypnosis for relief of pain and anxiety in adult cancer patients undergoing bone marrow procedures. *J Psychosoc Oncol*. 2012;30(3):281-93.
- 82) Sobana R, Sundar S, Dixit P. A study on the effects of music therapy in patients posted for upper gastrointestinal endoscopy. *Int J Pharm Bio Sci*. 2015 April; 6(2):307–313.
- 83) Trambert R, Kowalski MO, Wu B, Mehta N, Friedman P. A Randomized Controlled Trial Provides Evidence to Support Aromatherapy to Minimize Anxiety in Women Undergoing Breast Biopsy. *Worldviews Evid Based Nurs*. 2017 Oct;14(5):394-402.

- 84) Uğraş GA, Yıldırım G, Yüksel S, Öztürkçü Y, Kuzdere M, Öztekin SD. The effect of different types of music on patients' preoperative anxiety: A randomized controlled trial. *Complement Ther Clin Pract*. 2018 May; 31:158-163.
- 85) Walker MR, Kallingal GJ, Musser JE, Folen R, Stetz MC, Clark J. Treatment efficacy of virtual reality distraction in the reduction of pain and anxiety during cystoscopy. *Mil Med*. 2014 Aug;179(8):891-6.
- 86) Weeks BP & Nilsson U. Music interventions in patients during coronary angiographic procedures: a randomized controlled study of the effect on patients' anxiety and well-being. *Eur J Cardiovasc Nurs*. 2011 Jun;10(2):88-93.
- 87) Wu KL, Chen SR, Ko WC, Kuo SY, Chen PL, Su HF, et al. The effectiveness of an accessibility-enhanced multimedia informational educational programme in reducing anxiety and increasing satisfaction of patients undergoing cardiac catheterisation. *J Clin Nurs*. 2014 Jul;23(13-14):2063-73.
- 88) Xiaolian J, Xiaolin L, Lan ZH. Effects of visual and audiovisual distraction on pain and anxiety among patients undergoing colonoscopy. *Gastroenterol Nurs*. 2015 Jan-Feb;38(1):55-61.
- 89) Spielberger CD. *Manual for the State-Trait-Anxiety Inventory: STAI (form Y)*. Palo Alto, CA: Consulting Psychologists Press. 1983
- 90) Marteau TM, & Bekker H. The development of a six-item short-form of the state scale of the Spielberger State-Trait Anxiety Inventory (STAI). *Br J Clin Psychol*. 1992 Sep;31 (Pt 3):301-6.
- 91) Ulusoy M, Sahin NH, Erkmen H. Turkish version of the Beck Anxiety Inventory: psychometric properties. *J Cogn Psychother*. 1998 12:163–172.
- 92) Atwal R, Hayes J, Nanalal C. Lavender Aromatherapy to Reduce Anxiety Levels in Brain, Head and Neck and Breast Patients during Their CT Simulation Appointment for Radiation Therapy. *Journal of Medical Imaging and Radiation Sciences*. 2016 March; 47 (1) S25
- 93) Chen LC, Wang TF, Shih YN, Wu LJ. Fifteen-minute music intervention reduces pre-radiotherapy anxiety in oncology patients. *Eur J Oncol Nurs*. 2013 Aug;17(4):436-41.
- 94) Schnur JB, David D, Kangas M, Green S, Bovbjerg DH, Montgomery GH. A randomized trial of a cognitive-behavioral therapy and hypnosis intervention on positive and negative affect during breast cancer radiotherapy. *J Clin Psychol*. 2009 Apr;65(4):443-55
- 95) Darabpour S, Kheirkhah M, Ghasemi E. Effects of Swedish Massage on the Improvement of Mood Disorders in Women with Breast Cancer undergoing Radiotherapy. *Iran Red Crescent Med J*. 2016 Aug 10;18(11):e25461

- 96) Gibon AS, Merckaert I, Liénard A, Libert Y, Delvaux N, Marchal S, et al. Is it possible to improve radiotherapy team members' communication skills? A randomized study assessing the efficacy of a 38-h communication skills training program. *Radiother Oncol.* 2013 Oct;109(1):170-7.
- 97) Humphrey P, Bennett C, Cramp F. The experiences of women receiving brachytherapy for cervical cancer: A systematic literature review. *Radiography (Lond).* 2018 Nov;24(4):396-403
- 98) O'Callaghan C, Sexton M, Wheeler G. Music therapy as a non-pharmacological anxiolytic for paediatric radiotherapy patients. *Australas Radiol.* 2007 Apr;51(2):159-62
- 99) Olausson K, Hansson AH, Zackrisson B, Edvardsson D, Östlund U, Nyholm T. Development and psychometric testing of an instrument to measure the patient's experience of external radiotherapy: The Radiotherapy Experience Questionnaire (RTEQ). *Technical Innovations & Patient Support in Radiation Oncology.* 2017 July;3-4: 7-12
- 100) Goswami S, Pal S, Goldsworthy S, Basu T. An Effective Machine Learning Framework for Data Elements Extraction from the Literature of Anxiety Outcome Measures to Build Systematic Review. In: Abramowicz W, Corchuelo R. (eds) *Business Information Systems. BIS 2019. Lecture Notes in Business Information Processing*, vol 353. Springer, Cham




Figure 1. Flow diagram of the strategy search.




Figure. 2. Cochrane risk of bias summary of randomised controlled trials (n=46)

Table 1 – Intervention delivery characteristics

Comfort intervention	Rationale	Materials	Delivery features	Delivered by
AUDIO-VISUAL TECHNOLOGY INTERVENTIONS				
Audio →	<ul style="list-style-type: none"> ♦ Reduce anxiety^{43-48,55, 60, 64, 66, 71-73, 77, 84, 86, 88}, discomfort⁴⁸, stress⁶¹, pain^{42, 60, 64, 71-72, 77}, heart rate⁶⁸, analgesics/anxiolytics⁷¹ ♦ Improve satisfaction^{48, 60, 74}, compliance^{48, 82, 92}, relaxation^{42, 70-71}, comfort⁷⁰⁻⁷¹, wellbeing⁵⁹, reactive hyperaemia index⁷⁴ 	<ul style="list-style-type: none"> ♦ A range of music genres.^{42, 44-48, 55, 60-61, 64, 68, 70-73, 77, 82} ♦ Vedic chants⁷³ ♦ Nature sounds ♦ Music therapy (meditative, relaxing)^{66, 68, 74, 86} 	<ul style="list-style-type: none"> ♦ Music player (e.g. CD player/computer)^{44-48, 60-61, 64, 68, 70-73, 77, 82, 86,} ♦ Loudspeaker^{70-72, 82} ♦ Earphones^{45-48, 60-61, 64, 68, 73, 77} ♦ Cushion with speaker^{74, 86} ♦ Not specified^{42,55, 68} ♦ Most at 50-80bpm^{42, 44-48, 55, 60-61, 64, 68, 70-73, 77, 82} 	<ul style="list-style-type: none"> ♦ Technician⁴² ♦ Music therapist⁴⁵ ♦ Research nurse⁴⁴ ♦ Student nurses⁴⁶ ♦ Nurses^{46-47,55,60, 70} ♦ Investigators^{48, 55} ♦ Physicians^{42, 46} ♦ CT technologists⁶⁸ ♦ Study personnel⁸⁶ ♦ Not specified^{50, 61, 64, 66, 68, 72-74, 77, 82, 84, 88}
Audio-visual →	<ul style="list-style-type: none"> ♦ Reduce anxiety^{50-53, 60} ♦ Reduce pain^{50-52,60} ♦ Improve experience⁵²⁻⁵³ ♦ Improve satisfaction⁵² ♦ Tolerate procedure⁶⁹ 	<ul style="list-style-type: none"> ♦ Nature sounds such as a waterfall^{50-51, 88} ♦ Nature scene such as a mountain stream, tropical beach, general landscape scenery and animation^{50-51,69, 88} ♦ Videos included documentaries and movies^{60, 88} ♦ Comedies, documentaries and panel-based quiz shows ♦ Iranian music⁶⁹ 	<ul style="list-style-type: none"> ♦ Music player (e.g. CD player/computer)^{50-51, 69} ♦ loudspeaker ♦ Earphones^{50-52, 69} ♦ Wall or ceiling mounted murals of nature scenes with/without lighting^{50-51,53,69, 80}. ♦ Video goggles connected to DVD⁵²⁻⁵³ ♦ Wall mounted monitor connected to DVD ♦ Projector connected to DVD⁶⁰ 	<ul style="list-style-type: none"> ♦ Standard clinic staff⁵¹ ♦ Nurse^{52,60} ♦ Not specified^{50, 52-53, 69}
Visual →	<ul style="list-style-type: none"> ♦ Distraction from pain, anxiety, and tolerate procedure⁸⁸ 	<ul style="list-style-type: none"> ♦ Scenery, animation, to film⁸⁸ 	<ul style="list-style-type: none"> ♦ Monitor with DVD player (no sound)⁸⁸ 	<ul style="list-style-type: none"> ♦ Not specified⁸⁸
Virtual reality (VR) →	<ul style="list-style-type: none"> ♦ Reduce pain⁸⁵ ♦ Reduce anxiety⁸⁵ ♦ Reduce opioid use⁶⁷ 	<ul style="list-style-type: none"> ♦ Throwing/shooting snowballs at objects by clicking a computer mouse button^{67, 85} 	<ul style="list-style-type: none"> ♦ Headset goggles, earphones, DVD player, VR system⁶⁷ ♦ VR group donned a VR helmet and track ball hand controller⁸⁵ 	<ul style="list-style-type: none"> ♦ Nurses⁶⁷ ♦ Not specified⁸⁵

Comfort interventions	Rationale	Materials	Delivery features	Delivered by
PSYCHOLOGICAL INTERVENTIONS				
Breathing techniques →	♦Reduce discomfort, pain & anxiety ⁸⁰	♦Verbal coaching and slow breathing instructed ⁸⁰	♦Face to face ⁸⁰	♦Nurses ⁸⁰
Cognitive behavioural therapy →	♦Improve relaxation ⁷⁹	♦ Live guided imagery ⁷⁹ ♦ Recorded guided imagery ⁷⁹	♦ Face to face ⁷⁹ ♦ CD player ⁷⁹	♦Trained therapist ⁷⁹
Distraction →	♦Reduce pain & anxiety ⁶⁴	♦ Participant reads a book ⁶⁴	♦ A book ⁶⁴	♦Research nurses ⁶⁴
Empathic attention →	♦Reduce pain ^{60, 76} ♦Reduce anxiety ^{49, 60, 76} ♦Improve satisfaction ⁶⁰ ♦Reduce discomfort ⁶⁵ ♦Reduce adverse effects ⁶⁵	♦ Verbal empathy ^{49,65,76} & touch ⁴⁹ ♦ Non-verbal attention ⁷⁶ ♦Engage in conversation ^{60, 76} ♦Attentive listening, Perception of control, Emotionally neutral Avoid negative suggestion ⁷⁶	♦ Face to face ^{49, 60, 65, 76}	♦Nurse ⁶⁰ ♦Medical student ^{65, 76} ♦Psychology graduate ^{65, 76} ♦Therapist ⁴⁹
Hypnosis →	♦Reduce pain ^{76, 81} ♦Reduce anxiety ^{57, 76, 81} ♦Reduce discomfort & Reduce adverse effects ⁶⁵	♦ Progressive relaxation, visualisation, & deep trance ^{57,65, 81}	♦ Face to face ^{65, 81} ♦Self hypnosis ^{57,76}	♦Nurse ^{65, 76} ♦Medical student ^{65, 76} ♦Psychology graduate ^{65, 76} ♦Not specified ⁵⁷ ♦Social worker ⁸¹
	Rationale	Materials	Delivery features	Delivered by

Comfort interventions				
PHYSICAL INTERVENTIONS				
Comfort interventions	Rationale	Materials	Delivery features	Delivered by
Massage, therapeutic touch & reflexology →	<ul style="list-style-type: none"> ♦ Reduce pain^{75,80} ♦ Reduce anxiety^{56, 75, 78} ♦ Reduce distress⁵⁴ ♦ Reduce discomfort^{54,80} 	<ul style="list-style-type: none"> ♦ Massage^{75, 80} ♦ “energy repatterning” hand movements over parts of the patient’s anatomy (often the torso) where energy field abnormalities are detected⁵⁴ ♦ Three reflexology acupressure points for the pituitary gland, heart and solar plexus were stimulated by hand^{56,78} 	<ul style="list-style-type: none"> ♦ Face to face light finger⁸⁰ & 20 minutes Effleurage strokes across different parts of the body Massage⁷⁵ ♦ Face to face Kriegler and Kunz Therapeutic touch Massage⁵⁴ ♦ Face to face foot reflexology (both feet) for 10 minutes^{56,78} 	<ul style="list-style-type: none"> ♦ Nurse⁸⁰ ♦ Four trained practitioners⁵⁴ ♦ Massage therapist⁷⁵ ♦ Reflexologist^{56,78}
Distraction →	<ul style="list-style-type: none"> ♦ Reduce pain, anxiety and improve patient satisfaction⁶⁰ 	<ul style="list-style-type: none"> ♦ Stress balls⁶⁰ 	<ul style="list-style-type: none"> ♦ Stress balls manipulated during clinical procedure by participant⁶⁰ 	<ul style="list-style-type: none"> ♦ self-directed by patient⁶⁰

OTHER INTERVENTIONS				
Education/ information →	<ul style="list-style-type: none"> ♦ Improve experience⁴³ ♦ Reduce anxiety^{62, 87} ♦ Reduce psychophysiological arousal⁶³ ♦ Increase satisfaction⁸⁷ 	<ul style="list-style-type: none"> ♦ Participant watches live examination⁶³ ♦ Video education/ information⁴³ ♦ Audio information about procedure⁶² ♦ Instructional Accessibility-enhanced multimedia informational education (AEMIE)⁸⁷ 	<ul style="list-style-type: none"> ♦ Monitor screen of examination⁶³ ♦ Monitor screen with DVD player^{43, 87} ♦ Music player & headphones⁶² ♦ Head mounted display with headphones⁶³ 	<ul style="list-style-type: none"> ♦ Radiographer⁴³ ♦ Research assistant⁶² ♦ Nurse^{63, 87}
Aromatherapy →	<ul style="list-style-type: none"> ♦ Reduce anxiety^{52, 58-59, 83} ♦ Reduce physiology parameters^{59, 78} 	<ul style="list-style-type: none"> ♦ Lavandula angustifolia Miller, citrus aurantium L. essences⁵² ♦ Lavender oil, grapefruit oil, and Osmanthus fragrans+B7 oil for diffusion⁵⁸ ♦ Neroli essences were poured on gauze⁵⁹ ♦ Essential oils lavender/sandalwood on tab or orange/peppermint on tab⁸³ 	<ul style="list-style-type: none"> ♦ Participants inhaled aroma from the tissue paper for 20 minutes from a 20cm distance⁵² ♦ Diffuser used⁵⁸ ♦ Delivered via handheld-nebulizer with oxygen mask which pneumatically pump the oil into the mask; the oxygen masks were placed on the participants nose to smell for five minutes⁵⁹ ♦ Tabs placed on participant gown⁸³ 	<ul style="list-style-type: none"> ♦ Study researchers⁵² ♦ Endoscopist⁵⁸ ♦ Nurse⁸³ ♦ Not specified^{59,78}

Table 2 – Clinical significance of interventions before & after clinical procedures (based on anxiety outcome measures)

Source	Comfort intervention category	Outcome measure		Mean difference Before-after clinical procedure				Mean difference between groups	% difference between groups	Effect size with CI (95%)	Intervention Clinically significant
		Type	Minimal important difference (MID)	Intervention Group Mean difference	≥MID	Comparator Group Mean difference	≥MID				
Angioli R, et al 2014 ⁴⁴	Audio-visual technology interventions	STAI	10	3.4	X	1.1	X	2.2	66%	4.2 (3.8 to 4.5)	No
Buffum MD, et al 2006 ⁴⁷		STAI	10	3.4	X	1.1	X	2.2	66%	4.1 (3.5 to 4.6)	No
Chlan L, et al 2000 ⁴⁸		STAI	10	2.4	X	-1.6	X	4.0	167%	0.7 (0.2 to 1.2) *	No
Hayes A, et al 2003 ⁵⁵		STAI	10	4.4	X	1.5	X	2.9	66%	1.2 (0.9 to 1.5)	No
Hudson BF, et al 2015 ⁶⁰ (music)		STAI	10	0.0	X	-2.3	X	2.3	102%	1.3 (1.7 to 1.0)	No
Kwekkeboom KL, et al 2003 ⁶⁴		STAI	10	4.1	X	7.0	X	-2.9	-71%	-5.0 (-3.8 to -6.2)	No
LEE WL, et al 2017 ⁶⁶		STAI	10	5.3	X	-0.7	X	5.9	88%	5.6 (4.6 to 6.6)	No
Ng MY, Et al 2016 ⁶⁸		STAI	10	2.0	X	1.2	X	0.8	41%	0.6 (0.3 to 0.9)	No
Nilsson U, et al 2009 ⁷¹		Short STAI	10	14.7	✓	14.3	✓	0.4	2%	0.5 (0.5 to 0.5) *	Yes
Padam A, et al 2017 ⁷³		STAI	10	1.9	-	1.4	X	0.5	26%	0.4 (0.0 to 0.7)	No
		STAI	10	3.8	X	1.4	X	2.4	63%	2.6 (2.1 to 3.1)	No
Shabanloei R, et al 2010 ⁷⁷		STAI	10	9.7	X	5.8	X	3.9	40%	3.6 (2.9 to 4.3)	No
Sobana R, et al 2015 ⁸²	Short STAI	10	6.1	X	0.1	X	6.1	99%	2.0 (2.6 to 1.3)	No	

Diette GB, et al 2003 ⁵⁰	Audio-visual technology interventions	STAI	10	13.5	✓	12.0	✓	1.5	11%	-1.8 (-1.3 to -2.4)	No
Drahota A, et al 2008 ⁵¹		STAI	10	13.5	✓	12.0	✓	1.5	11%	0.2 (-0.3 to 0.6) *	No
Fang AS, et al 2016 ⁵³		STAI	10	6.1	X	0.1	X	6.1	99%	2.0 (2.5 to 1.5)	No
Hudson BF, et al 2015 ⁶⁰ (DVD)		STAI	10	2.3	X	-2.3	X	4.6	199%	3.3 (3.8 to 2.8)	No
Xiaolian J, et al 2015 ⁸⁸		STAI	10	5.0	X	4.1	X	0.8	17%	0.7 (0.3 to 1.0)	No
		STAI	10	2.5	X	-2.3	X	4.7	7%	3.3 (3.8 to 2.8)	No
Hızlı F, et al 2015 ⁵⁷	Psychological interventions	BAI	8.8	3.0	X	-1.9	X	4.8	38%	0.9 (0.6 to 1.2) *	No
		HAS	8.2	4.6	X	-2.8	X	7.4	40%	0.9 (0.6 to 1.3) *	No
Snow A, et al 2012 ⁸¹		VAS-A (0-100mm)	46	22.0	X	13.0	X	9.0	41%	0.7 (1.2 to 0.3)	No
Kwekkeboom KL, et al 2003 ⁶⁴		STAI	10	6.3	X	7.0	X	0.7	11%	-1.2 (-0.5 to -1.9)	No
Hudson BF, et al 2015 ⁶⁰		STAI	10	2.5	X	-2.3	X	4.7	193%	3.3 (3.8 to 2.8)	No
Heidaria F, et al 2017 ⁵⁶	Physical interventions	STAI	10	4.4	X	1.5	X	2.9	66%	1.0 (0.9 to 1.0) *	No
Rosen J, et al 2013 ⁷⁵		STAI	10	6.5	X	8.6	X	-2.1	-32%	-0.2 (0.5 to -0.9)	No
		STAI	10	12.1	✓	9.5	X	2.6	21%	0.2 (0.9 to -0.4)	No
Hudson BF, et al 2015 ⁶⁰		STAI	10	3.0	X	-2.3	X	5.3	176%	2.4 (2.8 to 1.9)	No
Ahlander BM, et al 2018 ⁴³		Other: Education/information	STAI	10	6.5	X	1.1	X	5.4	83%	-1.0 (-1.4 to -0.6)
Kola S, et al 2013 ⁶³			10	-4.0	X	4.5	X	-8.5	212%	1.0 (-1.2 to -0.7)*	No
	STAI		10	6.2	X	3.5	X	2.7	44%	0.4 (-0.3 to 1.1)*	No
			10	-6.2	X	4.5	X	-10.7	-173%	-1.0 (-1.3 to -0.5)*	No

			10	4.1	X	3.9	X	0.1	3%	0.0 (-0.3 to 0.4) *	No
Wu KL, et al 2014 ⁸⁷		STAI	10	16.3	✓	10.2	✓	6.2	38%	0.9 (0.6 to 1.2) *	Yes
				13.5	✓	10.2	✓	3.3	25%	0.5 (0.0 to 1.0) *	Yes
Eslami J, et al 2018 ⁵²		STAI	10	12.8	✓	-1.0	X	13.8	92%	5.9 (4.7 to 7.1)	Yes
		STAI	10	13.7	✓	-1.0	X	14.7	93%	9.0 (4.7 to 10.7)	Yes
Hu PH, et al 2010 ⁵⁹	Other: Aromatherapy	STAI	10	11.0	✓	7.1	X	3.9	35%	0.3 (-2.6 to 2.1) *	No
Trambert T, et al 2014 ⁸³		STAI	10	14.2	✓	2.9	X	11.3	79%	0.5 (-2.4 to 3.3) *	Yes
		STAI	10	6.5	X	2.9	X	3.6	55%	0.2 (-2.8 to 3.1) *	No

Abbreviations: STAI, State Trait Anxiety Inventory. BAI, Becks Anxiety Inventory. HAS, Hamilton Anxiety Scale. VAS-A, Anxiety Visual Analogue Scale.

Effect size is calculated using Cohen's D except where * indicates r-Stat has been used for non-parametric data. Comfort intervention was considered to demonstrate clinical significance when the effect size exceeded >0.4, mean differences were greater than the MID and RoB was acceptable.